

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
11 November 2004 (11.11.2004)

PCT

(10) International Publication Number
WO 2004/096226 A1

(51) International Patent Classification⁷: **A61K 31/517**, C07D 403/12, 403/14, A61P 35/00 (74) Agent: **ASTRAZENECA**; Global Intellectual Property, S-151 85 Sodertälje (SE).

(21) International Application Number:
PCT/GB2004/001799

(22) International Filing Date: 27 April 2004 (27.04.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0309850.6 30 April 2003 (30.04.2003) GB

(71) Applicant (for all designated States except MG, US): **ASTRAZENECA AB** [SE/SE]; S-151 85 Sodertälje (SE).

(71) Applicant (for MG only): **ASTRAZENECA UK LIMITED** [GB/GB]; 15 Stanhope Gate, London, Greater London W1K 1LN (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **HENNEQUIN, Laurent, Francois, Andre** [FR/GB]; AstraZeneca R & D Alderley, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). **PLOWRIGHT, Allyn** [GB/GB]; AstraZeneca R & D Alderley, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **QUINAZOLINE DERIVATIVES AND THEIR USE IN THE TREATMENT OF CANCER**



(57) Abstract: The invention concerns quinazoline derivatives of the formula: (I); wherein X¹, Q¹, Z, R¹, R², Y, a and m are as defined in the description, which are erbB tyrosine kinase inhibitors, particularly EGFR tyrosine kinase inhibitors. Also claimed are processes for their preparation; pharmaceutical compositions containing them; and their use as therapeutic agents in the treatment of erbB tyrosine kinase mediated diseases such as cancer.

QUINAZOLINE DERIVATIVES AND THEIR USE IN THE TREATMENT OF CANCER

The invention concerns certain novel quinazoline derivatives, or

pharmaceutically-acceptable salts thereof, which possess anti-tumour activity and are

5 accordingly useful in methods of treatment of the human or animal body. The invention also concerns processes for the manufacture of said quinazoline derivatives, to pharmaceutical compositions containing them and to their use in therapeutic methods, for example in the manufacture of medicaments for use in the prevention or treatment of solid tumour disease in a warm-blooded animal such as man.

10 Many of the current treatment regimes for diseases resulting from the abnormal regulation of cellular proliferation such as psoriasis and cancer, utilise compounds that inhibit DNA synthesis and cellular proliferation. To date, compounds used in such treatments are generally toxic to cells however their enhanced effects on rapidly dividing cells such as tumour cells can be beneficial. Alternative approaches to these cytotoxic anti-tumour agents
15 are currently being developed, for example selective inhibitors of cell signalling pathways. These types of inhibitors are likely to have the potential to display an enhanced selectivity of action against tumour cells and so are likely to reduce the probability of the therapy possessing unwanted side effects.

Eukaryotic cells are continually responding to many diverse extracellular signals that
20 enable communication between cells within an organism. These signals regulate a wide variety of physical responses in the cell including proliferation, differentiation, apoptosis and motility. The extracellular signals take the form of a diverse variety of soluble factors including growth factors as well as paracrine and endocrine factors. By binding to specific transmembrane receptors, these ligands integrate the extracellular signal to the intracellular
25 signalling pathways, therefore transducing the signal across the plasma membrane and allowing the individual cell to respond to its extracellular signals. Many of these signal transduction processes utilise the reversible process of the phosphorylation of proteins that are involved in the promotion of these diverse cellular responses. The phosphorylation status of target proteins is regulated by specific kinases and phosphatases that are responsible for the
30 regulation of about one third of all proteins encoded by the mammalian genome. As phosphorylation is such an important regulatory mechanism in the signal transduction process, it is therefore not surprising that aberrations in these intracellular pathways result in abnormal

cell growth and differentiation and so promote cellular transformation (reviewed in Cohen *et al*, Curr Opin Chem Biol, 1999, 3, 459-465).

It has been widely shown that a number of these tyrosine kinases are mutated to constitutively active forms and/or when over-expressed result in the transformation of a variety of human cells. These mutated and over-expressed forms of the kinase are present in a large proportion of human tumours (reviewed in Kolibaba *et al*, Biochimica et Biophysica Acta, 1997, 133, F217-F248). As tyrosine kinases play fundamental roles in the proliferation and differentiation of a variety of tissues, much focus has centred on these enzymes in the development of novel anti-cancer therapies. This family of enzymes is divided into two groups - receptor and non-receptor tyrosine kinases e.g. EGF Receptors and the SRC family respectively. From the results of a large number of studies including the Human Genome Project, about 90 tyrosine kinase have been identified in the human genome, of this 58 are of the receptor type and 32 are of the non-receptor type. These can be compartmentalised in to 20 receptor tyrosine kinase and 10 non-receptor tyrosine kinase sub-families (Robinson *et al*, Oncogene, 2000, 19, 5548-5557).

The receptor tyrosine kinases are of particular importance in the transmission of mitogenic signals that initiate cellular replication. These large glycoproteins, which span the plasma membrane of the cell possess an extracellular binding domain for their specific ligands (such as Epidermal Growth Factor (EGF) for the EGF Receptor). Binding of ligand results in the activation of the receptor's kinase enzymatic activity that is encoded by the intracellular portion of the receptor. This activity phosphorylates key tyrosine amino acids in target proteins, resulting in the transduction of proliferative signals across the plasma membrane of the cell.

It is known that the erbB family of receptor tyrosine kinases, which include EGFR, erbB2, erbB3 and erbB4, are frequently involved in driving the proliferation and survival of tumour cells (reviewed in Olayioye *et al*, EMBO J, 2000, 19, 3159). One mechanism in which this can be accomplished is by over expression of the receptor at the protein level, generally as a result of gene amplification. This has been observed in many common human cancers (reviewed in Klapper *et al*, Adv. Cancer Res, 2000, 77, 25) such as breast cancer (Sainsbury *et al*, Brit. J. Cancer, 1988, 58, 458; Guerin *et al*, Oncogene Res, 1988, 3, 21; Slamon *et al*, Science, 1989, 244, 707; Klijn *et al*, Breast Cancer Res. Treat., 1994, 29, 73 and reviewed in Salomon *et al*, Crit. Rev. Oncol. Hematol, 1995, 19, 183), non-small cell lung cancers (NSCLCs) including adenocarcinomas (Cerny *et al*, Brit. J. Cancer, 1986, 54,

265; Reubi *et al.*, Int. J. Cancer, 1990, 45, 269; Rusch *et al.*, Cancer Research, 1993, 53, 2379; Brabender *et al.*, Clin. Cancer Res., 2001, 7, 1850) as well as other cancers of the lung (Hendler *et al.*, Cancer Cells, 1989, 7, 347; Ohsaki *et al.*, Oncol. Rep., 2000, 7, 603), bladder cancer (Neal *et al.*, Lancet, 1985, 366; Chow *et al.*, Clin. Cancer Res., 2001, 7, 1957, Zhau *et al.*, Mol Carcinog., 3, 254), oesophageal cancer (Mukaida *et al.*, Cancer, 1991, 68, 142),
5 gastrointestinal cancer such as colon, rectal or stomach cancer (Bolen *et al.*, Oncogene Res., 1987, 1, 149; Kapitanovic *et al.*, Gastroenterology, 2000, 112, 1103; Ross *et al.*, Cancer Invest., 2001, 19, 554), cancer of the prostate (Visakorpi *et al.*, Histochem. J., 1992, 24, 481; Kumar *et al.*, 2000, 32, 73; Scher *et al.*, J. Natl. Cancer Inst., 2000, 92, 1866), leukaemia
10 (Konaka *et al.*, Cell, 1984, 37, 1035, Martin-Subero *et al.*, Cancer Genet Cytogenet., 2001, 127, 174), ovarian (Hellstrom *et al.*, Cancer Res., 2001, 61, 2420), head and neck (Shiga *et al.*, Head Neck, 2000, 22, 599) or pancreatic cancer (Ovotny *et al.*, Neoplasma, 2001, 48, 188).
As more human tumour tissues are tested for expression of the erbB family of receptor tyrosine kinases it is expected that their widespread prevalence and importance will be further
15 enhanced in the future.

As a consequence of the mis-regulation of one or more of these receptors, it is widely believed that many tumours become clinically more aggressive and so correlate with a poorer prognosis for the patient (Brabender *et al.*, Clin. Cancer Res., 2001, 7, 1850; Ross *et al.*, Cancer Investigation, 2001, 19, 554, Yu *et al.*, Bioessays, 2000, 22.7, 673). In addition to these
20 clinical findings, a wealth of pre-clinical information suggests that the erbB family of receptor tyrosine kinases are involved in cellular transformation. This includes the observations that many tumour cell lines over express one or more of the erbB receptors and that EGFR or erbB2 when transfected into non-tumour cells have the ability to transform these cells. This tumourigenic potential has been further verified as transgenic mice that over express erbB2
25 spontaneously develop tumours in the mammary gland. In addition to this, a number of pre-clinical studies have demonstrated that anti-proliferative effects can be induced by knocking out one or more erbB activities by small molecule inhibitors, dominant negatives or inhibitory antibodies (reviewed in Mendelsohn *et al.*, Oncogene, 2000, 19, 6550). Thus it has been recognised that inhibitors of these receptor tyrosine kinases should be of value as a
30 selective inhibitor of the proliferation of mammalian cancer cells (Yaish *et al.*, Science, 1988, 242, 933, Kolibaba *et al.*, Biochimica et Biophysica Acta, 1997, 133, F217-F248; Al-Obeidi *et al.*, 2000, Oncogene, 19, 5690-5701; Mendelsohn *et al.*, 2000, Oncogene, 19, 6550-6565). In addition to this pre-clinical data, findings using inhibitory antibodies against EGFR and erbB2

(c-225 and trastuzumab respectively) have proven to be beneficial in the clinic for the treatment of selected solid tumours (reviewed in Mendelsohn *et al*, 2000, Oncogene, 19, 6550-6565).

Amplification and/or activity of members of the erbB type receptor tyrosine kinases have been detected and so have been implicated to play a role in a number of non-malignant proliferative disorders such as psoriasis (Ben-Bassat, Curr. Pharm. Des., 2000, 6, 933; Elder *et al.*, Science, 1989, 243, 811), benign prostatic hyperplasia (BPH) (Kumar *et al.*, Int. Urol. Nephrol., 2000, 32, 73), atherosclerosis and restenosis (Bokemeyer *et al.*, Kidney Int., 2000, 58, 549). It is therefore expected that inhibitors of erbB type receptor tyrosine kinases will be useful in the treatment of these and other non-malignant disorders of excessive cellular proliferation. International Patent Applications WO 96/33977, WO 96/33978, WO 96/33979, WO 96/33980 and WO 96/33981 disclose that certain quinazoline derivatives which bear an anilino substituent at the 4-position possess receptor tyrosine kinase inhibitory activity.

A review of the structure activity relationship of various quinazoline derivatives is disclosed by G. W. Rewcastle *et al* (J. Med. Chem. 1995, 38, 3428-3487), including a number of 5-substituted compounds. However, such 5-substituted compounds are stated to have low in-vitro activity as EGFR tyrosine kinase inhibitors compared to quinazolines substituted at the 6- and 7- positions.

WO 96/09294 discloses 4-anilinoquinazoline derivatives, including 5-chloro and 5-methoxy substituted quinazoline derivatives as protein tyrosine kinase inhibitors.

WO96/15118 discloses certain 4-anilinoquinazoline derivatives that are substituted on the aniline by certain aryl or heteroaryl groups. The compounds are stated to be Class 1 receptor tyrosine kinase inhibitors. International Patent Application WO 97/03069 also discloses certain 4-substituted quinazoline derivatives and states that the compounds are erbB2 tyrosine kinase inhibitors.

WO97/30034 describes 4-anilinoquinazoline derivatives that are substituted on the aniline by certain aryl or heteroaryl groups and which are also substituted at the 6-position on the quinazoline by certain aryl or heteroaryl groups. These compounds are also Class I receptor tyrosine kinase inhibitors.

WO01/21596 discloses the use of 4-anilinoquinazoline derivatives as aurora 2 kinase inhibitors.

There is no disclosure in WO96/15118, WO 97/03069, WO97/30034 or WO01/21596 of any compounds substituted at the 5-position on the quinazoline ring.

International Patent Application WO01/94341 discloses that certain quinazoline derivatives which carry a 5-substituent are inhibitors of the Src family of non-receptor tyrosine
5 kinases, such as c-Src, c-Yes and c-Fyn.


None of the prior art discloses 4-anilinoquinazolines which are substituted at the 5-position by a pyrrolidinylmethoxy substituent, which is substituted on a ring nitrogen atom by an optionally substituted alkanoyl group.

Surprisingly we have now found that certain 5-substituted quinazoline derivatives
10 possess potent anti-tumour activity. In particular the compounds of the present invention are possess potent anti-tumour activity. Without wishing to imply that the compounds disclosed in the present invention possess pharmacological activity only by virtue of an effect on a single biological process, it is believed that the compounds provide an anti-tumour effect by way of inhibition of one or more of the erbB family of receptor tyrosine kinases that are
15 involved in the signal transduction steps which lead to the proliferation of tumour cells. In particular, it is believed that the compounds of the present invention provide an anti-tumour effect by way of inhibition of EGFR and/or erbB2 receptor tyrosine kinases.

Generally the compounds of the present invention possess potent inhibitory activity against the erbB receptor tyrosine kinase family, for example by inhibition of EGFR and/or
20 erbB2 and/or erbB4 receptor tyrosine kinases, whilst possessing less potent inhibitory activity against other kinases. Furthermore, many of the compounds of the present invention possess substantially better potency against the EGFR over that of the erbB2 tyrosine kinase. This has the advantage of allowing highly selective EGFR tyrosine kinase inhibition, even within the erbB family of receptors. The invention also includes compounds that are active against all or
25 a combination of EGFR, erbB2 and erbB4 receptor tyrosine kinases, thus potentially providing treatments for conditions mediated by one or more of these receptor tyrosine kinases.

Generally the compounds of the present invention exhibit favourable physical properties such as a high solubility whilst retaining high antiproliferative activity.

According to a first aspect of the invention there is provided a quinazoline derivative
30 of the formula I:



I

wherein:

- X^1 is $C(R^3)_2$, wherein each R^3 , which may be the same or different, is selected from
- 5 hydrogen and (1-4C)alkyl;
- Q^1 is pyrrolidinyl;
- and wherein Q^1 is linked to the group X^1-O by a ring carbon atom,
- and wherein Q^1 is substituted at the 1-position by the group of the formula $ZC(O)$;
- Z is selected from (1-4C)alkyl, halogeno-(1-4C)alkyl, hydroxy-(1-4C)alkyl, (1-
- 10 4C)alkoxy-(1-4C)alkyl, amino-(1-4C)alkyl, (1-4C)alkylamino-(1-4C)alkyl and
- di-[(1-4C)alkyl]amino-(1-4C)alkyl,
- or Z is Q^2 , wherein Q^2 is heterocyclyl or heterocyclyl-(1-4C)alkyl,
- and wherein any CH_2 or CH_3 group within a Z group, other than a CH_2 group within a
- heterocyclyl ring, optionally bears on each said CH_2 or CH_3 group one or more halogeno or
- 15 (1-6C)alkyl substituents or a substituent selected from hydroxy, cyano, amino, carboxy,
- carbamoyl, formyl, mercapto, sulfamoyl, (2-6C)alkenyl, (2-6C)alkynyl, (1-6C)alkoxy,
- (1-6C)alkylthio, (1-6C)alkylsulfinyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino,
- di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl,
- N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino,
- 20 N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulfamoyl,
- N,N-di-[(1-6C)alkyl]sulfamoyl, (1-6C)alkanesulfonylamino and
- N-(1-6C)alkyl-(1-6C)alkanesulfonylamino,
- and wherein any heterocyclyl group within Q^1 or Z optionally bears one or more (for
- example 1, 2 or 3) substituents, which may be the same or different selected from halogeno,
- 25 trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, sulfamoyl, formyl,
- mercapto, (1-6C)alkyl, (2-6C)alkenyl, (2-6C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy,
- (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulfinyl, (1-6C)alkylsulfonyl,
- (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl,

N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulfamoyl, N,N-di-[(1-6C)alkyl]sulfamoyl, (1-6C)alkanesulfonylamino, and N-(1-6C)alkyl-(1-6C)alkanesulfonylamino, or from a group of the formula:




wherein X^2 is a direct bond or is selected from O, CO and $N(R^5)$, wherein R^5 is hydrogen or (1-6C)alkyl, and R^4 is halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl, carboxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl, cyano-(1-6C)alkyl, amino-(1-6C)alkyl, N-(1-6C)alkylamino-(1-6C)alkyl, N,N-di-[(1-6C)alkyl]amino-(1-6C)alkyl, (2-6C)alkanoylamino-(1-6C)alkyl, (1-6C)alkoxycarbonylamino-(1-6C)alkyl, carbamoyl-(1-6C)alkyl, N-(1-6C)alkylcarbamoyl-(1-6C)alkyl, N,N-di-[(1-6C)alkyl]carbamoyl-(1-6C)alkyl, (2-6C)alkanoyl-(1-6C)alkyl, (2-6C)alkanoyloxy-(1-6C)alkyl and (1-6C)alkoxycarbonyl-(1-6C)alkyl,

and wherein any heterocyclyl group within Q^1 or Z optionally bears 1 or 2 oxo or thioxo substituents;

m is 0, 1 or 2 and R^1 when present is located at the 6- and/or 7- position;


each R^1 group, which may be the same or different, is selected from hydroxy, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, or from a group of the formula:



20 wherein X^3 is a direct bond or is O, and Q^3 is (3-7C)cycloalkyl, (3-7C)cycloalkyl-(1-6C)alkyl, (3-7C)cycloalkenyl, (3-7C)cycloalkenyl-(1-6C)alkyl, heterocyclyl or heterocyclyl-(1-6C)alkyl, or $(R^1)_m$ is (1-3C)alkylenedioxy,


and wherein adjacent carbon atoms in any (2-6C)alkylene chain within a R^1 substituent are optionally separated by the insertion into the chain of a group selected from O, S, SO, SO₂, $N(R^6)$, CO, CH(OR⁶), CON(R⁶), $N(R^6)$ CO, SO₂N(R⁶), $N(R^6)$ SO₂, CH=CH and C≡C wherein R^6 is hydrogen or (1-6C)alkyl,

and wherein any CH₂=CH- or HC≡C- group within a R^1 substituent optionally bears at the terminal CH₂= or HC≡ position a substituent selected from halogeno, carboxy, carbamoyl, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl and di-[(1-6C)alkyl]amino-(1-6C)alkyl or from a group of the formula :




wherein X^4 is a direct bond or is selected from CO and $N(R^7)CO$, wherein R^7 is hydrogen or (1-6C)alkyl, and Q^4 is heterocyclyl or heterocyclyl-(1-6C)alkyl,

and wherein any CH_2 or CH_3 group within a R^1 substituent, other than a CH_2 group within a heterocyclyl ring, optionally bears on each said CH_2 or CH_3 group one or more
 5 halogeno or (1-6C)alkyl substituents or a substituent selected from hydroxy, cyano, amino, carboxy, carbamoyl, sulfamoyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulfinyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino,
 10 N-(1-6C)alkylsulfamoyl, N,N-di-[(1-6C)alkyl]sulfamoyl, (1-6C)alkanesulfonylamino and N-(1-6C)alkyl-(1-6C)alkanesulfonylamino, or from a group of the formula:



wherein X^5 is a direct bond or is selected from O, S, SO, SO_2 , $N(R^8)$, CO, $CH(OR^8)$, $CON(R^8)$, $N(R^8)CO$, $SO_2N(R^8)$, $N(R^8)SO_2$, $C(R^8)_2O$, $C(R^8)_2S$ and $C(R^8)_2N(R^8)$, wherein R^8 is
 15 hydrogen or (1-6C)alkyl, and Q^5 is (3-7C)cycloalkyl, (3-7C)cycloalkyl-(1-6C)alkyl, (3-7C)cycloalkenyl, (3-7C)cycloalkenyl-(1-6C)alkyl, heterocyclyl or heterocyclyl-(1-6C)alkyl,

and wherein any heterocyclyl group within a substituent on R^1 optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from halogeno, trifluoromethyl,
 20 cyano, nitro, hydroxy, amino, carboxy, carbamoyl, formyl, mercapto, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulfinyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino,
 25 N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulfamoyl, N,N-di-[(1-6C)alkyl]sulfamoyl, (1-6C)alkanesulfonylamino, and N-(1-6C)alkyl-(1-6C)alkanesulfonylamino, or from a group of the formula:



wherein X^6 is a direct bond or is selected from O, $N(R^{10})$ and C(O), wherein R^{10} is hydrogen
 30 or (1-6C)alkyl, and R^9 is halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl, carboxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl, cyano-(1-6C)alkyl, amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl, di-[(1-6C)alkyl]amino-(1-6C)alkyl, (2-6C)alkanoylamino-(1-6C)alkyl, (1-6C)alkoxycarbonylamino-(1-6C)alkyl,

carbamoyl-(1-6C)alkyl, N-(1-6C)alkylcarbamoyl-(1-6C)alkyl,
N,N-di-[(1-6C)alkyl]carbamoyl-(1-6C)alkyl, (2-6C)alkanoyl-(1-6C)alkyl or
 (1-6C)alkoxycarbonyl-(1-6C)alkyl,

and wherein any heterocyclyl group within a substituent on R¹ optionally bears 1 or 2
 5 oxo or thioxo substituents;

Y is selected from halogeno, cyano, trifluoromethyl, (1-6C)alkyl, (2-6C)alkenyl,
 (2-6C)alkynyl and (1-6C)alkoxy;

each R², which may be the same or different, is selected from halogeno, cyano, nitro,
 hydroxy, amino, carboxy, carbamoyl, sulfamoyl, trifluoromethyl, (1-6C)alkyl, (2-8C)alkenyl,
 10 (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylthio,
 (1-6C)alkylsulfinyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino,
 (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl,
N-(1-6C)alkylsulfamoyl, N,N-di-[(1-6C)alkyl]sulfamoyl; and

a is 0, 1, 2, 3 or 4;

15 or a pharmaceutically acceptable salt thereof.

According to another aspect of the invention there is provided a quinazoline derivative
 of the formula I as hereinbefore defined wherein X¹, Q¹, R¹, R², a, m and Y are as
 hereinbefore defined; and


Z is selected from halogeno-(1-4C)alkyl, hydroxy-(1-4C)alkyl, (1-
 20 4C)alkoxy-(1-4C)alkyl, amino-(1-4C)alkyl, (1-4C)alkylamino-(1-4C)alkyl and
 di-[(1-4C)alkyl]amino-(1-4C)alkyl,

or Z is Q², wherein Q² is heterocyclyl or heterocyclyl-(1-4C)alkyl, and wherein the
 heterocyclyl in Q² contains at least 1 nitrogen heteroatom, and optionally 1 or 2 further
 heteroatoms selected from oxygen, nitrogen and sulfur,

25 and wherein any CH₂ or CH₃ group within a Z group, other than a CH₂ group within a
 heterocyclyl ring, optionally bears on each said CH₂ or CH₃ group one or more halogeno or
 (1-6C)alkyl substituents or a substituent selected from hydroxy, cyano, amino, carboxy,
 carbamoyl, formyl, mercapto, sulfamoyl, (2-6C)alkenyl, (2-6C)alkynyl, (1-6C)alkoxy,
 (1-6C)alkylthio, (1-6C)alkylsulfinyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino,
 30 di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl,
N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino,
N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulfamoyl,

N,N-di-[(1-6C)alkyl]sulfamoyl, (1-6C)alkanesulfonylamino and
N-(1-6C)alkyl-(1-6C)alkanesulfonylamino,

and wherein any heterocyclyl group within Q¹ or Z optionally bears one or more (for
 example 1, 2 or 3) substituents, which may be the same or different selected from halogeno,
 5 trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, sulfamoyl, formyl,
 mercapto, (1-6C)alkyl, (2-6C)alkenyl, (2-6C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy,
 (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulfinyl, (1-6C)alkylsulfonyl,
 (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl,
N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino,
 10 N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulfamoyl,
N,N-di-[(1-6C)alkyl]sulfamoyl, (1-6C)alkanesulfonylamino, and
N-(1-6C)alkyl-(1-6C)alkanesulfonylamino, or from a group of the formula:



wherein X² is a direct bond or is selected from O, CO and N(R⁵), wherein R⁵ is
 15 hydrogen or (1-6C)alkyl, and R⁴ is halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl, carboxy-(1-
 6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl, cyano-(1-6C)alkyl, amino-(1-6C)alkyl,
N-(1-6C)alkylamino-(1-6C)alkyl, N,N-di-[(1-6C)alkyl]amino-(1-6C)alkyl,
 (2-6C)alkanoylamino-(1-6C)alkyl, (1-6C)alkoxycarbonylamino-(1-6C)alkyl,
 carbamoyl-(1-6C)alkyl, N-(1-6C)alkylcarbamoyl-(1-6C)alkyl,
 20 N,N-di-[(1-6C)alkyl]carbamoyl-(1-6C)alkyl, (2-6C)alkanoyl-(1-6C)alkyl,
 (2-6C)alkanoyloxy-(1-6C)alkyl and (1-6C)alkoxycarbonyl-(1-6C)alkyl,

and wherein any heterocyclyl group within Q¹ or Z optionally bears 1 or 2 oxo or
 thioxo substituents;

or a pharmaceutically acceptable salt thereof.

25 In this specification the generic term "alkyl" includes both straight-chain and
 branched-chain alkyl groups such as propyl, isopropyl and tert-butyl, and (3-7C)cycloalkyl
 groups such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl. However
 references to individual alkyl groups such as "propyl" are specific for the straight-chain
 version only, references to individual branched-chain alkyl groups such as "isopropyl" are
 30 specific for the branched-chain version only and references to individual cycloalkyl groups
 such as "cyclopentyl" are specific for that 5-membered ring only. An analogous convention
 applies to other generic terms, for example (1-6C)alkoxy includes methoxy, ethoxy,
 cyclopropyloxy and cyclopentyloxy, (1-6C)alkylamino includes methylamino, ethylamino,

cyclobutylamino and cyclohexylamino, and di-[(1-6Calkyl)amino includes dimethylamino, diethylamino, N-cyclobutyl-N-methylamino and N-cyclohexyl-N-ethylamino.

It is to be understood that, insofar as certain of the compounds of formula I defined above may exist in optically active or racemic forms by virtue of one or more asymmetric
5 carbon atoms, the invention includes in its definition any such optically active or racemic form which possesses the above-mentioned activity. The synthesis of optically active forms may be carried out by standard techniques of organic chemistry well known in the art, for example by synthesis from optically active starting materials or by resolution of a racemic form. Similarly, the above-mentioned activity may be evaluated using the standard laboratory
10 techniques referred to hereinafter.

It is to be understood that the present invention includes in its definition any and all tautomeric forms of the compounds of the formula I which possess the above mentioned activity.

It is also to be understood that in so far as certain compounds of the formula I may
15 exist in solvated forms as well as unsolvated forms, for example, hydrated forms, the present invention includes any and all such solvated forms, which possess the above mentioned activity.

Suitable values for the generic radicals referred to above include those set out below.

A suitable value for any one of the 'Q' groups (Q^3 and Q^5) when it is
20 (3-7C)cycloalkyl or for the (3-7C)cycloalkyl group within a 'Q' group is, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or bicyclo[2.2.1]heptyl and a suitable value for any one of the 'Q' groups (Q^3 and Q^5) when it is (3-7C)cycloalkenyl or for the (3-7C)cycloalkenyl group within a 'Q' group is, for example, cyclobutenyl, cyclopentenyl, cyclohexenyl or cycloheptenyl.

25 A suitable value for any one of the 'Q' groups (Q^2 to Q^5) when it is heterocyclyl or for the heterocyclyl group within a 'Q' group is, for example, a non-aromatic saturated or partially saturated 3 to 10 membered monocyclic or bicyclic ring with up to five heteroatoms selected from oxygen, nitrogen and sulfur, which, unless specified otherwise, may be carbon or nitrogen linked. Examples of suitable values of "heterocyclyl" include oxiranyl, oxetanyl,
30 azetidiny, tetrahydrofuranyl, tetrahydropyranyl, oxepanyl, pyrrolinyl, pyrrolidinyl, morpholinyl, tetrahydro-1,4-thiazinyl, 1,1-dioxotetrahydro-1,4-thiazinyl, piperidinyl, homopiperidinyl, piperazinyl, homopiperazinyl, dihydropyridinyl, tetrahydropyridinyl, dihydropyrimidinyl, tetrahydropyrimidinyl, tetrahydrothienyl, tetrahydrothiopyranyl,

decahydroisoquinolinyl or decahydroquinolinyl, particularly tetrahydrofuranyl, tetrahydropyranyl, pyrrolidinyl, morpholinyl, 1,4-oxazepanyl, thiomorpholinyl 1,1-dioxotetrahydro-4H-1,4-thiazinyl, piperidinyl or piperazinyl, more particularly tetrahydrofuran-3-yl, tetrahydropyran-4-yl, tetrahydrothien-3-yl, tetrahydrothiopyran-4-yl, 5 pyrrolidin-1-yl, pyrrolidin-2-yl, pyrrolidin-3-yl, morpholino, morpholin-2-yl, morpholin-3-yl, 1,1-dioxotetrahydro-4H-1,4-thiazin-4-yl, piperidino, piperidin-4-yl, piperidin-3-yl, piperidin-2-yl, piperazin-1-yl or piperazin-2-yl. A nitrogen or sulfur atom within a heterocyclyl group may be oxidized to give the corresponding N or S oxide, for example 1,1-dioxotetrahydrothienyl, 1-oxotetrahydrothienyl, 1,1-dioxotetrahydrothiopyranyl or 10 1-oxotetrahydrothiopyranyl. A suitable value for such a group which bears 1 or 2 oxo or thioxo substituents is, for example, 2-oxopyrrolidinyl, 2-thioxopyrrolidinyl, 2-oxoimidazolidinyl, 2-thioxoimidazolidinyl, 2-oxopiperidinyl, 2,5-dioxopyrrolidinyl, 2,5-dioxoimidazolidinyl or 2,6-dioxopiperidinyl.

Particular values for Q¹ is pyrrolidin-2-yl and pyrrolidin-3-yl.

15 Particular values for Q² when it is heterocyclyl containing at least 1 nitrogen atom and optionally 1 or 2 further heteroatoms selected from oxygen, nitrogen and sulfur, include the heterocycles above containing at least 1 nitrogen atom, particularly a fully saturated 4, 5, 6 or 7-membered monocyclic or bicyclic (particularly monocyclic) heterocyclyl group containing 1 nitrogen heteroatom and optionally 1 further heteroatom selected from oxygen, nitrogen and 20 sulfur, particularly such groups that are linked by a ring nitrogen atom to the alkyl or carbonyl in formula I. When Q² is linked via a ring nitrogen this nitrogen atom is not quaternised, i.e. a neutral compound is formed. Suitable values represented by Q² include the heterocyclyl groups listed above, more particularly pyrrolidinyl, piperidinyl, piperazinyl or morpholinyl, still more particularly pyrrolidin-1-yl, piperidino, piperazin-1-yl or morpholino.

25 A suitable value for a 'Q' group when it is heterocyclyl-(1-6C)alkyl is, for example, heterocyclylmethyl, 2-heterocyclylethyl and 3-heterocyclylpropyl. The invention comprises corresponding suitable values for 'Q' groups when, for example, rather than a heterocyclyl-(1-6C)alkyl group, an (3-7C)cycloalkyl-(1-6C)alkyl or (3-7C)cycloalkenyl-(1-6C)alkyl group is present. It is to be understood that when a 'Q' group, 30 for example Q² is heterocyclyl-(1-4C)alkyl, Q² is attached to the carbonyl group in formula I via the (1-4C)alkyl portion of the heterocyclyl-(1-4C)alkyl group. For example when Q² is morpholinomethyl, the group Z-C(O)- in formula I so formed is a morpholinoacetyl group.

Suitable values for any of the 'R' groups (R^1 to R^{10}), or for various groups within an R^1 substituent, or for various groups within Q^1 , or for Z or for various groups within Z, or for Y include:-

- | | | |
|----|--|--|
| | for halogeno | fluoro, chloro, bromo and iodo; |
| 5 | for (1-6C)alkyl: | methyl, ethyl, propyl, isopropyl and <u>tert</u> -butyl; |
| | for (2-8C)alkenyl: | vinyl, isopropenyl, allyl and but-2-enyl; |
| | for (2-8C)alkynyl: | ethynyl, 2-propynyl and but-2-ynyl; |
| | for (1-6C)alkoxy: | methoxy, ethoxy, propoxy, isopropoxy and butoxy; |
| | for (2-6C)alkenyloxy: | vinylloxy and allyloxy; |
| 10 | for (2-6C)alkynyloxy: | ethynyloxy and 2-propynyloxy; |
| | for (1-6C)alkylthio: | methylthio, ethylthio and propylthio; |
| | for (1-6C)alkylsulfinyl: | methylsulfinyl and ethylsulfinyl; |
| | for (1-6C)alkylsulfonyl: | methylsulfonyl and ethylsulfonyl; |
| | for (1-6C)alkylamino: | methylamino, ethylamino, propylamino, |
| 15 | | isopropylamino and butylamino; |
| | for di-[(1-6C)alkyl]amino: | dimethylamino, diethylamino, <u>N</u> -ethyl-
<u>N</u> -methylamino and diisopropylamino; |
| | for (1-6C)alkoxycarbonyl: | methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl
and <u>tert</u> -butoxycarbonyl; |
| 20 | for <u>N</u> -(1-6C)alkylcarbamoyl: | <u>N</u> -methylcarbamoyl, <u>N</u> -ethylcarbamoyl and
<u>N</u> -propylcarbamoyl; |
| | for <u>N,N</u> -di-[(1-6C)alkyl]carbamoyl: | <u>N,N</u> -dimethylcarbamoyl, <u>N</u> -ethyl-
<u>N</u> -methylcarbamoyl and <u>N,N</u> -diethylcarbamoyl; |
| | for (2-6C)alkanoyl: | acetyl and propionyl; |
| 25 | for (2-6C)alkanoyloxy: | acetoxy and propionyloxy; |
| | for (2-6C)alkanoylamino: | acetamido and propionamido; |
| | for <u>N</u> -(1-6C)alkyl-(2-6C)alkanoylamino: | <u>N</u> -methylacetamido and <u>N</u> -methylpropionamido; |
| | for amino(2-6C)alkanoyl: | aminoacetyl and 2-aminopropionyl; |
| | for <u>N</u> -(1-6C)alkylamino(2-6C)alkanoyl: | <u>N</u> -methylaminoacetyl and 2-(<u>N</u> -
30 methylaminopropionyl; |
| | for <u>N,N</u> -di-[(1-6C)alkyl]amino(2-6C)alkanoyl: | <u>N,N</u> -di-methylaminoacetyl; |
| | for <u>N</u> -(1-6C)alkylsulfamoyl: | <u>N</u> -methylsulfamoyl and <u>N</u> -ethylsulfamoyl; |
| | for <u>N,N</u> -di-[(1-6C)alkyl]sulfamoyl: | <u>N,N</u> -dimethylsulfamoyl; |

- for (1-6C)alkanesulfonylamino: methanesulfonylamino and ethanesulfonylamino;
for N-(1-6C)alkyl-(1-6C)alkanesulfonylamino: N-methylmethanesulfonylamino and
N-methylethanesulfonylamino;
for (3-6C)alkenoylamino: acrylamido, methacrylamido and crotonamido;
5 for N-(1-6C)alkyl-(3-6C)alkenoylamino: N-methylacrylamido and N-methylcrotonamido;
for (3-6C)alkynoylamino: propiolamido;
for N-(1-6C)alkyl-(3-6C)alkynoylamino: N-methylpropiolamido;
for amino-(1-6C)alkyl: aminomethyl, 2-aminoethyl, 1-aminoethyl and
3-aminopropyl;
10 for (1-6C)alkylamino-(1-6C)alkyl: methylaminomethyl, ethylaminomethyl,
1-methylaminoethyl, 2-methylaminoethyl,
2-ethylaminoethyl and 3-methylaminopropyl;
for di-[(1-6C)alkyl]amino-(1-6C)alkyl: dimethylaminomethyl, diethylaminomethyl,
1-dimethylaminoethyl, 2-dimethylaminoethyl and
15 3-dimethylaminopropyl;
for halogeno-(1-6C)alkyl: chloromethyl, 2-chloroethyl, 1-chloroethyl and
3-chloropropyl;
for hydroxy-(1-6C)alkyl: hydroxymethyl, 2-hydroxyethyl, 1-hydroxyethyl and
3-hydroxypropyl;
20 for (1-6C)alkoxy-(1-6C)alkyl: methoxymethyl, ethoxymethyl, 1-methoxyethyl,
2-methoxyethyl, 2-ethoxyethyl and
3-methoxypropyl;
for cyano-(1-6C)alkyl: cyanomethyl, 2-cyanoethyl, 1-cyanoethyl and
3-cyanopropyl;
25 for carboxy-(1-6C)alkyl: carboxymethyl, 2-carboxyethyl, 1-carboxyethyl and
3-carboxypropyl;
for (2-6C)alkanoylamino-(1-6C)alkyl: acetamidomethyl, propionamidomethyl and
2-acetamidoethyl;
for (1-6C)alkoxycarbonyl-(1-6C)alkyl: methoxycarbonylmethyl, 2-methoxycarbonylethyl
and 2-ethoxycarbonylethyl;
30 for (2-6C)alkanoyl-(1-6C)alkyl: acetylmethyl and 2-acetylethyl;
for (2-6C)alkanoyloxy-(1-6C)alkyl: acetoxymethyl, 2-acetoxyethyl, propionyloxymethyl
and 2-propionyloxyethyl;

- for (1-6C)alkoxycarbonylamino-(1-6C)alkyl: methoxycarbonylaminomethyl,
ethoxycarbonylaminomethyl,
tert-butoxycarbonylaminomethyl and
2-methoxycarbonylaminoethyl;
- 5 for carbamoyl-(1-6C)alkyl: carbamoylmethyl, 1-carbamoylethyl,
2-carbamoylethyl and 3-carbamoylpropyl;
- for N-(1-6C)alkylcarbamoyl-(1-6C)alkyl: N-methylcarbamoylmethyl,
N-ethylcarbamoylmethyl, N-propylcarbamoylmethyl,
10 1-(N-methylcarbamoyl)ethyl,
1-(N-ethylcarbamoyl)ethyl,
2-(N-methylcarbamoyl)ethyl,
2-(N-ethylcarbamoyl)ethyl and
3-(N-methylcarbamoyl)propyl; and
- 15 for N,N-di[(1-6C)alkyl]carbamoyl-(1-6C)alkyl: N,N-dimethylcarbamoylmethyl,
N,N-diethylcarbamoylmethyl,
2-(N,N-dimethylcarbamoyl)ethyl, and
3-(N,N-dimethylcarbamoyl)propyl.

A suitable value for $(R^1)_m$ when it is a (1-3C)alkylenedioxy group is, for example,
20 methylenedioxy or ethylenedioxy and the oxygen atoms thereof occupy adjacent ring
positions.

When in this specification reference is made to a (1-4C)alkyl group it is to be
understood that such groups refer to alkyl groups containing up to 4 carbon atoms. A skilled
person will realise that representative examples of such groups are those listed above under
25 (1-6C)alkyl that contain up to 4 carbon atoms, such as methyl, ethyl, propyl and butyl.

Similarly, reference to a (1-3C)alkyl group refers to alkyl groups containing up to 3 carbon
atoms such as methyl, ethyl and propyl. A similar convention is adopted for the other groups
listed above such as (1-4C)alkoxy, (2-4C)alkenyl, (2-4C)alkynyl and (2-4C)alkanoyl.

When, as defined hereinbefore, a CH_3 group within a R^1 substituent bears a group of
30 the formula $-X^5-Q^5$ and, for example, X^5 is a $C(R^8)_2O$ linking group, it is the carbon atom, not
the oxygen atom, of the $C(R^8)_2O$ linking group which is attached to the CH_3 group and the
oxygen atom is linked to the Q^5 group. Similarly when, as defined hereinbefore, a terminal
 $CH_2=$ group within a R^1 substituent bears a group of the formula Q^4-X^4 and, for example, X^4

is $N(R^7)CO$, it is the carbonyl group of the $N(R^7)CO$ group which is attached to the $CH_2=$ group and the $N(R^7)$ group which is attached to Q^4 .

As defined hereinbefore, adjacent carbon atoms in any (2-6C)alkylene chain within a R^1 substituent may be optionally separated by the insertion into the chain of a group such as
 5 O, $CON(R^5)$, $N(R^5)$ or $C\equiv C$. For example, insertion of a $C\equiv C$ group into the ethylene chain within a 2-morpholinoethoxy group gives rise to a 4-morpholinobut-2-ynyloxy group and, for example, insertion of a CONH group into the ethylene chain within a 3-methoxypropoxy group gives rise to, for example, a 2-(2-methoxyacetamido)ethoxy group. It is to be understood that the term (2-6C)alkylene chain refers to any CH_2CH_2 group within R^1 and
 10 includes, for example alkylene chains within a (1-6C)alkyl, (1-6C)alkoxy, (2-8C)alkenyl, (2-8C)alkenyloxy, (2-8C)alkynyl and (2-8C)alkynyloxy group. For example the insertion of a $N(CH_3)$ group between the third and fourth carbon atoms in a hex-5-enyloxy group in R^1 gives rise to a 3-(N-methyl-N-allylamino)propoxy group.

When, as defined hereinbefore, any $CH_2=CH-$ or $HC\equiv C-$ group within a R^1 substituent
 15 optionally bears at the terminal $CH_2=$ or $HC\equiv$ position a substituent such as a group of the formula Q^4-X^4 - wherein X^4 is, for example, $NHCO$ and Q^4 is a heterocyclyl-(1-6C)alkyl group, suitable R^1 substituents so formed include, for example, \underline{N} -[heterocyclyl-(1-6C)alkyl]carbamoylvinyl groups such as \underline{N} -(2-pyrrolidin-1-ylethyl)carbamoylvinyl or
N-[heterocyclyl-(1-6C)alkyl]carbamoylethynyl groups such as \underline{N} -(2-pyrrolidin-
 20 1-ylethyl)carbamoylethynyl.

When, as defined hereinbefore, any CH_2 or CH_3 group within a R^1 substituent optionally bears on each said CH_2 or CH_3 group one or more halogeno or (1-6C)alkyl substituents, there are suitably 1 or 2 halogeno or (1-6C)alkyl substituents present on each said CH_2 group and there are suitably 1, 2 or 3 such substituents present on each said CH_3 group.

25 When, as defined hereinbefore, any CH_2 or CH_3 group within a R^1 substituent optionally bears on each said CH_2 or CH_3 group a substituent as defined hereinbefore, suitable R^1 substituents so formed include, for example, hydroxy-substituted heterocyclyl-(1-6C)alkoxy groups such as 2-hydroxy-3-piperidinopropoxy and 2-hydroxy-3-morpholinopropoxy, hydroxy-substituted amino-(2-6C)alkoxy groups such as 3-amino-
 30 2-hydroxypropoxy, hydroxy-substituted (1-6C)alkylamino-(2-6C)alkoxy groups such as 2-hydroxy-3-methylaminopropoxy, hydroxy-substituted di-[(1-6C)alkyl]amino-(2-6C)alkoxy groups such as 3-dimethylamino-2-hydroxypropoxy, hydroxy-substituted (1-6C)alkoxy groups

such as 2-hydroxyethoxy, (1-6C)alkoxy-substituted (1-6C)alkoxy groups such as 2-methoxyethoxy and 3-ethoxypropoxy, (1-6C)alkylsulfonyl-substituted (1-6C)alkoxy groups such as 2-methylsulfonylethoxy and heterocyclyl-substituted (1-6C)alkylamino-(1-6C)alkoxy groups such as 2-(2-morpholinoethylamino)ethoxy, 2-(2-piperazin-1-ylethylamino)ethoxy and
 5 2-(3-morpholinopropylamino)ethoxy.


Similar considerations apply to the attachments and substitutions within the Z, Q¹ and X¹ groups.

It is to be understood that the 2- and 8- positions on the quinazoline ring are unsubstituted.

10 A suitable pharmaceutically-acceptable salt of a compound of the formula I is, for example, an acid-addition salt of a compound of the formula I, for example an acid-addition salt with an inorganic or organic acid such as hydrochloric, hydrobromic, sulfuric, trifluoroacetic, citric or maleic acid; or, for example, a salt of a compound of the formula I which is sufficiently acidic, for example an alkali or alkaline earth metal salt such as
 15 calcium or magnesium salt, or an ammonium salt, or a salt with an organic base such as methylamine, dimethylamine, trimethylamine, piperidine, morpholine or tris-(2-hydroxyethyl)amine.

Particular novel compounds of the invention include, for example, quinazoline derivatives of the formula I, or pharmaceutically-acceptable salts thereof, wherein, unless
 20 otherwise stated, each of R¹, R², Y, Z, Q¹, X¹, m and a has any of the meanings defined hereinbefore or in paragraphs (a) to (zzz) hereinafter:-

(a) each R¹ group, which may be the same or different, is selected from hydroxy, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, or from a group of the formula :




25 wherein X³ is a direct bond or is O, and Q³ is (3-7C)cycloalkyl, (3-7C)cycloalkyl-(1-6C)alkyl, heterocyclyl or heterocyclyl-(1-6C)alkyl,

and wherein adjacent carbon atoms in any (2-6C)alkylene chain within a R¹ substituent are optionally separated by the insertion into the chain of a group selected from O, N(R⁶), CON(R⁶), N(R⁶)CO, CH=CH and C≡C wherein R⁶ is hydrogen or (1-6C)alkyl,


30 and wherein any CH₂=CH- or HC≡C- group within a R¹ substituent optionally bears at the terminal CH₂= or HC≡ position a substituent selected from carbamoyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, amino-(1-6C)alkyl,

(1-6C)alkylamino-(1-6C)alkyl and di-[(1-6C)alkyl]amino-(1-6C)alkyl or from a group of the formula :



wherein X^4 is a direct bond or is selected from CO and $N(R^7)CO$, wherein R^7 is hydrogen or
 5 (1-6C)alkyl, and Q^4 is heterocyclyl or heterocyclyl-(1-6C)alkyl,

and wherein any CH_2 or CH_3 group within a R^1 substituent, other than a CH_2 group within a heterocyclyl ring, optionally bears on each said CH_2 or CH_3 group one or more halogeno or (1-6C)alkyl substituents or a substituent selected from hydroxy, amino, cyano, carbamoyl, (1-6C)alkoxy, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, N-(1-6C)alkylcarbamoyl
 10 and N,N-di-[(1-6C)alkyl]carbamoyl, or from a group of the formula :



wherein X^5 is a direct bond or is selected from O, $N(R^8)$, $CON(R^8)$, $N(R^8)CO$ and $C(R^8)_2O$, wherein R^8 is hydrogen or (1-6C)alkyl, and Q^5 is heterocyclyl or heterocyclyl-(1-6C)alkyl,

and wherein any heterocyclyl group within a substituent on R^1 optionally bears 1, 2 or
 15 3 substituents, which may be the same or different, selected from halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carbamoyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, or from a group of the formula:



wherein X^6 is a direct bond or is selected from O and $N(R^{10})$, wherein R^{10} is hydrogen or (1-6C)alkyl, and R^9 is halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl, cyano-(1-6C)alkyl, amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl, di-[(1-6C)alkyl]amino-(1-6C)alkyl, carbamoyl-(1-6C)alkyl,

25 N-(1-6C)alkylcarbamoyl-(1-6C)alkyl and N,N-di-[(1-6C)alkyl]carbamoyl-(1-6C)alkyl,

and wherein any heterocyclyl group within a substituent on R^1 optionally bears 1 or 2 oxo substituents;

(b) each R^1 group, which may be the same or different, is selected from hydroxy, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, or from a group of the formula :




wherein X^3 is a direct bond or is O, and Q^3 is heterocyclyl or heterocyclyl-(1-6C)alkyl,

and wherein adjacent carbon atoms in any (2-6C)alkylene chain within a R^1 substituent are optionally separated by the insertion into the chain of a group selected from O, $N(R^6)$, $CON(R^6)$, $N(R^6)CO$, $CH=CH$ and $C\equiv C$ wherein R^6 is hydrogen or (1-6C)alkyl,

and wherein any $CH_2=CH-$ or $HC\equiv C-$ group within a R^1 substituent optionally bears


- 5 at the terminal $CH_2=$ or $HC\equiv$ position a substituent selected from carbamoyl, \underline{N} -(1-6C)alkylcarbamoyl, $\underline{N,N}$ -di-[(1-6C)alkyl]carbamoyl, amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl and di-[(1-6C)alkyl]amino-(1-6C)alkyl

- and wherein any CH_2 or CH_3 group within a R^1 substituent, other than a CH_2 group within a heterocyclyl ring, optionally bears on each said CH_2 or CH_3 group one or more
10 halogeno or (1-6C)alkyl substituents or a substituent selected from hydroxy, amino, cyano, carbamoyl, (1-6C)alkoxy, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, \underline{N} -(1-6C)alkylcarbamoyl and $\underline{N,N}$ -di-[(1-6C)alkyl]carbamoyl, or from a group of the formula :



- wherein X^5 is a direct bond or is selected from O, $N(R^8)$, $CON(R^8)$, $N(R^8)CO$ and $C(R^8)_2O$,
15 wherein R^8 is hydrogen or (1-6C)alkyl, and Q^5 is heterocyclyl or heterocyclyl-(1-6C)alkyl,
and wherein any heterocyclyl group within a substituent on R^1 optionally bears 1, 2 or

- 3 substituents, which may be the same or different, selected from halogeno, trifluoromethyl, hydroxy, amino, carbamoyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, \underline{N} -(1-6C)alkylcarbamoyl,
20 $\underline{N,N}$ -di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, or from a group of the formula:



- wherein X^6 is a direct bond or is selected from O and $N(R^{10})$, wherein R^{10} is hydrogen or (1-6C)alkyl, and R^9 is halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl, cyano-(1-6C)alkyl, amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl and
25 di-[(1-6C)alkyl]amino-(1-6C)alkyl,

and wherein any heterocyclyl group within a substituent on R^1 optionally bears 1 or 2 oxo substituents;


- (c) each R^1 group, which may be the same or different, is selected from hydroxy, (1-6C)alkoxy, (2-6C)alkenyloxy and (2-6C)alkynyloxy,

- 30 and wherein adjacent carbon atoms in any (2-6C)alkylene chain within a R^1 substituent are optionally separated by the insertion into the chain of a group selected from O, $N(R^6)$, $CON(R^6)$, $N(R^6)CO$, $CH=CH$ and $C\equiv C$ wherein R^6 is hydrogen or (1-6C)alkyl,

and wherein any CH₂ or CH₃ group within a R¹ substituent, other than a CH₂ group within a heterocyclyl ring, optionally bears on each said CH₂ or CH₃ group one or more halogeno or (1-6C)alkyl substituents or a substituent selected from hydroxy, amino, cyano, carbamoyl, (1-6C)alkoxy, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, N-(1-6C)alkylcarbamoyl
 5 and N,N-di-[(1-6C)alkyl]carbamoyl;

- (d) each R¹ which may be the same or different, are selected from hydroxy, methoxy, ethoxy, propoxy, isopropoxy, 2-hydroxyethoxy, 2-fluoroethoxy, cyclopropylmethoxy, 2-cyclopropylethoxy, vinyloxy, allyloxy, ethynyloxy, 2-propynyloxy, tetrahydrofurfuryloxy, 2-(tetrahydrofuran-2-yl)ethoxy, 3-(tetrahydrofuran-2-yl)propoxy,
 10 2-(tetrahydrofuran-3-yl)ethoxy, 3-(tetrahydrofuran-3-yl)propoxy, 2-pyrrolidin-1-ylethoxy, 3-pyrrolidin-1-ylpropoxy, pyrrolidin-3-yloxy, pyrrolidin-2-ylmethoxy, 2-pyrrolidin-2-ylethoxy, 3-pyrrolidin-2-ylpropoxy, 2-morpholinoethoxy, 3-morpholinopropoxy, 2-(1,1-dioxotetrahydro-4H-1,4-thiazin-4-yl)ethoxy, 3-(1,1-dioxotetrahydro-4H-1,4-thiazin-4-yl)propoxy, 2-piperidinoethoxy,
 15 3-piperidinopropoxy, piperidin-3-yloxy, piperidin-4-yloxy, piperidin-3-ylmethoxy, 2-piperidin-3-ylethoxy, piperidin-4-ylmethoxy, 2-piperidin-4-ylethoxy, 2-homopiperidin-1-ylethoxy, 3-homopiperidin-1-ylpropoxy, 2-piperazin-1-ylethoxy, 3-piperazin-1-ylpropoxy, 2-homopiperazin-1-ylethoxy, 3-homopiperazin-1-ylpropoxy, pyrrolidin-1-yl, morpholino, piperidino and piperazin-1-yl,
 20 and wherein adjacent carbon atoms in any (2-6C)alkylene chain within a R¹ substituent are optionally separated by the insertion into the chain of a group selected from O, NH, N(CH₃), CH=CH and C≡C,

and when R¹ is a vinyloxy, allyloxy, ethynyloxy or 2-propynyloxy group, the R¹ substituent optionally bears at the terminal CH₂= or HC≡ position a substituent selected from
 25 N-(2-dimethylaminoethyl)carbamoyl, N-(3-dimethylaminopropyl)carbamoyl, methylaminomethyl, 2-methylaminoethyl, 3-methylaminopropyl, 4-methylaminobutyl, dimethylaminomethyl, 2-dimethylaminoethyl, 3-dimethylaminopropyl and 4-dimethylaminobutyl, or from a group of the formula :



- 30 wherein X⁴ is a direct bond or is NHCO or N(CH₃)CO and Q⁴ is pyrrolidin-1-ylmethyl, 2-pyrrolidin-1-ylethyl, 3-pyrrolidin-1-ylpropyl, 4-pyrrolidin-1-ylbutyl, pyrrolidin-2-ylmethyl, 2-pyrrolidin-2-ylethyl, 3-pyrrolidin-2-ylpropyl, morpholinomethyl,

2-morpholinoethyl, 3-morpholinopropyl, 4-morpholinobutyl, piperidinomethyl,
2-piperidinoethyl, 3-piperidinopropyl, 4-piperidinobutyl, piperidin-3-ylmethyl,
2-piperidin-3-ylethyl, piperidin-4-ylmethyl, 2-piperidin-4-ylethyl, piperazin-1-ylmethyl,
2-piperazin-1-ylethyl, 3-piperazin-1-ylpropyl or 4-piperazin-1-ylbutyl,

5 and wherein any CH₂ group which is attached to 2 carbon atoms (other than a CH₂ group within a heterocyclyl ring) or any CH₃ group which is attached to a carbon atom within a R¹ substituent optionally bears on each said CH₂ or CH₃ group a substituent selected from hydroxy, amino, methoxy, ethoxy, methylsulfonyl, methylamino and dimethylamino,

and wherein any heterocyclyl group within a substituent on R¹ optionally bears 1 or 2
10 substituents, which may be the same or different, selected from fluoro, chloro, trifluoromethyl, hydroxy, amino, methylamino, ethylamino, dimethylamino, diethylamino, carbamoyl, methyl, ethyl, n-propyl, isopropyl and methoxy, and any piperidin-3-ylmethyl, piperidin-4-ylmethyl, 2-piperazin-1-ylethylamino, 3-piperazin-1-ylpropylamino, or piperazin-1-yl group within a R¹ substituent is optionally N-substituted with 2-methoxyethyl, 3-methoxypropyl, 2-aminoethyl,
15 3-aminopropyl, 2-methylaminoethyl, 3-methylaminopropyl, 2-dimethylaminoethyl, 3-dimethylaminopropyl, acetyl or propionyl,

and wherein any heterocyclyl group within a substituent on R¹ optionally bears 1 or 2 oxo substituents;

(e) m is 1 and the R¹ group is located at the 7-position and is selected from
20 hydroxy, methoxy, ethoxy, propoxy, butoxy,

and wherein adjacent carbon atoms in any (2-6C)alkylene chain within a R¹ substituent are optionally separated by the insertion into the chain of a group selected from O, NH, N(CH₃), CO, CONH and NHCO,

and wherein any CH₂ group which is attached to 2 carbon atoms (other than a CH₂
25 group within a heterocyclyl ring) or any CH₃ group which is attached to a carbon atom within a R¹ substituent optionally bears on each said CH₂ or CH₃ group a substituent selected from hydroxy, methoxy, ethoxy and methylsulfonyl, or from a group of the formula:



wherein X⁵ is a direct bond or is selected from O, NH and N(CH₃) and Q⁵ is selected
30 from pyrrolidin-1-yl, pyrrolidin-2-yl, pyrrolidin-3-yl, morpholino, piperidino, piperidin-3-yl, piperidin-4-yl, homopiperidin-1-yl, piperazin-1-yl and homopiperazin-1-yl,

and wherein any heterocyclyl group within a substituent on R¹ optionally bears 1 or 2 substituents, which may be the same or different, selected from fluoro, chloro, trifluoromethyl,

hydroxy, methyl, ethyl, n-propyl, isopropyl, methoxy, ethoxy, 2-methoxyethoxy, 2-hydroxyethoxy, 3-hydroxypropoxy, 3-methoxypropoxy, acetyl and methylsulfonyl, and wherein any heterocyclyl group within a substituent on R¹ optionally bears 1 or 2 oxo substituents;

- 5 (f) m is 1 and the R¹ group is located at the 7-position and is selected from hydroxy, methoxy, ethoxy, propoxy, butoxy, 2-pyrrolidin-1-ylethoxy, 3-pyrrolidin-1-ylpropoxy, pyrrolidin-3-yloxy, pyrrolidin-2-ylmethoxy, 2-pyrrolidin-2-ylethoxy, 3-pyrrolidin-2-ylpropoxy, 2-morpholinoethoxy, 3-morpholinopropoxy, 2-piperidinoethoxy, 3-piperidinopropoxy, piperidin-3-yloxy, piperidin-4-yloxy, piperidin-3-ylmethoxy, 10 2-piperidin-3-ylethoxy, piperidin-4-ylmethoxy, 2-piperidin-4-ylethoxy, 2-homopiperidin-1-ylethoxy, 3-homopiperidin-1-ylpropoxy, 2-piperazin-1-ylethoxy, 3-piperazin-1-ylpropoxy, 2-homopiperazin-1-ylethoxy or 3-homopiperazin-1-ylpropoxy, and wherein adjacent carbon atoms in any (2-6C)alkylene chain within a R¹ substituent are optionally separated by the insertion into the chain of a group selected from O, NH,

- 15 N(CH₃), CH=CH and C≡C,

and wherein any CH₂ group which is attached to 2 carbon atoms (other than a CH₂ group within a heterocyclyl ring) or any CH₃ group which is attached to a carbon atom within a R¹ substituent optionally bears on each said CH₂ or CH₃ group a substituent selected from hydroxy, methoxy and methylsulfonyl,

- 20 and wherein any heterocyclyl group within a substituent on R¹ optionally bears 1 or 2 substituents, which may be the same or different, selected from fluoro, chloro, trifluoromethyl, hydroxy, amino, methyl, ethyl, methoxy, methylamino and dimethylamino, and wherein any heterocyclyl group within a substituent on R¹ optionally bears 1 or 2 oxo substituents;

- 25 (g) m is 1 and the R¹ group is located at the 7-position and is selected from hydroxy, methoxy, ethoxy, isopropoxy, propoxy, butoxy, 2-pyrrolidin-1-ylethoxy, 3-pyrrolidin-1-ylpropoxy, 2-piperidinoethoxy, 3-piperidinopropoxy, 2-piperidin-3-ylethoxy, 3-piperidin-3-ylpropoxy, 2-piperidin-4-ylethoxy, 3-piperidin-4-ylpropoxy, 2-piperazin-1-ylethoxy, 3-piperazin-1-ylpropoxy, 2-morpholinoethoxy, 3-morpholinopropoxy, 30 2-homopiperidinoethoxy, 3-homopiperidinopropoxy, 2-homopiperazin-1-ylethoxy and 3-homopiperazin-1-ylpropoxy,

and wherein any terminal CH₃ group within a (1-6C)alkoxy chain in a R¹ substituent optionally bears on the terminal CH₃ group 1, 2 or 3 fluoro substituents or a substituent selected from hydroxy and methoxy,

and wherein any CH₂ group which is attached to 2 carbon atoms (other than a CH₂ group within a heterocyclyl ring) within a R¹ substituent optionally bears on each said CH₂ a hydroxy substituent,

and wherein any heterocyclyl group within a R¹ substituent optionally bears one or more substituents (for example 1, 2 or 3 substituents) selected from fluoro and chloro, or a substituent selected from hydroxy, methyl, ethyl, isopropyl, 2-hydroxyethyl and 2-

10 methoxyethyl;

(h) m is 1 and R¹ is located at the 7-position and is selected from methoxy, ethoxy, propoxy, isopropoxy, cyclopropylmethoxy, 2-hydroxyethoxy, 2-fluoroethoxy, 2-methoxyethoxy, 2-ethoxyethoxy, 2,2-difluoroethoxy, 2,2,2-trifluoroethoxy and 3-hydroxy-3-methylbutoxy;

15 (i) m is 1 and R¹ is located at the 7-position and is selected from (1-4C)alkoxy, hydroxy-(1-4C)alkoxy and (1-4C)alkoxy(1-4C)alkoxy;

(j) m is 1 and R¹ is located at the 7-position and is selected from methoxy, ethoxy, propoxy, isopropoxy, 2-hydroxyethoxy, 2-methoxyethoxy 2-ethoxyethoxy, 3-hydroxypropoxy, 2-hydroxypropoxy, 3-methoxypropoxy and 2-methoxypropoxy;

20 (k) m is 1 and R¹ is located at the 7-position and is selected from methoxy, ethoxy, 2-methoxyethoxy and 2-ethoxyethoxy;

(l) m is 1 and R¹ is located at the 7-position and is (1-4C)alkoxy (particularly methoxy);

(m) m is 0;

(n) m is 1 and R¹ is located at the 7-position and R¹ has any of the values defined in (a) to

25 (d) above;

(o) X¹ is C(R³)₂, wherein each R³, which may be the same or different, is selected from hydrogen and methyl;

(p) X¹ is CH₂;

(q) Q¹ is selected from pyrrolidin-2-yl and pyrrolidin-3-yl,

30 and wherein the group of the formula ZC(O) is attached at the 1-position on Q¹, and wherein Q¹ optionally bears one or more (for example 1, 2 or 3) substituents, which may be the same or different selected from halogeno, trifluoromethyl, hydroxy, carbamoyl, (1-4C)alkyl, (1-4C)alkoxy, N-(1-4C)alkylcarbamoyl and N,N-di-[(1-4C)alkyl]carbamoyl,

and wherein Q¹ optionally bears an oxo substituent;

- (r) Q¹ is selected from pyrrolidin-2-yl and pyrrolidin-3-yl,
and wherein the group of the formula ZC(O) is attached at the 1-position on Q¹,
and wherein Q¹ optionally bears one or more (for example 1, 2 or 3) substituents,

5 which may be the same or different selected from fluoro, chloro, hydroxy, carbamoyl,
(1-4C)alkyl, (1-4C)alkoxy, N-(1-4C)alkylcarbamoyl and N,N-di-[(1-4C)alkyl]carbamoyl,
and wherein Q¹ optionally bears an oxo substituent;

- (s) Q¹ is pyrrolidin-2-yl,
and wherein the group of the formula ZC(O) is attached at the 1-position in Q¹,
10 and wherein Q¹ optionally bears one or more (for example 1, 2 or 3) substituents,
which may be the same or different selected from fluoro, chloro, hydroxy, carbamoyl,
(1-4C)alkyl, (1-4C)alkoxy, N-(1-4C)alkylcarbamoyl and N,N-di-[(1-4C)alkyl]carbamoyl,
and wherein Q¹ optionally bears an oxo substituent;

- (t) Q¹ is pyrrolidin-3-yl,
15 and wherein the group of the formula ZC(O) is attached at the 1-position in Q¹,
and wherein Q¹ optionally bears one or more (for example 1, 2 or 3) substituents,
which may be the same or different selected from fluoro, chloro, hydroxy, carbamoyl,
(1-4C)alkyl, (1-4C)alkoxy, N-(1-4C)alkylcarbamoyl and N,N-di-[(1-4C)alkyl]carbamoyl,
and wherein Q¹ optionally bears an oxo substituent;

20 and wherein Q¹ optionally bears an oxo substituent;

- (u) Q¹ is selected from pyrrolidin-2-yl and pyrrolidin-3-yl,
and wherein the group of the formula ZC(O) is attached at the 1-position in Q¹,
and wherein Q¹ optionally bears 1 or 2 substituents, which may be the same or
different selected from hydroxy, (1-4C)alkyl and (1-4C)alkoxy,

25 and wherein Q¹ optionally bears an oxo substituent;

- (v) Q¹ is pyrrolidin-2-yl,
and wherein the group of the formula ZC(O) is attached at the 1-position in Q¹,
and wherein Q¹ optionally bears 1 or 2 substituents, which may be the same or
different, selected from hydroxy, (1-4C)alkyl, (1-4C)alkoxy, hydroxy-(1-4C)alkyl,

30 (1-4C)alkoxy-(1-4C)alkyl, hydroxy-(1-4C)alkoxy, (1-4C)alkoxy-(1-4C)alkoxy,
and wherein Q¹ optionally bears an oxo substituent;

- (w) Q¹ is pyrrolidin-2-yl,
and wherein the group of the formula ZC(O) is attached at the 1-position in Q¹,

- and wherein Q^1 optionally bears 1 or 2 substituents, which may be the same or different selected from hydroxy, (1-4C)alkyl and (1-4C)alkoxy,
- and wherein Q^1 optionally bears an oxo substituent;
- (x) Q^1 is pyrrolidin-2-yl,
- 5 and wherein the group of the formula $ZC(O)$ is attached at the 1-position in Q^1 ,
and wherein Q^1 optionally bears an oxo substituent;
- (y) Q^1 is pyrrolidin-3-yl,
- and wherein the group of the formula $ZC(O)$ is attached at the 1-position in Q^1 ,
and wherein Q^1 optionally bears an oxo substituent;
- 10 (z) Q^1-X^1 is selected from pyrrolidin-2-ylmethyl and pyrrolidin-3-ylmethyl,
and wherein Q^1 optionally bears one or more (for example 1, 2 or 3) substituents,
which may be the same or different selected from fluoro, chloro, hydroxy, carbamoyl, methyl,
methoxy, N-methylcarbamoyl and N,N-dimethylcarbamoyl,
and wherein any heterocyclyl group within Q^1 optionally bears an oxo substituent,
- 15 and wherein the group of the formula $ZC(O)$ is attached at the 1-position in Q^1 ;
- (aa) Q^1-X^1 is selected from pyrrolidin-2-ylmethyl and pyrrolidin-2-ylmethyl,
and wherein Q^1 optionally bears one or more (for example 1, 2 or 3) substituents,
which may be the same or different selected from fluoro, chloro, hydroxy, methyl and
methoxy,
- 20 and wherein Q^1 optionally bears an oxo substituent,
and wherein the group of the formula $ZC(O)$ is attached at the 1-position in Q^1 ;
- (bb) Q^1-X^1 is pyrrolidin-2-ylmethyl
and wherein Q^1 optionally bears one or more (for example 1, 2 or 3) substituents,
which may be the same or different selected from hydroxy, (1-4C)alkyl and (1-4C)alkoxy
- 25 (particularly hydroxy, methyl and methoxy),
and wherein Q^1 optionally bears an oxo substituent,
and wherein the group of the formula $ZC(O)$ is attached at the 1-position in Q^1 ;
- (cc) Q^1-X^1 is pyrrolidin-2-ylmethyl
and wherein Q^1 optionally bears one or more (for example 1 or 2, particularly 1)
- 30 substituents, which may be the same or different selected from hydroxy, methyl, ethyl,
methoxy, ethoxy, hydroxymethyl, methoxymethyl, 2-hydroxyethoxy and 2-methoxyethoxy (
particularly Q^1 optionally bears one of the hereinabove defined substituents at the 4-position

on the pyrrolidinyl ring, more particularly Q¹ optionally bears a substituent at the 4-position selected from hydroxy, methyl and methoxy),

and wherein the group of the formula ZC(O) is attached at the 1-position in Q¹;

(dd) Q¹-X¹ is (2R)-pyrrolidin-2-ylmethyl,

5 and wherein Q¹ optionally bears one or more (for example 1, 2 or 3) substituents, which may be the same or different selected from halogeno, hydroxy, (1-4C)alkyl and (1-4C)alkoxy (particularly hydroxy, methyl and methoxy),

and wherein Q¹ optionally bears an oxo substituent,

and wherein the group of the formula ZC(O) is attached at the 1-position in Q¹;

10 (ee) Q¹-X¹ is (2S)-pyrrolidin-2-ylmethyl,

and wherein Q¹ optionally bears one or more (for example 1, 2 or 3) substituents, which may be the same or different selected from halogeno, hydroxy, (1-4C)alkyl and (1-4C)alkoxy (particularly hydroxy, methyl and methoxy),

and wherein Q¹ optionally bears an oxo substituent,

15 and wherein the group of the formula ZC(O) is attached at the 1-position in Q¹;

(ff) Q¹-X¹ is (2R)-pyrrolidin-2-ylmethyl,

and wherein the pyrrolidinyl group optionally bears one or two substituents, which may be the same or different selected from fluoro, chloro, hydroxy, carbamoyl, (1-3C)alkyl (particularly methyl), (1-3C)alkoxy (particularly methoxy), N-methylcarbamoyl and N,N-

20 dimethylcarbamoyl,

and wherein the pyrrolidinyl group optionally bears an oxo substituent,

and wherein the group of the formula ZC(O) is attached at the 1-position of the pyrrolidinyl group;

(gg) Q¹-X¹ is (2S)-pyrrolidin-2-ylmethyl,

25 and wherein the pyrrolidinyl group optionally bears one or two substituents, which may be the same or different selected from fluoro, chloro, hydroxy, carbamoyl, (1-3C)alkyl (particularly methyl), (1-3C)alkoxy (particularly methoxy), N-methylcarbamoyl and N,N-dimethylcarbamoyl,

and wherein the pyrrolidinyl group optionally bears an oxo substituent,

30 and wherein the group of the formula ZC(O) is attached at the 1-position of the pyrrolidinyl group;

(hh) Q¹-X¹ is selected from (3S)-pyrrolidin-3-ylmethyl and (3R)-pyrrolidin-3-ylmethyl,

and wherein the pyrrolidinyl group optionally bears one or two substituents, which may be the same or different selected from fluoro, chloro, hydroxy, carbamoyl, (1-3C)alkyl (particularly methyl), (1-3C)alkoxy (particularly methoxy), N-methylcarbamoyl and N,N-dimethylcarbamoyl,

5 and wherein the pyrrolidinyl group optionally bears an oxo substituent,
and wherein the group of the formula ZC(O) is attached at the 1-position of the pyrrolidinyl group;

(ii) Z is selected from halogeno-(1-2C)alkyl, hydroxy-(1-2C)alkyl, (1-4C)alkoxy-(1-2C)alkyl, amino-(1-2C)alkyl, (1-4C)alkylamino-(1-2C)alkyl and

10 di-[(1-4C)alkyl]amino-(1-2C)alkyl,


or Z is Q², wherein Q² is heterocyclyl or heterocyclyl-(1-2C)alkyl, and wherein the heterocyclyl in Q² contains at least 1 nitrogen heteroatom, and optionally 1 or 2 further heteroatoms selected from oxygen, nitrogen and sulfur,

and wherein Q² is linked to the (1-2C)alkyl or C(O) group by a ring nitrogen,

15 and wherein any CH₂ or CH₃ group within a Z group, other than a CH₂ group within a heterocyclyl ring, optionally bears on each said CH₂ or CH₃ group one or more halogeno or (1-6C)alkyl substituents or a substituent selected from hydroxy, cyano, amino, carbamoyl, formyl, mercapto, (2-6C)alkenyl, (2-6C)alkynyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulfinyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino,

20 N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino and N-(1-6C)alkyl-(2-6C)alkanoylamino,

and wherein any heterocyclyl group within Z optionally bears one or more (for example 1, 2 or 3) substituents, which may be the same or different selected from halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carbamoyl, formyl, (1-6C)alkyl, (2-6C)alkenyl, (2-6C)alkynyl, (1-6C)alkoxy, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, N-(1-6C)alkylsulfamoyl and N,N-di-[(1-6C)alkyl]sulfamoyl, or from a group of the formula:



30 wherein X² is a direct bond or is selected from O, CO and N(R⁵), wherein R⁵ is hydrogen or (1-6C)alkyl, and R⁴ is halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl, cyano-(1-6C)alkyl, amino-(1-6C)alkyl, N-(1-6C)alkylamino-(1-6C)alkyl, N,N-di-[(1-6C)alkyl]amino-(1-6C)alkyl,

carbamoyl-(1-6C)alkyl, N-(1-6C)alkylcarbamoyl-(1-6C)alkyl and
N,N-di-[(1-6C)alkyl]carbamoyl-(1-6C)alkyl,

and wherein any heterocyclyl group within Z optionally bears 1 or 2 oxo substituents;

(jj) Z is selected from halogeno-(1-2C)alkyl, hydroxy-(1-2C)alkyl, (1-
 5 4C)alkoxy-(1-2C)alkyl, amino-(1-2C)alkyl, (1-4C)alkylamino-(1-2C)alkyl and
 di-[(1-4C)alkyl]amino-(1-2C)alkyl,


or Z is Q², wherein Q² is heterocyclyl or heterocyclyl-(1-2C)alkyl, and wherein the
 heterocyclyl in Q² is a fully saturated 4, 5, 6 or 7 membered monocyclic heterocyclyl group
 which contains at least 1 nitrogen heteroatom, and optionally 1 or 2 further heteroatoms
 10 selected from oxygen, nitrogen and sulfur,

and wherein Q² is linked to the (1-2C)alkyl or C(O) group by a ring nitrogen,

and wherein any CH₂ or CH₃ group within a Z group, other than a CH₂ group within a
 heterocyclyl ring, optionally bears on each said CH₂ or CH₃ group one or more (1-4C)alkyl
 substituents or a substituent selected from cyano, carbamoyl, formyl, (2-6C)alkenyl,
 15 (2-6C)alkynyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl and (2-6C)alkanoyl,

and wherein any CH₂ which is attached to 2 carbon atoms or a CH₃ group which is
 attached to a carbon atom within a Z group, other than a CH₂ group within a heterocyclyl
 ring, optionally bears on each said CH₂ or CH₃ group one or more halogeno substituents or a
 substituent selected from hydroxy, amino, mercapto, (1-6C)alkoxy, (1-6C)alkylthio,
 20 (1-6C)alkylsulfinyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino,
 (2-6C)alkanoyloxy, (2-6C)alkanoylamino and N-(1-6C)alkyl-(2-6C)alkanoylamino,

and wherein any heterocyclyl group within Z optionally bears one or more (for
 example 1, 2 or 3) substituents, which may be the same or different selected from halogeno,
 trifluoromethyl, cyano, nitro, hydroxy, amino, carbamoyl, formyl, (1-6C)alkyl, (2-6C)alkenyl,
 25 (2-6C)alkynyl, (1-6C)alkoxy, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino,
 (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl,
 (2-6C)alkanoyl, N-(1-6C)alkylsulfamoyl and N,N-di-[(1-6C)alkyl]sulfamoyl, or from a group
 of the formula:



30 wherein X² is a direct bond or is selected from O, CO and N(R⁵), wherein R⁵ is
 hydrogen or (1-4C)alkyl, and R⁴ is halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl,
 (1-6C)alkoxy-(1-6C)alkyl, cyano-(1-6C)alkyl, amino-(1-6C)alkyl,
N-(1-6C)alkylamino-(1-6C)alkyl, N,N-di-[(1-6C)alkyl]amino-(1-6C)alkyl,

carbamoyl-(1-6C)alkyl, N-(1-6C)alkylcarbamoyl-(1-6C)alkyl and

N,N-di-[(1-6C)alkyl]carbamoyl-(1-6C)alkyl,

and wherein any heterocyclyl group within Z optionally bears 1 or 2 oxo substituents;

(kk) Z is selected from hydroxy-(1-2C)alkyl, (1-4C)alkoxy-(1-2C)alkyl, amino-(1-2C)alkyl,

5 (1-4C)alkylamino-(1-2C)alkyl and di-[(1-4C)alkyl]amino-(1-2C)alkyl,

or Z is Q², wherein Q² is heterocyclyl or heterocyclyl-(1-2C)alkyl, and wherein the heterocyclyl in Q² is a fully saturated 4, 5, 6 or 7 membered monocyclic heterocyclyl group which contains at least 1 nitrogen heteroatom, and optionally 1 or 2 further heteroatoms selected from oxygen, nitrogen and sulfur,

10 and wherein Q² is linked to the (1-2C)alkyl or C(O) group by a ring nitrogen,


and wherein any CH₂ or CH₃ group within a Z group, other than a CH₂ group within a heterocyclyl ring, optionally bears on each said CH₂ or CH₃ group one or more (1-4C)alkyl substituents or a substituent selected from cyano, carbamoyl, formyl, (2-6C)alkenyl, (2-6C)alkynyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl and (2-6C)alkanoyl,

15 and wherein any CH₂ which is attached to 2 carbon atoms or a CH₃ group which is attached to a carbon atom within a Z group, other than a CH₂ group within a heterocyclyl ring, optionally bears on each said CH₂ or CH₃ group one or more halogeno substituents or a substituent selected from hydroxy, amino, mercapto, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulfinyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino,

20 (2-6C)alkanoyloxy, (2-6C)alkanoylamino and N-(1-6C)alkyl-(2-6C)alkanoylamino,

and wherein any heterocyclyl group within Z optionally bears one or more (for example 1, 2 or 3) substituents, which may be the same or different selected from halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carbamoyl, formyl, (1-6C)alkyl, (2-6C)alkenyl, (2-6C)alkynyl, (1-6C)alkoxy, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino,

25 (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, N-(1-6C)alkylsulfamoyl and N,N-di-[(1-6C)alkyl]sulfamoyl, or from a group of the formula:



wherein X² is a direct bond or is selected from O, CO and N(R⁵), wherein R⁵ is

30 hydrogen or (1-4C)alkyl, and R⁴ is halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl,

(1-6C)alkoxy-(1-6C)alkyl, cyano-(1-6C)alkyl, amino-(1-6C)alkyl,

N-(1-6C)alkylamino-(1-6C)alkyl, N,N-di-[(1-6C)alkyl]amino-(1-6C)alkyl,

carbamoyl-(1-6C)alkyl, N-(1-6C)alkylcarbamoyl-(1-6C)alkyl and
N,N-di-[(1-6C)alkyl]carbamoyl-(1-6C)alkyl,

and wherein any heterocyclyl group within Z optionally bears 1 or 2 oxo substituents;

(II) Z is selected from halogeno-(1-2C)alkyl, hydroxy-(1-2C)alkyl, (1-
 5 4C)alkoxy-(1-2C)alkyl, amino-(1-2C)alkyl, (1-4C)alkylamino-(1-2C)alkyl and
 di-[(1-4C)alkyl]amino-(1-2C)alkyl,

or Z is Q², wherein Q² is heterocyclyl or heterocyclyl-(1-2C)alkyl, and wherein the
 heterocyclyl in Q² is selected from azetidin-1-yl, pyrrolidin-1-yl, piperidino, piperazin-1-yl,
 morpholino, homopiperidin-1-yl and homopiperazin-1-yl,

10 and wherein Q² is linked to the (1-2C)alkyl or C(O) group by a ring nitrogen,

and wherein any CH₂ or CH₃ group within a Z group, other than a CH₂ group within a
 heterocyclyl ring, optionally bears on each said CH₂ or CH₃ group one or more (1-4C)alkyl
 substituents or a substituent selected from cyano, carbamoyl, (2-6C)alkenyl, (2-6C)alkynyl,
N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl and (2-6C)alkanoyl,

15 and wherein any CH₂ which is attached to 2 carbon atoms or a CH₃ group which is
 attached to a carbon atom within a Z group, other than a CH₂ group within a heterocyclyl
 ring, optionally bears on each said CH₂ or CH₃ group one or more halogeno substituents or a
 substituent selected from hydroxy, amino, (1-6C)alkoxy, (1-6C)alkylsulfonyl,
 (1-6C)alkylamino and di-[(1-6C)alkyl]amino,

20 and wherein any heterocyclyl group within Z optionally bears one or more (for
 example 1, 2 or 3) substituents, which may be the same or different selected from halogeno,
 trifluoromethyl, cyano, hydroxy, amino, carbamoyl, (1-6C)alkyl, (1-6C)alkoxy,
 (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, N-(1-6C)alkylcarbamoyl,
N,N-di-[(1-6C)alkyl]carbamoyl and (2-6C)alkanoyl, or from a group of the formula:

25 $-X^2-R^4$

wherein X² is a direct bond or is selected from O, CO and N(R⁵); wherein R⁵ is
 hydrogen or (1-4C)alkyl, and R⁴ is halogeno-(1-4C)alkyl, hydroxy-(1-4C)alkyl,
 (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, amino-(1-4C)alkyl,
N-(1-4C)alkylamino-(1-4C)alkyl, N,N-di-[(1-4C)alkyl]amino-(1-4C)alkyl,

30 carbamoyl-(1-4C)alkyl, N-(1-4C)alkylcarbamoyl-(1-4C)alkyl and
N,N-di-[(1-4C)alkyl]carbamoyl-(1-4C)alkyl,

and wherein any heterocyclyl group within Z optionally bears 1 or 2 oxo substituents;

(mm) Z is selected from halogeno-(1-2C)alkyl, hydroxy-(1-2C)alkyl, (1-4C)alkoxy-(1-2C)alkyl, amino-(1-2C)alkyl, (1-4C)alkylamino-(1-2C)alkyl and di-[(1-4C)alkyl]amino-(1-2C)alkyl,


or Z is Q², wherein Q² is heterocyclyl or heterocyclyl-(1-2C)alkyl, and wherein the
 5 heterocyclyl in Q² is selected from pyrrolidin-1-yl, piperidino, piperazin-1-yl, morpholino, homopiperidin-1-yl and homopiperazin-1-yl

and wherein Q² is linked to the (1-2C)alkyl or C(O) group by a ring nitrogen,

and wherein any CH₂ or CH₃ group within a Z group, other than a CH₂ group within a heterocyclyl ring, optionally bears on each said CH₂ or CH₃ group a substituent selected from
 10 (2-6C)alkenyl and (2-6C)alkynyl,

and wherein any CH₂ which is attached to 2 carbon atoms or a CH₃ group which is attached to a carbon atom within a Z group, other than a CH₂ group within a heterocyclyl ring, optionally bears on each said CH₂ or CH₃ group one or more halogeno substituents or a substituent selected from hydroxy, amino, (1-4C)alkoxy, (1-4C)alkylamino and
 15 di-[(1-4C)alkyl]amino,

and wherein any heterocyclyl group within Z optionally bears one or more (for example 1, 2 or 3) substituents, which may be the same or different selected from fluoro, chloro, cyano, hydroxy, amino, carbamoyl, (1-4C)alkyl, (1-4C)alkoxy, (1-4C)alkylamino, di-[(1-4C)alkyl]amino, N-(1-4C)alkylcarbamoyl, N,N-di-[(1-4C)alkyl]carbamoyl and
 20 (2-6C)alkanoyl, or from a group of the formula:



wherein X² is a direct bond or is selected from O, CO and N(R⁵), wherein R⁵ is hydrogen or (1-4C)alkyl, and R⁴ is halogeno-(1-4C)alkyl, hydroxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, amino-(1-4C)alkyl,
 25 N-(1-4C)alkylamino-(1-4C)alkyl and N,N-di-[(1-4C)alkyl]amino-(1-4C)alkyl,

and wherein any heterocyclyl group within Z optionally bears an oxo substituent;

(nn) Z is selected from hydroxy-(1-2C)alkyl, (1-4C)alkoxy-(1-2C)alkyl, amino-(1-2C)alkyl, (1-4C)alkylamino-(1-2C)alkyl and di-[(1-4C)alkyl]amino-(1-2C)alkyl,


or Z is Q², wherein Q² is heterocyclyl or heterocyclyl-(1-2C)alkyl, and wherein the
 30 heterocyclyl in Q² is selected from pyrrolidin-1-yl, piperidino, piperazin-1-yl, morpholino, homopiperidin-1-yl and homopiperazin-1-yl

and wherein Q² is linked to the (1-2C)alkyl or C(O) group by a ring nitrogen,

and wherein any CH₂ or CH₃ group within a Z group, other than a CH₂ group within a heterocyclyl ring, optionally bears on each said CH₂ or CH₃ group a substituent selected from (2-6C)alkenyl and (2-6C)alkynyl,

and wherein any CH₂ which is attached to 2 carbon atoms or a CH₃ group which is
 5 attached to a carbon atom within a Z group, other than a CH₂ group within a heterocyclyl ring, optionally bears on each said CH₂ or CH₃ group one or more halogeno substituents or a substituent selected from hydroxy, amino, (1-4C)alkoxy, (1-4C)alkylamino and di-[(1-4C)alkyl]amino,

and wherein any heterocyclyl group within Z optionally bears one or more (for
 10 example 1, 2 or 3) substituents, which may be the same or different selected from fluoro, chloro, cyano, hydroxy, amino, carbamoyl, (1-4C)alkyl, (1-4C)alkoxy, (1-4C)alkylamino, di-[(1-4C)alkyl]amino, N-(1-4C)alkylcarbamoyl, N,N-di-[(1-4C)alkyl]carbamoyl and (2-6C)alkanoyl, or from a group of the formula:




15 wherein X² is a direct bond or is selected from O, CO and N(R⁵), wherein R⁵ is hydrogen or (1-4C)alkyl, and R⁴ is halogeno-(1-4C)alkyl, hydroxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, amino-(1-4C)alkyl, N-(1-4C)alkylamino-(1-4C)alkyl and N,N-di-[(1-4C)alkyl]amino-(1-4C)alkyl, and wherein any heterocyclyl group within Z optionally bears an oxo substituent;

20 (oo) Z is selected from (1-2C)alkyl, hydroxy-(1-2C)alkyl, amino-(1-2C)alkyl, (1-4C)alkylamino-(1-2C)alkyl and di-[(1-4C)alkyl]amino-(1-2C)alkyl,

or Z is Q², wherein Q² is heterocyclyl-(1-2C)alkyl, and wherein the heterocyclyl in Q² is selected from pyrrolidin-1-yl, piperidino and piperazin-1-yl (particularly pyrrolidin-1-yl),

and wherein Q² is linked to the (1-2C)alkyl group by a ring nitrogen,

25 and wherein any heterocyclyl group within Z optionally bears one or more (for example 1, 2 or 3) substituents, which may be the same or different selected from fluoro, chloro, hydroxy, (1-3C)alkyl, (1-3C)alkoxy, or from a group of the formula:



wherein X² is a direct bond or is O, and R⁴ is halogeno-(1-3C)alkyl,
 30 hydroxy-(1-3C)alkyl or (1-3C)alkoxy-(1-3C)alkyl,

and wherein any heterocyclyl group within Z optionally bears an oxo substituent;


(pp) Z is selected from fluoromethyl, hydroxymethyl, (1-4C)alkoxy-methyl, aminomethyl, (1-4C)alkylaminomethyl, di-[(1-4C)alkyl]aminomethyl, pyrrolidin-1-ylmethyl,

piperidinomethyl, piperazin-1-ylmethyl, morpholinomethyl, homopiperidin-1-ylmethyl and homopiperazin-1-ylmethyl

and wherein any CH₂ or CH₃ group within a Z group, other than a CH₂ group within a heterocyclyl ring, optionally bears on each said CH₂ or CH₃ group a substituent selected from
5 vinyl and ethynyl,

and wherein any CH₂ which is attached to 2 carbon atoms or a CH₃ group which is attached to a carbon atom within a Z group, other than a CH₂ group within a heterocyclyl ring, optionally bears on each said CH₂ or CH₃ group one or more halogeno substituents or a substituent selected from hydroxy, amino, (1-4C)alkoxy, (1-4C)alkylamino and
10 di-[(1-4C)alkyl]amino,

and wherein any heterocyclyl group within Z optionally bears one or more (for example 1, 2 or 3) substituents, which may be the same or different selected from fluoro, chloro, cyano, hydroxy, amino, carbamoyl, (1-4C)alkyl, (1-4C)alkoxy, (1-4C)alkylamino, di-[(1-4C)alkyl]amino, N-(1-4C)alkylcarbamoyl, N,N-di-[(1-4C)alkyl]carbamoyl and
15 (2-6C)alkanoyl, or from a group of the formula:



wherein X² is a direct bond or is selected from O, CO and N(R⁵), wherein R⁵ is hydrogen or (1-4C)alkyl, and R⁴ is halogeno-(1-4C)alkyl, hydroxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, amino-(1-4C)alkyl,
20 N-(1-4C)alkylamino-(1-4C)alkyl and N,N-di-[(1-4C)alkyl]amino-(1-4C)alkyl,
and wherein any heterocyclyl group within Z optionally bears an oxo substituent;
(qq) Z is selected from hydroxymethyl, (1-4C)alkoxy-methyl, aminomethyl, (1-4C)alkylaminomethyl, N-allylaminomethyl, N-(2-propynyl)aminomethyl, di-[(1-4C)alkyl]aminomethyl, N-allyl-N-[(1-4C)alkyl]aminomethyl, N-(2-propynyl)-N-
25 [(1-4C)alkyl]aminomethyl, pyrrolidin-1-ylmethyl, piperidinomethyl, piperazin-1-ylmethyl, morpholinomethyl, homopiperidin-1-ylmethyl and homopiperazin-1-ylmethyl

and wherein any CH₂ which is attached to 2 carbon atoms or a CH₃ group which is attached to a carbon atom within a Z group, other than a CH₂ group within a heterocyclyl ring, optionally bears on each said CH₂ or CH₃ group one or more substituents selected from
30 fluoro and chloro, or a substituent selected from hydroxy and (1-4C)alkoxy,

and wherein any heterocyclyl group within Z optionally bears one or more (for example 1, 2 or 3) substituents, which may be the same or different selected from fluoro, chloro, cyano, hydroxy, amino, carbamoyl, (1-4C)alkyl, (1-4C)alkoxy, (1-4C)alkylamino,

di-[(1-4C)alkyl]amino, N-(1-4C)alkylcarbonyl, N,N-di-[(1-4C)alkyl]carbonyl, acetyl, propionyl, 2-fluoroethyl, 2-hydroxyethyl, 2-methoxyethyl, cyanomethyl, hydroxyacetyl, aminoacetyl, methylaminoacetyl, ethylaminoacetyl, dimethylaminoacetyl, N-methyl-N-ethylaminoacetyl and fluoroacetyl,

5 and wherein any heterocyclyl group within Z optionally bears an oxo substituent;

(rr) Z is selected from hydroxymethyl, methoxymethyl, ethoxymethyl, aminomethyl, (1-4C)alkylaminomethyl, N-[hydroxy-(2-4C)alkyl]-aminomethyl, N-[methoxy-(2-4C)alkyl]-aminomethyl, di-[(1-4C)alkyl]aminomethyl, N-[hydroxy-(2-4C)alkyl]-N-[(1-4C)alkyl]aminomethyl, N,N-di-[hydroxy-(2-4C)alkyl]-aminomethyl, N-[methoxy-(2-4C)alkyl]-N-[(1-4C)alkyl]aminomethyl, pyrrolidin-1-ylmethyl, piperidinomethyl, 10 piperazin-1-ylmethyl, morpholinomethyl, homopiperidin-1-ylmethyl and homopiperazin-1-ylmethyl,

and wherein any heterocyclyl group within Z optionally bears one or more (for example 1, 2 or 3) substituents, which may be the same or different selected from fluoro, 15 chloro, cyano, hydroxy, amino, (1-4C)alkyl, (1-4C)alkoxy, (1-4C)alkylamino and di-[(1-4C)alkyl]amino,

and wherein any heterocyclyl group within Z optionally bears an oxo substituent; (ss) Z is selected from hydroxymethyl, methoxymethyl, ethoxymethyl, aminomethyl, methylaminomethyl, ethylaminomethyl, N-(2-hydroxyethyl)aminomethyl, N-(2-methoxyethyl)aminomethyl, dimethylaminomethyl, N-methyl-N-ethylaminomethyl, di-ethylaminomethyl, N-(2-hydroxyethyl)-N-methylaminomethyl, N-(2-hydroxyethyl)-N-ethylaminomethyl, N,N-di-(2-hydroxyethyl)aminomethyl, N-(2-methoxyethyl)-N-methylaminomethyl, N-(2-methoxyethyl)-N-ethylaminomethyl, pyrrolidin-1-ylmethyl, piperidinomethyl, piperazin-1-ylmethyl and morpholinomethyl,

25 and wherein any heterocyclyl group within Z optionally bears 1 or 2 substituents, which may be the same or different, selected from fluoro, chloro, hydroxy, (1-4C)alkyl and (1-4C)alkoxy;

(tt) Z is selected from hydroxymethyl, methoxymethyl, ethoxymethyl, aminomethyl, (1-4C)alkylaminomethyl, N-[hydroxy-(2-4C)alkyl]-aminomethyl, N-[methoxy-(2-4C)alkyl]-aminomethyl, di-[(1-4C)alkyl]aminomethyl, N-[hydroxy-(2-4C)alkyl]-N-[(1-4C)alkyl]aminomethyl, N,N-di-[hydroxy-(2-4C)alkyl]-aminomethyl and N-[methoxy-(2-4C)alkyl]-N-[(1-4C)alkyl]aminomethyl;

- (uu) Z is selected from pyrrolidin-1-ylmethyl, piperidinomethyl, piperazin-1-ylmethyl, morpholinomethyl, homopiperidin-1-ylmethyl, homopiperazin-1-ylmethyl, 2-pyrrolidin-1-ylethyl, 2-piperidinoethyl, 2-piperazin-1-ylethyl, 2-morpholinoethyl, 2-homopiperidin-1-ylethyl and 2-homopiperazin-1-ylethyl (particularly Z is selected from
5 pyrrolidin-1-ylmethyl, piperidinomethyl, piperazin-1-ylmethyl and morpholinomethyl, and wherein the heterocyclyl group within Z optionally bears one or more (for example 1, 2 or 3) substituents, which may be the same or different selected from fluoro, chloro, cyano, hydroxy, amino, carbamoyl, (1-4C)alkyl, (1-4C)alkoxy, (1-4C)alkylamino, di-[(1-4C)alkyl]amino, N-(1-4C)alkylcarbamoyl, N,N-di-[(1-4C)alkyl]carbamoyl, acetyl,
10 propionyl, 2-fluoroethyl, 2-hydroxyethyl, 2-methoxyethyl, cyanomethyl, hydroxyacetyl, aminoacetyl, methylaminoacetyl, ethylaminoacetyl, dimethylaminoacetyl, N-methyl-N-ethylaminoacetyl and fluoroacetyl,
- (vv) Z is selected from methyl, hydroxymethyl, (1-3C)alkoxymethyl, aminomethyl, (1-3C)alkylaminomethyl, N-di-[(1-3C)alkyl]aminomethyl and pyrrolidin-1-ylmethyl,
15 and wherein any heterocyclyl group within Z optionally bears one or more (for example 1, 2 or 3) substituents, which may be the same or different selected from fluoro, chloro, hydroxy, (1-3C)alkyl and (1-3C)alkoxy;
- (ww) Q¹-X¹ is selected from (2R)-pyrrolidin-2-ylmethyl and (2S)-pyrrolidin-2-ylmethyl, and wherein the group of the formula ZC(O) is attached at the 1-position of the
20 pyrrolidinyl group,
- Z is selected from hydroxymethyl, methoxymethyl, ethoxymethyl, aminomethyl, (1-4C)alkylaminomethyl, N-[hydroxy-(2-4C)alkyl]-aminomethyl, N-[methoxy-(2-4C)alkyl]-aminomethyl, di-[(1-4C)alkyl]aminomethyl, N-[hydroxy-(2-4C)alkyl]-N-[(1-4C)alkyl]aminomethyl, N,N-di-[hydroxy-(2-4C)alkyl]-aminomethyl, N-[methoxy-(2-4C)alkyl]-N-[(1-4C)alkyl]aminomethyl, pyrrolidin-1-ylmethyl, piperidinomethyl,
25 piperazin-1-ylmethyl, morpholinomethyl, homopiperidin-1-ylmethyl and homopiperazin-1-ylmethyl,
- and wherein any heterocyclyl group within Z optionally bears one or more (for example 1, 2 or 3) substituents, which may be the same or different selected from fluoro,
30 chloro, hydroxy, (1-4C)alkyl and (1-4C)alkoxy,
- and wherein any heterocyclyl group within Z optionally bears an oxo substituent;
- (xx) Q¹-X¹ is selected from (3R)-pyrrolidin-3-ylmethyl and (3S)-pyrrolidin-3-ylmethyl,

and wherein the group of the formula ZC(O) is attached at the 1-position of the pyrrolidinyl group,

Z is selected from hydroxymethyl, methoxymethyl, ethoxymethyl, aminomethyl, (1-4C)alkylaminomethyl, N-[hydroxy-(2-4C)alkyl]-aminomethyl, N-[methoxy-(2-4C)alkyl]-aminomethyl, di-[(1-4C)alkyl]aminomethyl, N-[hydroxy-(2-4C)alkyl]-N-[(1-4C)alkyl]aminomethyl, N,N-di-[hydroxy-(2-4C)alkyl]-aminomethyl, N-[methoxy-(2-4C)alkyl]-N-[(1-4C)alkyl]aminomethyl, pyrrolidin-1-ylmethyl, piperidinomethyl, piperazin-1-ylmethyl, morpholinomethyl, homopiperidin-1-ylmethyl and homopiperazin-1-ylmethyl,

10 and wherein any heterocyclyl group within Z optionally bears one or more (for example 1, 2 or 3) substituents, which may be the same or different selected from fluoro, chloro, hydroxy, (1-4C)alkyl and (1-4C)alkoxy,

and wherein any heterocyclyl group within Z optionally bears an oxo substituent;

(yy) Q¹-X¹ is selected from (2R)-pyrrolidin-2-ylmethyl and (2S)-pyrrolidin-2-ylmethyl,

15 and wherein the group of the formula ZC(O) is attached at the 1-position of the pyrrolidinyl group,

Z is selected from hydroxymethyl, methoxymethyl, ethoxymethyl, aminomethyl, (1-4C)alkylaminomethyl, N-[hydroxy-(2-4C)alkyl]-aminomethyl, N-[methoxy-(2-4C)alkyl]-aminomethyl, di-[(1-4C)alkyl]aminomethyl, N-[hydroxy-(2-4C)alkyl]-N-[(1-4C)alkyl]aminomethyl, N,N-di-[hydroxy-(2-4C)alkyl]-aminomethyl, N-[methoxy-(2-4C)alkyl]-N-[(1-4C)alkyl]aminomethyl, pyrrolidin-1-ylmethyl, piperidinomethyl, piperazin-1-ylmethyl, morpholinomethyl, homopiperidin-1-ylmethyl and homopiperazin-1-ylmethyl,

25 and wherein any heterocyclyl group within Z optionally bears one or more (for example 1, 2 or 3) substituents, which may be the same or different selected from fluoro, chloro, hydroxy, (1-4C)alkyl and (1-4C)alkoxy,

and wherein any heterocyclyl group within Z optionally bears an oxo substituent;

(zz) Q¹-X¹ is selected from (2R)-pyrrolidin-2-ylmethyl and (2S)-pyrrolidin-2-ylmethyl (particularly Q¹-X¹ is (2R)-pyrrolidin-2-ylmethyl),

30 and wherein the group of the formula ZC(O) is attached at the 1-position of the pyrrolidin-2-yl group,

Z is selected from methyl, hydroxymethyl, aminomethyl, (1-2C)alkylaminomethyl, di-[(1-2C)alkyl]aminomethyl and pyrrolidin-1-ylmethyl,

and wherein any heterocyclyl group within Z or Q¹ optionally bears one or more (for example 1, 2 or 3) substituents, which may be the same or different selected from hydroxy, (1-3C)alkyl and (1-3C)alkoxy,

and wherein any heterocyclyl group within Z optionally bears an oxo substituent;

- 5 (aaa) Q¹-X¹ is selected from (2R)-pyrrolidin-2-ylmethyl and (2S)-pyrrolidin-2-ylmethyl (particularly Q¹-X¹ is (2R)-pyrrolidin-2-ylmethyl),

and wherein the group of the formula ZC(O) is attached at the 1-position of the pyrrolidin-2-yl group,

Z is selected from hydroxymethyl, aminomethyl, (1-2C)alkylaminomethyl,

- 10 di-[(1-2C)alkyl]aminomethyl and pyrrolidin-1-ylmethyl,

and wherein any heterocyclyl group within Z or Q¹ optionally bears one or more (for example 1, 2 or 3) substituents, which may be the same or different selected from hydroxy, (1-3C)alkyl and (1-3C)alkoxy;

- (bbb) Q¹-X¹ is selected from (2R)-pyrrolidin-2-ylmethyl and (2S)-pyrrolidin-2-ylmethyl
15 (particularly Q¹-X¹ is (2R)-pyrrolidin-2-ylmethyl),

and wherein the group of the formula ZC(O) is attached at the 1-position of the pyrrolidin-2-yl group,

Z is methyl,

and wherein Q¹ optionally bears one or more (for example 1, 2 or 3) substituents,

- 20 which may be the same or different selected from hydroxy, (1-3C)alkyl and (1-3C)alkoxy;

(ccc) Y is selected from halogeno, cyano, (1-6C)alkyl, (1-6C)alkoxy and (2-6C)alkynyl;

(ddd) Y is selected from fluoro, chloro, bromo, iodo, cyano, methyl, ethyl, isopropyl, methoxy, ethoxy, vinyl, allyl, ethynyl, 1-propynyl and 2-propynyl;

(eee) Y is selected from fluoro, chloro, bromo, methyl and ethynyl;

- 25 (fff) Y is halogeno (particularly fluoro, chloro or bromo);

(ggg) Y is methyl;

(hhh) Y is ethynyl;

(iii) each R², which may be the same or different, is selected from halogeno, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, trifluoromethyl, (1-6C)alkyl, (2-8C)alkenyl,

- 30 (2-8C)alkynyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulfinyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl and N,N-di-[(1-6C)alkyl]carbamoyl;

- (jjj) each R^2 , which may be the same or different, is selected from fluoro, chloro, bromo, iodo, cyano, hydroxy, amino, carbamoyl, trifluoromethyl, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (1-4C)alkoxy, (1-4C)alkylthio, (1-4C)alkylsulfinyl, (1-4C)alkylsulfonyl, (1-4C)alkylamino, di-[(1-4C)alkyl]amino, (1-4C)alkoxycarbonyl, N-(1-4C)alkylcarbamoyl, 5 N,N-di-[(1-4C)alkyl]carbamoyl;
- (kkk) each R^2 , which may be the same or different, is selected from fluoro, chloro, bromo, iodo, cyano, hydroxy, trifluoromethyl, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl and (1-4C)alkoxy;
- (lll) each R^2 , which may be the same or different, is selected from fluoro, chloro, bromo, 10 iodo, cyano, carbamoyl, hydroxy, trifluoromethyl, methyl, ethyl, isopropyl, methoxy, ethoxy, vinyl, allyl, ethynyl, 1-propynyl, 2-propynyl, N-methylcarbamoyl, N-ethylcarbamoyl and N,N-dimethylcarbamoyl;
- (mmm) each R^2 , which may be the same or different, is selected from fluoro, chloro, bromo, iodo, cyano, hydroxy, trifluoromethyl, methyl, ethyl, isopropyl, methoxy, ethoxy, 15 vinyl, allyl, ethynyl, 1-propynyl, and 2-propynyl;
- (nnn) each R^2 , which may be the same or different, is selected from fluoro, chloro, bromo, cyano, hydroxy, trifluoromethyl, methyl, ethyl, methoxy, ethoxy and ethynyl;
- (ooo) each R^2 , which may be the same or different, is selected from halogeno (particularly fluoro, chloro and bromo);
- 20 (ppp) a is 0, 1, 2 or 3, each R^2 is as defined in any of (iii) to (ooo) above and at least 1 R^2 is in an ortho position to the NH group;
- (qqq) a is 0, 1 or 2, each R^2 is as defined in any of (iii) to (ooo) above and wherein R^2 is not in an ortho position to the NH group;
- (rrr) a is 1, R^2 is in an ortho position to the NH group and is selected from halogeno 25 (particularly fluoro, chloro or bromo);
- (sss) a is 1, R^2 is in the para position to the NH group and is selected from halogeno (particularly fluoro, chloro or bromo);
- (ttt) a is 1, R^2 is in the 2-position on the anilino group and is selected from halogeno (particularly fluoro, chloro or bromo, more particularly fluoro), and Y is halogeno 30 (particularly fluoro, chloro or bromo);
- (uuu) a is 1, R^2 is in the para to the NH group and is selected from halogeno (particularly fluoro, chloro or bromo), and Y is halogeno (particularly fluoro, chloro or bromo);
- (vvv) a is 0 and Y is as defined in any of (ccc) to (hhh);

- (www) Y is selected from fluoro, chloro, bromo, iodo, methyl, ethyl, isopropyl, methoxy, ethoxy, vinyl, allyl, ethynyl, 1-propynyl and 2-propynyl, a is 0, 1, 2 or 3, and each R^2 , which may be the same or different, is selected from fluoro, chloro, bromo, iodo, cyano, trifluoromethyl, methyl, ethyl, isopropyl, methoxy, ethoxy, vinyl, allyl, ethynyl, 1-propynyl and 2-propynyl;
- (xxx) the aniline at the 4-position in the quinazoline ring in formula I is selected from 3-chloro-4-fluoroanilino, 3-chloro-2-fluoroanilino, 2-fluoro-5-chloroanilino, 3-bromoanilino, 3-methylanilino and 3-ethynylanilino;
- (yyy) the aniline at the 4-position in the quinazoline ring in formula I is 3-chloro-2-fluoroanilino; and
- (zzz) the aniline at the 4-position in the quinazoline ring in formula I is 3-chloro-4-fluoroanilino.

A particular embodiment of the present invention is a quinazoline derivative of the formula I wherein:

- m is 1 and R^1 is located at the 7-position and is selected from (1-4C)alkoxy, hydroxy-(1-4C)alkoxy and (1-4C)alkoxy(1-4C)alkoxy;
- X^1 is $C(R^3)_2$, wherein each R^3 , which may be the same or different, is selected from hydrogen and (1-4C)alkyl;
- Q^1 is selected from pyrrolidin-2-yl and pyrrolidin-3-yl,
- and wherein the group of the formula $ZC(O)$ is attached at the 1-position in Q^1 , and wherein Q^1 optionally bears one or more (for example 1, 2 or 3) substituents, which may be the same or different selected from fluoro, chloro, hydroxy, carbamoyl, (1-4C)alkyl, (1-4C)alkoxy, N-(1-4C)alkylcarbamoyl and N,N-di-[(1-4C)alkyl]carbamoyl,
- and wherein Q^1 optionally bears an oxo substituent;
- Z is selected from halogeno-(1-2C)alkyl, hydroxy-(1-2C)alkyl, (1-4C)alkoxy-(1-2C)alkyl, amino-(1-2C)alkyl, (1-4C)alkylamino-(1-2C)alkyl and di-[(1-4C)alkyl]amino-(1-2C)alkyl,
- or Z is Q^2 , wherein Q^2 is heterocyclyl or heterocyclyl-(1-2C)alkyl, and wherein the heterocyclyl in Q^2 is selected from azetidin-1-yl, pyrrolidin-1-yl, piperidino, piperazin-1-yl, morpholino, homopiperidin-1-yl and homopiperazin-1-yl,
- and wherein Q^2 is linked to the (1-2C)alkyl or C(O) group by a ring nitrogen,

and wherein any CH₂ or CH₃ group within a Z group, other than a CH₂ group within a heterocyclyl ring, optionally bears on each said CH₂ or CH₃ group one or more (1-4C)alkyl substituents or a substituent selected from cyano, carbamoyl, (2-6C)alkenyl, (2-6C)alkynyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl and (2-6C)alkanoyl,

5 and wherein any CH₂ which is attached to 2 carbon atoms or a CH₃ group which is attached to a carbon atom within a Z group, other than a CH₂ group within a heterocyclyl ring, optionally bears on each said CH₂ or CH₃ group one or more halogeno substituents or a substituent selected from hydroxy, amino, (1-6C)alkoxy, (1-6C)alkylsulfonyl, (1-6C)alkylamino and di-[(1-6C)alkyl]amino,

10 and wherein any heterocyclyl group within Z optionally bears one or more (for example 1, 2 or 3) substituents, which may be the same or different selected from halogeno, trifluoromethyl, cyano, hydroxy, amino, carbamoyl, (1-6C)alkyl, (1-6C)alkoxy, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl and (2-6C)alkanoyl, or from a group of the formula:

15 $-X^2-R^4$

wherein X² is a direct bond or is selected from O, CO and N(R⁵), wherein R⁵ is hydrogen or (1-4C)alkyl, and R⁴ is halogeno-(1-4C)alkyl, hydroxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, amino-(1-4C)alkyl, N-(1-4C)alkylamino-(1-4C)alkyl, N,N-di-[(1-4C)alkyl]amino-(1-4C)alkyl,

20 carbamoyl-(1-4C)alkyl, N-(1-4C)alkylcarbamoyl-(1-4C)alkyl and N,N-di-[(1-4C)alkyl]carbamoyl-(1-4C)alkyl,

and wherein any heterocyclyl group within Z optionally bears 1 or 2 oxo substituents;

Y is selected from halogeno, cyano, trifluoromethyl, (1-6C)alkyl, (1-6C)alkoxy and (2-6C)alkynyl;

25 each R², which may be the same or different, is selected from halogeno, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, trifluoromethyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulfinyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl and N,N-di-[(1-6C)alkyl]carbamoyl; and

30 a is 0, 1 or 2;

or a pharmaceutically acceptable salt thereof.

A further embodiment of the invention is a quinazoline derivative of the formula I wherein:

m is 1 and R¹ is located at the 7-position and is selected from (1-4C)alkoxy, hydroxy-(1-4C)alkoxy, methoxy-(1-4C)alkoxy and ethoxy-(1-4C)alkoxy (particularly R¹ is

5 (1-4C)alkoxy, more particularly methoxy);

Q¹-X¹ is selected from pyrrolidin-2-ylmethyl and pyrrolidin-3-ylmethyl,

and wherein Q¹ optionally bears one or more (for example 1, 2 or 3) substituents, which may be the same or different selected from fluoro, chloro, hydroxy, carbamoyl, (1-3C)alkyl, (1-3C)alkoxy, N-methylcarbamoyl and N,N-dimethylcarbamoyl,

10 and wherein Q¹ optionally bears an oxo substituent,

and wherein the group of the formula ZC(O) is attached at the 1-position in Q¹;

Z is selected from hydroxymethyl, methoxymethyl, ethoxymethyl, aminomethyl, (1-4C)alkylaminomethyl, N-[hydroxy-(2-4C)alkyl]-aminomethyl, N-[methoxy-(2-4C)alkyl]-aminomethyl, di-[(1-4C)alkyl]aminomethyl, N-[hydroxy-(2-4C)alkyl]-N-

15 [(1-4C)alkyl]aminomethyl, N,N-di-[hydroxy-(2-4C)alkyl]-aminomethyl, N-[methoxy-(2-4C)alkyl]-N-[(1-4C)alkyl]aminomethyl, pyrrolidin-1-ylmethyl, piperidinomethyl, piperazin-1-ylmethyl and morpholinomethyl,

and wherein any heterocyclyl group within Z optionally bears one or more (for example 1, 2 or 3) substituents, which may be the same or different selected from fluoro, chloro, cyano, hydroxy, amino, carbamoyl, (1-4C)alkyl, (1-4C)alkoxy, (1-4C)alkylamino, di-[(1-4C)alkyl]amino, N-(1-4C)alkylcarbamoyl, N,N-di-[(1-4C)alkyl]carbamoyl, acetyl, propionyl, 2-fluoroethyl, 2-hydroxyethyl, 2-methoxyethyl, cyanomethyl, hydroxyacetyl, aminoacetyl, methylaminoacetyl, ethylaminoacetyl, dimethylaminoacetyl, N-methyl-N-ethylaminoacetyl and fluoroacetyl,

25 and wherein any heterocyclyl group within Z optionally bears an oxo substituent;

Y is selected from fluoro, chloro, bromo, iodo, cyano, trifluoromethyl, methyl, ethyl, isopropyl, methoxy, ethoxy, vinyl, allyl, ethynyl, 1-propynyl and 2-propynyl;

each R², which may be the same or different, is selected from fluoro, chloro, bromo, iodo, cyano, hydroxy, trifluoromethyl, (1-4C)alkyl, (2-4C)alkynyl and (1-4C)alkoxy; and

30 a is 0, 1 or 2 (particularly 0 or 1);

or a pharmaceutically acceptable salt thereof.

A further embodiment of the invention is a quinazoline derivative of the formula I wherein:

- m** is 1 and **R**¹ is located at the 7-position and is selected from (1-4C)alkoxy, hydroxy-(1-4C)alkoxy, methoxy-(1-4C)alkoxy and ethoxy-(1-4C)alkoxy (particularly **R**¹ is (1-4C)alkoxy, more particularly methoxy);
- Q**¹-**X**¹ is selected from pyrrolidin-2-ylmethyl and pyrrolidin-3-ylmethyl, and wherein **Q**¹ optionally bears one or more (for example 1, 2 or 3) substituents, which may be the same or different selected from hydroxy and (1-3C)alkoxy, and wherein **Q**¹ optionally bears an oxo substituent,
- and wherein the group of the formula **ZC(O)** is attached at the 1-position in **Q**¹;
- Z** is selected from hydroxymethyl, methoxymethyl, ethoxymethyl, aminomethyl, (1-4C)alkylaminomethyl, N-[hydroxy-(2-4C)alkyl]-aminomethyl, N-[methoxy-(2-4C)alkyl]-aminomethyl, di-[(1-4C)alkyl]aminomethyl, N-[hydroxy-(2-4C)alkyl]-N-[(1-4C)alkyl]aminomethyl, N,N-di-[hydroxy-(2-4C)alkyl]-aminomethyl, N-[methoxy-(2-4C)alkyl]-N-[(1-4C)alkyl]aminomethyl, pyrrolidin-1-ylmethyl, piperidinomethyl, piperazin-1-ylmethyl and morpholinomethyl,
- and wherein any heterocyclyl group within **Z** optionally bears one or more (for example 1, 2 or 3) substituents, which may be the same or different selected from hydroxy, amino, carbamoyl, (1-4C)alkyl, (1-4C)alkoxy, (1-4C)alkylamino, di-[(1-4C)alkyl]amino, N-(1-4C)alkylcarbamoyl and N,N-di-[(1-4C)alkyl]carbamoyl
- and wherein any heterocyclyl group within **Z** optionally bears an oxo substituent;
- Y** is selected from fluoro, chloro, bromo, cyano, methyl, methoxy and ethynyl;
- each **R**², which may be the same or different, is selected from fluoro, chloro, bromo, cyano, hydroxy, trifluoromethyl, methyl, ethyl, methoxy, ethoxy and ethynyl; and
- a** is 0, 1 or 2 (particularly 0 or 1);
- or a pharmaceutically acceptable salt thereof.

A further embodiment of the invention is a quinazoline derivative of the formula I wherein:

- m** is 1 and **R**¹ is located at the 7-position and is selected from (1-4C)alkoxy, hydroxy-(1-4C)alkoxy, methoxy-(1-4C)alkoxy and ethoxy-(1-4C)alkoxy (particularly **R**¹ is (1-4C)alkoxy, 2-hydroxyethoxy or 2-methoxyethoxy more particularly **R**¹ is methoxy);
- Q**¹-**X**¹ is pyrrolidin-2-ylmethyl,

and wherein Q^1 optionally bears one or more (for example 1, 2 or 3) substituents,
which may be the same or different selected from hydroxy and (1-3C)alkoxy,

and wherein Q^1 optionally bears an oxo substituent,

and wherein the group of the formula $ZC(O)$ is attached at the 1-position in Q^1 ;

5 Z is selected from hydroxymethyl, methoxymethyl, aminomethyl, methylaminomethyl, di-methylaminomethyl, pyrrolidin-1-ylmethyl, piperidinomethyl and morpholinomethyl (particularly Z is selected from hydroxymethyl, aminomethyl, methylaminomethyl, di-methylaminomethyl and pyrrolidin-1-ylmethyl),

and wherein any heterocyclyl group within Z optionally bears 1 or 2 substituents,

10 which may be the same or different selected from hydroxy, amino, methyl, ethyl, methoxy, methylamino and di-methylamino;

Y is halogeno (particularly fluoro, chloro or bromo);

each R^2 , which may be the same or different, is selected from halogeno (particularly fluoro, chloro or bromo); and

15 a is 0, 1 or 2 (particularly 0 or 1);

or a pharmaceutically acceptable salt thereof.

A further embodiment of the invention is a quinazoline derivative of the formula I wherein:

m is 1 and R^1 is located at the 7-position and is selected from (1-4C)alkoxy, hydroxy-
20 (1-4C)alkoxy, (1-4C)alkoxy -(1-4C)alkoxy (particularly R^1 is (1-4C)alkoxy-(1-4C)alkoxy or (1-4C)alkoxy more particularly R^1 methoxy, 2-methoxyethoxy or 2-ethoxyethoxy);

Q^1-X^1 is selected pyrrolidin-2-ylmethyl,

and wherein Q^1 optionally bears one or more (for example 1, 2 or 3) substituents,
which may be the same or different selected from hydroxy, (1-3C)alkyl, (1-3C)alkoxy,
25 hydroxy-(1-3C)alkoxy and (1-3C)alkoxy- (1-3C)alkoxy;

and wherein Q^1 optionally bears an oxo substituent,

and wherein the group of the formula $ZC(O)$ is attached at the 1-position in Q^1 ;

Z is selected from hydroxymethyl, methoxymethyl, aminomethyl,
(1-4C)alkylaminomethyl, N-[hydroxy-(2-4C)alkyl]-aminomethyl, N-[methoxy-(2-4C)alkyl]-
30 aminomethyl, di-[(1-4C)alkyl]aminomethyl, N-[hydroxy-(2-4C)alkyl]-N-
[(1-4C)alkyl]aminomethyl, N,N-di-[hydroxy-(2-4C)alkyl]-aminomethyl, N-[methoxy-

(2-4C)alkyl]-N-[(1-4C)alkyl]aminomethyl, pyrrolidin-1-ylmethyl, piperidinomethyl, piperazin-1-ylmethyl and morpholinomethyl,

and wherein any heterocyclyl group within Z optionally bears one or more (for example 1, 2 or 3) substituents, which may be the same or different selected from halogeno for example fluoro or chloro), hydroxy, (1-4C)alkyl and (1-4C)alkoxy, and wherein any heterocyclyl group within Z optionally bears an oxo substituent;

Y is selected from fluoro, chloro, bromo, cyano, methyl, methoxy and ethynyl;

each R^2 , which may be the same or different, is selected from fluoro, chloro, bromo, cyano, hydroxy, trifluoromethyl, methyl, ethyl, methoxy, ethoxy and ethynyl; and

10 a is 0, 1 or 2 (particularly 0 or 1);

or a pharmaceutically acceptable salt thereof.

A further embodiment of the invention is a quinazoline derivative of the formula I wherein:

m is 1 and R^1 is located at the 7-position and is selected from (1-4C)alkoxy, 2-hydroxyethoxy and 2-methoxyethoxy (particularly R^1 is methoxy);

Q^1-X^1 is (2R)-pyrrolidin-2-ylmethyl,

and wherein the group of the formula $ZC(O)$ is attached at the 1-position of the pyrrolidinyl group,

Z is selected from hydroxymethyl, methoxymethyl, ethoxymethyl, aminomethyl, (1-4C)alkylaminomethyl, N-[hydroxy-(2-4C)alkyl]-aminomethyl, N-[methoxy-(2-4C)alkyl]-aminomethyl, di-[(1-4C)alkyl]aminomethyl, N-[hydroxy-(2-4C)alkyl]-N-[(1-4C)alkyl]aminomethyl, N,N-di-[hydroxy-(2-4C)alkyl]-aminomethyl, N-[methoxy-(2-4C)alkyl]-N-[(1-4C)alkyl]aminomethyl, pyrrolidin-1-ylmethyl, piperidinomethyl, piperazin-1-ylmethyl, morpholinomethyl, homopiperidin-1-ylmethyl and homopiperazin-1-ylmethyl,

and wherein any heterocyclyl group within Z optionally bears one or more (for example 1, 2 or 3) substituents, which may be the same or different selected from fluoro, chloro, hydroxy, (1-4C)alkyl and (1-4C)alkoxy,

and wherein any heterocyclyl group within Z optionally bears an oxo substituent;

30 Y is selected from fluoro, chloro, bromo, cyano, methyl, ethyl, isopropyl, methoxy, ethoxy, vinyl, allyl, ethynyl, 1-propynyl and 2-propynyl (particularly Y is selected from fluoro, chloro and bromo);

each R^2 , which may be the same or different, is selected from fluoro, chloro, bromo, cyano, hydroxy, trifluoromethyl, (1-4C)alkyl, (2-4C)alkynyl and (1-4C)alkoxy (particularly each R^2 , which may be the same or different is selected from fluoro, chloro and bromo); and

a is 0, 1 or 2 (particularly 0 or 1);

5 or a pharmaceutically acceptable salt thereof.

A further embodiment of the invention is a quinazoline derivative of the formula I wherein:

m is 1 and R^1 is located at the 7-position and is selected from (1-4C)alkoxy, 2-hydroxyethoxy and 2-methoxyethoxy (particularly R^1 is methoxy);

10 Q^1-X^1 is (2S)-pyrrolidin-2-ylmethyl,

and wherein the group of the formula ZC(O) is attached at the 1-position of the pyrrolidinyl group,

Z is selected from hydroxymethyl, methoxymethyl, ethoxymethyl, aminomethyl, (1-4C)alkylaminomethyl, N-[hydroxy-(2-4C)alkyl]-aminomethyl, N-[methoxy-(2-4C)alkyl]-aminomethyl, di-[(1-4C)alkyl]aminomethyl, N-[hydroxy-(2-4C)alkyl]-N-
15 [(1-4C)alkyl]aminomethyl, N,N-di-[hydroxy-(2-4C)alkyl]-aminomethyl, N-[methoxy-(2-4C)alkyl]-N-[(1-4C)alkyl]aminomethyl, pyrrolidin-1-ylmethyl, piperidinomethyl, piperazin-1-ylmethyl, morpholinomethyl, homopiperidin-1-ylmethyl and homopiperazin-1-ylmethyl,

20 and wherein any heterocyclyl group within Z optionally bears one or more (for example 1, 2 or 3) substituents, which may be the same or different selected from fluoro, chloro, hydroxy, (1-4C)alkyl and (1-4C)alkoxy,

and wherein any heterocyclyl group within Z optionally bears an oxo substituent;

Y is selected from fluoro, chloro, bromo, cyano, methyl, ethyl, isopropyl, methoxy, ethoxy, vinyl, allyl, ethynyl, 1-propynyl and 2-propynyl (particularly Y is selected from fluoro,
25 chloro and bromo);

each R^2 , which may be the same or different, is selected from fluoro, chloro, bromo, cyano, hydroxy, trifluoromethyl, (1-4C)alkyl, (2-4C)alkynyl and (1-4C)alkoxy (particularly each R^2 , which may be the same or different is selected from fluoro, chloro and bromo); and

30 a is 0, 1 or 2 (particularly 0 or 1);

or a pharmaceutically acceptable salt thereof.

A further embodiment of the invention is a quinazoline derivative of the formula I wherein:

m is 1 and **R¹** is located at the 7-position and is selected from (1-4C)alkoxy, 2-hydroxyethoxy and 2-methoxyethoxy (particularly **R¹** is methoxy);

5 **Q¹-X¹** is selected from (3R)-pyrrolidin-3-ylmethyl and (3S)-pyrrolidin-3-ylmethyl, and wherein the group of the formula **ZC(O)** is attached at the 1-position of the pyrrolidinyl group,

Z is selected from hydroxymethyl, methoxymethyl, ethoxymethyl, aminomethyl, (1-4C)alkylaminomethyl, N-[hydroxy-(2-4C)alkyl]-aminomethyl, N-[methoxy-(2-4C)alkyl]-aminomethyl, di-[(1-4C)alkyl]aminomethyl, N-[hydroxy-(2-4C)alkyl]-N-
10 [(1-4C)alkyl]aminomethyl, N,N-di-[hydroxy-(2-4C)alkyl]-aminomethyl, N-[methoxy-(2-4C)alkyl]-N-[(1-4C)alkyl]aminomethyl, pyrrolidin-1-ylmethyl, piperidinomethyl, piperazin-1-ylmethyl, morpholinomethyl, homopiperidin-1-ylmethyl and homopiperazin-1-ylmethyl,

15 and wherein any heterocyclyl group within **Z** optionally bears one or more (for example 1, 2 or 3) substituents, which may be the same or different selected from fluoro, chloro, hydroxy, (1-4C)alkyl and (1-4C)alkoxy,

and wherein any heterocyclyl group within **Z** optionally bears an oxo substituent;


Y is selected from fluoro, chloro, bromo, cyano, methyl, ethyl, isopropyl, methoxy, ethoxy, vinyl, allyl, ethynyl, 1-propynyl and 2-propynyl (particularly **Y** is selected from fluoro, chloro and bromo);

each **R²**, which may be the same or different, is selected from fluoro, chloro, bromo, cyano, hydroxy, trifluoromethyl, (1-4C)alkyl, (2-4C)alkynyl and (1-4C)alkoxy (particularly each **R²**, which may be the same or different is selected from fluoro, chloro and bromo); and

25 **a** is 0, 1 or 2 (particularly 0 or 1);

or a pharmaceutically acceptable salt thereof.

A further embodiment of the invention is a quinazoline derivative of the formula I of the formula Ia:



Ia


wherein:

m is 0 or m is 1 and R^{1a} is selected from (1-4C)alkoxy, hydroxy-(1-4C)alkoxy and
 5 (1-4C)alkoxy(1-4C)alkoxy;

R^{11} is selected from hydrogen, hydroxy, (1-4C)alkyl, (1-4C)alkoxy, hydroxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, hydroxy-(1-4C)alkoxy and (1-4C)alkoxy-(1-4C)alkoxy;

Z^1 is selected from (1-2C)alkyl, hydroxy-(1-2C)alkyl, (1-2C)alkoxy-(1-2C)alkyl,
 10 amino-(1-2C)alkyl, (1-4C)alkylamino-(1-2C)alkyl and di-[(1-4C)alkyl]amino-(1-2C)alkyl,
 or Z^1 is Q^2 , wherein Q^2 is pyrrolidinyl-(1-2C)alkyl (particularly pyrrolidin-1-yl-(1-2C)alkyl)),

and wherein any pyrrolidinyl group within Z^1 optionally bears one or more (for
 example 1, 2 or 3) substituents, which may be the same or different selected from fluoro,
 15 chloro, hydroxy, (1-3C)alkyl, (1-3C)alkoxy, or from a group of the formula:



wherein X^2 is a direct bond or is O, and R^4 is halogeno-(1-3C)alkyl,
 hydroxy-(1-3C)alkyl or (1-3C)alkoxy-(1-3C)alkyl,

and wherein any pyrrolidinyl group within Z^1 optionally bears an oxo substituent;

20 Y^1 is selected from fluoro, chloro, bromo, cyano, methyl, ethyl, isopropyl, methoxy, ethoxy, vinyl, allyl, ethynyl, 1-propynyl and 2-propynyl (particularly Y^1 is selected from fluoro, chloro, bromo and ethynyl);

each R^{2a} , which may be the same or different, is selected from fluoro, chloro, bromo, cyano, hydroxy, trifluoromethyl, (1-4C)alkyl, (2-4C)alkynyl and (1-4C)alkoxy (particularly
 25 each R^{2a} , which may be the same or different is selected from fluoro, chloro and bromo); and
 a is 0, 1 or 2 (particularly 0 or 1);

or a pharmaceutically acceptable salt thereof.

A further embodiment of the invention is a quinazoline derivative of the formula I of the formula Ia as hereinabove defined wherein Z^1 is selected from (1-2C)alkyl, hydroxy-(1-2C)alkyl, amino-(1-2C)alkyl, (1-4C)alkylamino-(1-2C)alkyl and di-[(1-4C)alkyl]amino-(1-2C)alkyl,

5 or Z^1 is Q^2 , wherein Q^2 is pyrrolidinyl-(1-2C)alkyl (particularly pyrrolidin-1-yl-(1-2C)alkyl)),

and wherein the pyrrolidinyl group within Z^1 optionally bears one or more (for example 1, 2 or 3) substituents, which may be the same or different selected from fluoro, chloro, hydroxy, (1-3C)alkyl, (1-3C)alkoxy, or from a group of the formula:

10 $-X^2-R^4$

wherein X^2 is a direct bond or is O, and R^4 is halogeno-(1-3C)alkyl, hydroxy-(1-3C)alkyl or (1-3C)alkoxy-(1-3C)alkyl,


and wherein any the pyrrolidinyl group within Z^1 optionally bears an oxo substituent; and wherein R^{1a} , m , R^{11} , Z^1 , Y^1 , R^{2a} and a are as defined above; or a pharmaceutically

15 acceptable salt thereof.

A further embodiment of the invention is a quinazoline derivative of the formula I of the formula Ia as hereinabove defined wherein m is 1 and R^{1a} is selected from (1-4C)alkoxy, hydroxy-(1-4C)alkoxy and (1-4C)alkoxy(1-4C)alkoxy; and R^{11} , Z^1 , Y^1 , R^{2a} and a are as defined above; or a pharmaceutically acceptable salt thereof.

20 A further embodiment of the invention is a quinazoline derivative of the formula I of the formula Ia as hereinabove defined wherein R^{11} is selected from hydrogen, hydroxy and (1-3C)alkoxy (particularly, R^{11} is selected from hydrogen, hydroxy and methoxy, more particularly R^{11} is hydrogen or hydroxy); and m , R^{1a} , Z^1 , Y^1 , R^{2a} and a are as defined above; or a pharmaceutically acceptable salt thereof.

25 A further embodiment of the invention is a quinazoline derivative of the formula I of the formula Ia as hereinabove defined wherein the aniline of the formula:



is selected from 3-chloro-4-fluoroanilino, 3-chloro-2-fluoroanilino, 2-fluoro-5-chloroanilino, 3-bromoanilino, 3-methylanilino and 3-ethynylanilino (particularly the aniline is 3-chloro-4-fluoroanilino or 3-chloro-2-fluoroanilino, more particularly 3-chloro-4-fluoroanilino); and m , Z^1 , R^{1a} and R^{11} are as defined above; or a pharmaceutically acceptable salt thereof.

A further embodiment of the invention is a quinazoline derivative of the formula I of the formula Ia as hereinabove defined wherein the pyrrolidin-2-ylmethoxy group at the 5-position in the quinazoline ring in formula Ia is the (2R)-pyrrolidin-2-ylmethoxy enantiomer and m , R^{1a} , R^{11} , Z^1 , Y^1 , R^{2a} and a are as defined above; or a pharmaceutically acceptable salt thereof.

A further embodiment of the invention is a quinazoline derivative of the formula I of the formula Ia as hereinabove defined wherein:

m is 1 and R^{1a} is selected from methoxy, ethoxy, 2-hydroxyethoxy, 2-methoxyethoxy and 2-ethoxyethoxy;

R^{11} is selected from hydrogen, hydroxy or methoxy (particularly hydrogen or hydroxy);

Z^1 is selected from methyl, hydroxymethyl, aminomethyl, methylaminomethyl, dimethylaminomethyl and pyrrolidin-1-ylmethyl (particularly Z^1 is selected from methyl, hydroxymethyl, dimethylaminomethyl and pyrrolidin-1-ylmethyl);

Y^1 is selected from fluoro, chloro, bromo and ethynyl (particularly Y^1 is fluoro or chloro);

R^{2a} is selected from fluoro, chloro and bromo (particularly R^{2a} is fluoro or chloro); and

a is 0 or 1;

or a pharmaceutically acceptable salt thereof.

In this embodiment a preferred aniline at the 4-position in the quinazoline ring in formula Ia is 3-chloro-4-fluoroanilino or 3-chloro-2-fluoroanilino (particularly 3-chloro-4-fluoroanilino).

A further embodiment of the invention is a quinazoline derivative of the formula I of the formula Ia as hereinabove defined wherein:

m is 1 and R^{1a} is selected from methoxy, ethoxy, 2-hydroxyethoxy, 2-methoxyethoxy and 2-ethoxyethoxy (particularly R^{1a} is selected from methoxy, and 2-ethoxyethoxy);

R^{11} is selected from hydrogen or hydroxy;

Z^1 is selected from methyl or hydroxymethyl;

Y^1 is selected from fluoro, chloro, bromo and ethynyl (particularly Y^1 is fluoro or chloro);

R^{2a} is selected from fluoro, chloro and bromo (particularly R^{2a} is fluoro or chloro); and

a is 0 or 1;

or a pharmaceutically acceptable salt thereof.

In this embodiment a preferred aniline at the 4-position in formula Ia is 3-chloro-4-
5 fluoroanilino or 3-chloro-2-fluoroanilino (particularly 3-chloro-4-fluoroanilino).

A particular compound of the invention is, for example, a quinazoline derivative of the formula I selected from:

2-((2*R*)-2-(((4-[3-Chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl)pyrrolidin-1-yl)-2-oxoethanol;

10 *N*-(3-chloro-4-fluorophenyl)-5-(((2*R*)-1-((dimethylamino)acetyl)pyrrolidin-2-yl)methoxy)-7-methoxyquinazolin-4-amine;

N-(3-chloro-4-fluorophenyl)-7-methoxy-5-(((2*S*)-1-(pyrrolidin-1-ylacetyl)pyrrolidin-2-yl)methoxy)quinazolin-4-amine;

(3*S*,5*S*)-5-(((4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl)-1-(*N,N*-
15 dimethylglycyl)pyrrolidin-3-ol;

(3*S*,5*R*)-5-(((4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl)-1-glycoloylpyrrolidin-3-ol;

(3*R*,5*R*)-5-(((4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl)-1-glycoloylpyrrolidin-3-ol;

20 (3*R*,5*R*)-5-(((4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl)-1-(*N,N*-dimethylglycyl)pyrrolidin-3-ol;

(3*S*,5*R*)-1-acetyl-5-(((4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl)pyrrolidin-3-ol;

2-((2*R*)-2-(((4-[3-chloro-4-fluoroanilino]-7-(2-hydroxyethoxy)quinazolin-5-
25 yl)oxy)methyl)pyrrolidin-1-yl)-2-oxoethanol;

2-((2*R*)-2-(((4-[3-chloro-4-fluoroanilino]-7-(2-ethoxyethoxy)quinazolin-5-yl)oxy)methyl)pyrrolidin-1-yl)-2-oxoethanol;

2-((2*R*)-2-(((4-[3-chloro-4-fluoroanilino]-7-ethoxyquinazolin-5-yl)oxy)methyl)-pyrrolidin-1-yl)-2-oxoethanol; and

30 *N*-(3-chloro-4-fluorophenyl)-7-methoxy-5-(((2*R*)-1-(methoxyacetyl)pyrrolidin-2-yl)methoxy)quinazolin-4-amine;

or a pharmaceutically acceptable salt thereof.

A further particular compound of the invention is, for example, a quinazoline derivative of the formula I selected from:

2-((2*R*)-2-(((4-(3-chloro-4-fluoroanilino)-7-methoxyquinazolin-5-yl)oxy)methyl)pyrrolidin-1-yl)-2-oxoethanol;

5 *N*-(3-chloro-4-fluorophenyl)-5-(((2*R*)-1-((dimethylamino)acetyl)pyrrolidin-2-yl)methoxy)-7-methoxyquinazolin-4-amine; and


N-(3-chloro-4-fluorophenyl)-7-methoxy-5-(((2*S*)-1-(pyrrolidin-1-ylacetyl)pyrrolidin-2-yl)methoxy)quinazolin-4-amine;

or a pharmaceutically acceptable salt thereof.

10 A quinazoline derivative of the formula I, or a pharmaceutically-acceptable salt thereof, may be prepared by any process known to be applicable to the preparation of chemically-related compounds. Suitable processes include, for example, those illustrated in European Patent Applications Nos. 0520722, 0566226, 0602851, 0635507 and 0635498, and International Patent Applications WO 96/15118, WO 96/16960 and WO01/94341. Such
15 processes, when used to prepare a quinazoline derivative of the formula I are provided as a further feature of the invention and are illustrated by the following representative process variants in which, unless otherwise stated, R^1 , R^2 , X^1 , Y, Q^1 , Z a and m have any of the meanings defined hereinbefore. Necessary starting materials may be obtained by standard procedures of organic chemistry. The preparation of such starting materials is described in
20 conjunction with the following representative process variants and within the accompanying Examples. Alternatively necessary starting materials are obtainable by analogous procedures to those illustrated which are within the ordinary skill of an organic chemist.

Process (a) The coupling, conveniently in the presence of a suitable base, of a quinazoline of the formula II:

25



II


wherein R^1 , R^2 , X^1 , Y, Q^1 , a and m have any of the meanings defined hereinbefore

except that any functional group is protected if necessary, with a carboxylic acid of the formula Z-COOH, or a reactive derivative thereof, wherein Z has any of the meanings defined hereinbefore except that any functional group is protected if necessary;

or

- 5 Process (b): for the preparation of those compounds of the formula I wherein Z is selected from amino-(1-4C)alkyl, (1-4C)alkylamino-(1-4C)alkyl, di-[(1-4C)alkyl]amino-(1-4C)alkyl, or Z is Q², wherein Q² is heterocyclyl-(1-4C)alkyl, and wherein the heterocyclyl in Q² contains at least 1 nitrogen heteroatom, and optionally 1 or 2 further heteroatoms selected from oxygen, nitrogen and sulfur, and wherein Q² is attached by a ring nitrogen atom,

- 10 the reaction of a quinazoline of the formula III:




III

- 15 wherein R¹, R², X¹, Y, Q¹, a and m have any of the meanings defined hereinbefore except that any functional group is protected if necessary, Z¹ is a direct bond or (1-4C)alkyl and L¹ is a displaceable group, with an amine or a compound of the formula Q²H wherein Q² has any of the meanings defined hereinbefore except that any functional group is protected if necessary;


- 20 or

Process (c): for the preparation of those compounds of the formula I wherein R¹ is a hydroxy group, the cleavage of a quinazoline derivative of the formula I wherein R¹ is a (1-6C)alkoxy; or

- Process (d): for the preparation of those compounds of the formula I wherein m is 1 and R¹ is optionally substituted (1-6C)alkoxy, optionally substituted (2-6C)alkenyloxy, optionally substituted (2-6C)alkynyloxy or a group of the formula :



wherein X^3 is O, and Q^3 is as hereinbefore defined except that any functional group is protected if necessary, and the optional substituents on R^1 are those that may be present on R^1 as hereinbefore defined in relation to formula I, the reaction, conveniently in the presence of a suitable base, of a quinazoline of the formula IV:




IV

wherein R^2 , X^1 , Y , Q^1 , Z and a have any of the meanings defined hereinbefore except that any functional group is protected if necessary, with a compound of the formula $R^{1a}-L^1$ or Q^3-L^1 wherein Q^3 is as hereinbefore defined except any functional group is protected if necessary, R^{1a} is selected from optionally substituted (1-6C)alkyl, optionally substituted (2-6C)alkenyl and optionally substituted (2-6C)alkynyl and L^1 is a displaceable group, and wherein the optional substituents on R^{1a} are the same as those that may be present on the corresponding groups on R^1 as hereinbefore defined in relation to formula I ;

or

- 15 Process (e): for the preparation of those compounds of the formula I wherein m is 1 and R^1 is optionally substituted (1-6C)alkoxy, optionally substituted (2-6C)alkenyloxy, optionally substituted (2-6C)alkynyloxy or a group of the formula :



wherein X^3 is O, and Q^3 is as hereinbefore defined and the optional substituents on R^1 are those that may be present on R^1 as hereinbefore defined in relation to formula I, the coupling of the a quinazoline of the formula IV as hereinbefore defined with an alcohol of the formula $R^{1a}OH$ or Q^3OH wherein Q^3 is as hereinbefore defined except any functional group is protected if necessary and R^{1a} is as hereinbefore defined in process (d);

and thereafter, if necessary:

- 25 (i) converting a quinazoline derivative of the formula I into another quinazoline derivative of the formula I;
- (ii) removing any protecting group that is present by conventional means;
- (iii) forming a pharmaceutically acceptable salt.

Specific conditions for the above reactions are as follows:

Process (a)

The coupling reaction is conveniently carried out in the presence of a suitable coupling agent, such as a carbodiimide, or a suitable peptide coupling agent, for example O-(7-
5 azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate or 4-(4,6-dimethoxy[1.3.5]triazin-2-yl)-4-methylmorpholinium chloride, or a carbodiimide such as dicyclohexylcarbodiimide, optionally in the presence of a catalyst such as dimethylaminopyridine or 4-pyrrolidinopyridine.


The coupling reaction is conveniently carried out in the presence of a suitable base. A
10 suitable base is, for example, an organic amine base such as, for example, pyridine, 2,6-lutidine, collidine, 4-dimethylaminopyridine, triethylamine, di-isopropylethylamine, N-methylmorpholine or diazabicyclo[5.4.0]undec-7-ene, or, for example, an alkali or alkaline earth metal carbonate, for example sodium carbonate, potassium carbonate, cesium carbonate, calcium carbonate, or, for example, an alkali metal hydride, for example sodium hydride.

15 The reaction is conveniently carried out in the presence of a suitable inert solvent or diluent, for example an alcohol or ester such as methanol, ethanol, isopropanol or ethyl acetate, a halogenated solvent such as methylene chloride, chloroform or carbon tetrachloride, an ether such as tetrahydrofuran or 1,4-dioxan, an aromatic solvent such as toluene, or a dipolar aprotic solvent such as N,N-dimethylformamide, N,N-dimethylacetamide,
20 N-methylpyrrolidin-2-one or dimethylsulfoxide. The reaction is conveniently carried out at a temperature in the range, for example, from 0 to 120°C, particularly at or near ambient temperature.

By the term "reactive derivative" of the carboxylic acid of the formula Z-COOH is meant a carboxylic acid derivative that will react with the quinazoline of formula II to give
25 the corresponding amide. A suitable reactive derivative of a carboxylic acid of the formula ZCCOH is, for example, an acyl halide, for example an acyl chloride formed by the reaction of the acid and an inorganic acid chloride, for example thionyl chloride; a mixed anhydride, for example an anhydride formed by the reaction of the acid and a chloroformate such as isobutyl chloroformate; an active ester, for example an ester formed by the reaction of the acid
30 and a phenol such as pentafluorophenol, an ester such as pentafluorophenyl trifluoroacetate, an alcohol such as methanol, ethanol, isopropanol, butanol or N-hydroxybenzotriazole, or an active ester formed by the reaction of the acid and an activated triazine such as 4-(4,6-

dimethoxy[1.3.5]triazin-2-yl)-4-methylmorpholinium chloride hydrate; or an acyl azide, for example an azide formed by the reaction of the acid and azide such as diphenylphosphoryl azide; an acyl cyanide, for example a cyanide formed by the reaction of an acid and a cyanide such as diethylphosphoryl cyanide. The reaction of these reactive derivatives of carboxylic acid with amines (such as a compound of the formula II) is well known in the art, for Example they may be reacted in the presence of a base, such as those described above, and in a suitable solvent, such as those described above. The reaction may conveniently be performed at a temperature as described above.

The quinazoline of the formula II may be obtained by conventional procedures. For example, a quinazoline derivative of the formula II wherein m is 1 and R¹ is optionally substituted (1-6C)alkoxy and is located at the 7-position may be prepared according to Reaction Scheme 1:



15

Reaction Scheme 1

Notes for Reaction Scheme 1:

L is a suitable displaceable group;

R is optionally substituted (1-6C)alkyl;

Pg is a protecting group; and

R², X¹, Y, Q¹, Z and a have any of the meanings defined hereinbefore, except any functional group is protected if necessary and any protecting groups present are removed either following
5 a particular reaction step or at the end of Reaction Scheme 1 to give the compound of formula II.

Step (i)

A suitable displaceable group represented by L includes, for example a halogeno or a sulfonyloxy group, for example fluoro, chloro, methylsulfonyloxy or toluene-4-sulfonyloxy
10 group. A particular group L is fluoro or chloro, more particularly fluoro.

The halogenation reaction (chlorination shown) may be carried out using a suitable chlorinating agent such as thionyl chloride (as shown), phosphoryl chloride or a mixture of carbon tetrachloride and triphenylphosphine.

The halogenation reaction is conveniently carried out in a suitable solvent, for example
15 1,2-dichloroethane or N,N-dimethylformamide conveniently in the presence of an base such as an organic base, for example di-isopropylethylamine. The reaction is conveniently carried out at a temperature in the range, for example, 0 to 150°C, preferably at or near the reflux temperature.

Compounds of the formula IIa are commercially available compounds or they are
20 known in the literature, or they can be prepared by standard processes known in the art. For example when each L in the compound of formula IIa is fluoro, the 5,7-difluoro-3,4-dihydroquinazolin-4-one starting material may be prepared using the method described in WO01/94341, example 4, footnote [5].

Step (ii):

25 The reaction is conveniently carried out in the presence of an acid. Suitable acids include, for example hydrogen chloride gas (conveniently dissolved in diethyl ether or dioxane) or hydrochloric acid.

Alternatively, the reaction may be carried out in the presence of a suitable base. A suitable base is, for example, an organic amine base such as, for example, one of the bases
30 described above in relation to process (a).

Alternatively the quinazoline derivative may be reacted with the aniline in the absence of an acid or a base. In this reaction displacement of the chloro leaving group results in the formation of the hydrochloric acid in-situ and the autocatalysis of the reaction.

The reaction in step (ii) is conveniently carried out in the presence of a suitable inert solvent or diluent, for example an alcohol or ester such as methanol, ethanol, isopropanol or ethyl acetate, a halogenated solvent such as methylene chloride, chloroform or carbon tetrachloride, an ether such as tetrahydrofuran or 1,4-dioxan, an aromatic solvent such as toluene, or a dipolar aprotic solvent such as N,N-dimethylformamide, N,N-dimethylacetamide, N-methylpyrrolidin-2-one or dimethylsulfoxide. The above reaction is conveniently carried out at a temperature in the range, for example, 0 to 250°C, conveniently in the range 40 to 80°C or, preferably, at or near the reflux temperature of the solvent when used. Conveniently, the above reaction may be performed in the presence of a suitable catalyst, for example a crown ether.

The anilines used as a starting material in step (ii) are commercially available compounds or they are known in the literature, or they can be prepared by standard processes known in the art.

Step (iii)

A suitable base for the reaction the quinazoline and the alcohol of the formula ROH includes, for example a strong non-nucleophilic base such as an alkali metal hydride, for example sodium hydride, an alkali metal amide, for example lithium di-isopropylamide (LDA) or an alkali metal alkoxide such as potassium tert-butoxide or sodium tert-butoxide.

The reaction is conveniently carried out in the presence of a suitable inert solvent or diluent, for example a halogenated solvent such as methylene chloride, chloroform or carbon tetrachloride, an ether such as tetrahydrofuran or 1,4-dioxane, an aromatic solvent such as toluene, or a dipolar aprotic solvent such as N,N-dimethylformamide, N,N-dimethylacetamide, N-methylpyrrolidin-2-one or dimethylsulfoxide. The reaction is conveniently carried out at a temperature in the range of, for example, from 10 to 250°C, preferably in the range of from 40 to 150°C.


Step (iv)

Cleavage of the (1-6C)alkoxy group may be performed using an analogous procedure to that described for process (c) below, for example by reaction with pyridine hydrochloride as shown in Reaction Scheme 1 .

Step (v)

- Coupling reaction under Mitsunobu conditions. The coupling of alcohols using the Mitsunobu coupling reaction is well known in the art and is discussed in detail in, for example, Tet. Letts., 31, 699, (1990); The Mitsunobu Reaction, D.L.Hughes, Organic Reactions, 1992, Vol.42, 335-656 and Progress in the Mitsunobu Reaction, D.L.Hughes, Organic Preparations and Procedures International, 1996, Vol.28, 127-164. For example suitable Mitsunobu conditions include, reaction in the presence of a suitable tertiary phosphine and a di-alkylazodicarboxylate, conveniently in the presence of a suitable organic solvent. A suitable tertiary phosphine includes for example tri-n-butylphosphine or tri-phenylphosphine (as shown). A suitable di-alkylazodicarboxylate includes for example diethyl azodicarboxylate (DEAD) or di-tert-butyl azodicarboxylate (DTAD, as shown in Reaction Scheme 1). Conveniently the reaction is carried out in an organic solvent, for example a halogenated solvent such as methylene chloride, or an ether such as tetrahydrofuran. The reaction is suitably carried out in the temperature range of from 0°C to 60°C, suitably at ambient temperature.
- Alcohols of the formula HQ^1-X^1-OH used as the starting material in step (ii) (prior to protecting with the protecting group Pg) are commercially available compounds or they are known in the literature, or they can be prepared by standard processes known in the art. The protecting group Pg is present on the NH group in Q^1 to which the group of the formula $ZC(O)$ is attached in the quinazoline of the formula I. The protecting group is a suitable amine protecting group as described herein, for example a tert-butoxycarbonyl protecting group (BOC). When Q^1 carries other functional groups, for example a hydroxy group, it may be necessary to protect such groups with additional suitable protecting groups for such functional groups as described herein. For example if additional hydroxy groups are present, a suitable hydroxy protecting group is tert-butyl(dimethyl)silyloxy group. The protecting groups present may be removed following the Mitsunobu coupling, for example by treatment with a suitable acid such as trifluoroacetic acid.

Compounds of the formula II wherein R^1 is at the 7-position may also be prepared according to Reaction Scheme 1a:



Reaction Scheme 1a

Notes for Reaction Scheme 1a:

L^1 is a suitable displaceable group as defined hereinafter in relation to Process (d);

5 Pg is a suitable amine protecting group as defined hereinbefore in relation to Reaction Scheme 1;

Pg^1 is a suitable hydroxy protecting group; and

R^1 , R^2 , X^1 , Y , Q^1 , Z and a have any of the meanings defined hereinbefore, except any functional group is protected if necessary and any protecting groups present are removed
 10 either following a particular reaction step or at the end of Reaction Scheme 1a to give the compound of formula **II**.

Step (i)

Pg^1 is a suitable hydroxy protecting group, for example an aryl-lower alkyl group such as benzyl. The protecting group Pg^1 may be removed using standard methods well known in
 15 the art. For example, when Pg^1 is benzyl, the protecting group may be removed by hydrogenation. Suitable hydrogenation conditions are well known, for example by hydrogenation in the presence of a palladium catalyst.

Compounds of formula **IIb** may be prepared by standard processes known in the art, for example using the process described in Reaction Scheme 3.

20 Step (ii)

Carried out under analogous conditions to those used in Process (d) described herein.

Step (iii)

The protecting group Pg may be removed using standard conditions, for example when Pg is *tert*-butoxycarbonyl, be treatment with trifluoroacetic acid.

- 5 Compounds of the formula **II** wherein R¹ is not at the 7-position, may be prepared using analogous methods to those described above in Reaction Schemes 1 and 1a using appropriate starting materials.

Process (b)


- Suitable displaceable groups represented by L¹ in the quinazoline of formula **II** include
- 10 halogeno, or a sulfonyloxy group, for example chloro or bromo, methane sulfonyloxy or toluenesulfonyloxy, particularly halogeno such as chloro. The reaction is suitably performed in the presence of a suitable inert solvent or diluent, for example a solvent(s) or diluent(s) described above in relation to process (a) such as a chlorinated solvent, for example dichloromethane or a dipolar aprotic solvent such as N,N-dimethylformamide. Suitably the
- 15 reaction is carried out at a temperature of, for example 0 to 180°C, particularly 20°C to the reflux temperature of the solvent/diluent.

The reaction is conveniently carried out in the presence of a base. Suitable bases include, for example, those described above in relation to process (a) such as a carbonate, for example cesium carbonate, or a tertiary amine, for example diisopropylethylamine.

- 20 Alternatively, an excess of the amine or the compound of the formula Q²H may be used to provide the required basic conditions.

The amines and compounds of the formula Q²H used in process (b) are commercially available compounds or they are known in the literature, or they can be prepared by standard processes known in the art.

- 25 The quinazoline of the formula **III** may be obtained by conventional procedures. For example a quinazoline of the formula **III** wherein m is 1, R¹ is (1-6C)alkoxy and is located at the 7-position may be prepared as described in Reaction Scheme 2:



Reaction Scheme 2

Notes for Reaction Scheme 2:

L^1 and L^2 are displaceable groups;

5 R is (1-6C)alkyl; and

R^2 , X^1 , Y, Q^1 , Z^1 and a have any of the meanings defined hereinbefore, except any functional group is protected if necessary and any protecting groups present are removed following the reaction to give the compound of formula III.

Suitable displaceable groups represented by L^1 and L^2 are as hereinbefore described
 10 for L^1 in process (b), provided that L^2 is more labile than L^1 . Suitably, L^1 and L^2 are both halogeno, for example both are chloro.

The reaction is conveniently carried out in the presence of a base. Suitable bases include, for example those described above in relation to process (a), such as an organic amine, for example di-isopropylethylamine. The reaction is suitably performed in the
 15 presence of a suitable inert solvent or diluent, for example a solvent(s) or diluent(s) described above in relation to process (a) such as a methylene chloride. Suitably the reaction is carried out at a temperature of, for example 0 to 180°C, particularly 20°C to the reflux temperature of the solvent/diluent.

Compound of the formula $\text{L}^1\text{Z}^1\text{C(O)L}^2$ used as the starting material in Reaction
 20 Scheme 2 are commercially available compounds or they are known in the literature, or they can be prepared by standard processes known in the art. The compound of the formula II used as a starting material in Reaction Scheme 2 may be prepared using conventional methods, for example the process described in Reaction Scheme 1.

In a variant to Process (b) the reaction may be telescoped starting from the compound
 25 of formula II by reacting the compound of the formula $\text{L}^1\text{Z}^1\text{C(O)L}^2$ and a compound of formula II followed by reaction with a suitable amine or the compound of the formula Q^2H as

described above. Such a process avoids the need to isolate the intermediate compound of the formula **III**.

Process (c)

The cleavage reaction may conveniently be carried out by any of the many procedures known for such a transformation. The cleavage reaction of a compound of the formula **I** wherein R^1 is a (1-6C)alkoxy group may be carried out, for example, by treatment of the quinazoline derivative with an alkali metal (1-6C)alkylsulfide such as sodium ethanethiolate or, for example, by treatment with an alkali metal diarylphosphide such as lithium diphenylphosphide. Alternatively the cleavage reaction may conveniently be carried out, for example, by treatment of the quinazoline derivative with a boron or aluminium trihalide such as boron tribromide, or by reaction with an organic or inorganic acid, for example trifluoroacetic acid. Such reactions are suitably carried out in the presence of a suitable inert solvent or diluent as defined hereinbefore. A preferred cleavage reaction is the treatment of a quinazoline derivative of the formula **I** with pyridine hydrochloride. The cleavage reactions are suitably carried out at a temperature in the range, for example, from 10 to 150°C, for example from 25 to 80°C.


Process (d)

Suitable displaceable groups represented by L^1 are as hereinbefore defined in relation to process (b), for example L^1 is halogeno such as chloro or bromo. Suitable bases are as hereinbefore defined in relation to process (a), for example an alkali or alkaline earth metal carbonate, for example sodium carbonate, potassium carbonate, cesium carbonate, calcium carbonate. The reaction is suitably carried out under analogous conditions and in the presence of the solvents described above for process (b).

The compounds of the formulae $R^{1a}-L^1$ and Q^3-L^1 used as the starting materials in Process (d) are commercially available compounds or they are known in the literature, or they can be prepared by standard processes known in the art.

The compound of formula **IV** used in process (d) may be prepared using conventional techniques, for example by cleavage of an alkoxy group from another compound of formula **I** using process (c).

Alternatively, the compound of formula **IV** may be prepared as described in Reaction Scheme 3:



Reaction Scheme 3

Notes for Reaction Scheme 3

5 Pg and Pg^2 are suitable amine protecting groups;

Pg^1 is a suitable hydroxy protecting group;

L^2 is a suitable displaceable group, for example halogeno such as chloro; and

R^1 , R^2 , X^1 , Y , Q^1 , Z and a have any of the meanings defined hereinbefore, except any functional group is protected if necessary and any protecting groups present are removed either following a particular reaction step or at the end of Reaction Scheme 3 to give the compound of formula IV.

- 5 Step (i): The reaction is carried out under Mitsunobu conditions as described herein in relation to Reaction Scheme 1 (step (v)).

Suitable hydroxy protecting groups represented by Pg^1 are as defined above in relation to Reaction Scheme 1a, for example benzyl.

- 10 Suitable amino protecting groups represented by Pg^2 are as well known in the art and include for example lower alkanoyloxyalkyl groups such as pivaloyloxymethyl. Suitable amino protecting groups represented by Pg are as hereinbefore defined, for example tert-butoxycarbonyl.

- Compounds of formula IVa used as the starting material in Reaction Scheme 3 are known and can be prepared using conventional methods, for example as described in Example 15 [8] of WO01/94341.

Step (ii): The protecting group Pg^2 may be removed using standard conditions. For example when Pg^2 is a lower alkanoyloxyalkyl group, by treatment with methanolic ammonia.

- Step (iii): Halogenation under analogous conditions to those used in step (i) of Reaction Scheme 1. Alternative halogenation agents to the $POCl_3$ shown in Reaction Scheme 3 may be 20 used if desired. Such halogenation agents are well known and include, for example thionyl chloride.

Step (iv): Aniline coupling using analogous conditions to those used in Step (ii) of Reaction Scheme 1.

- Step (v): Deprotection to remove Pg and Pg^1 using standard techniques well known in the art. 25 For example by treatment with trifluoroacetic acid.

Step (vi): Analogous conditions to those used in Reaction Scheme 2. Compounds of the formula $ZC(O)L^2$ are commercially available compounds or they are known in the literature, or they can be prepared by standard processes known in the art.

Process (e)

- 30 Suitably the coupling reaction is carried out under Mitsunobu conditions using an analogous procedure to that described above in step (iv) of Reaction Scheme 1.

The compounds of the formulae $R^{1a}OH$ and Q^3OH used as the starting materials in Process (e) are commercially available compounds or they are known in the literature, or they

can be prepared by standard processes known in the art.

The quinazoline derivative of the formula I may be obtained from the above processes in the form of the free base or alternatively it may be obtained in the form of a salt, for example in process (b) an acid addition salt may be produced with the acid of the formula
5 H-L¹ when L¹ is, for example halogeno such as chloro. When it is desired to obtain the free base from a salt of the compound of formula I, the salt may be treated with a suitable base, for example, an alkali or alkaline earth metal carbonate or hydroxide, for example sodium carbonate, potassium carbonate, calcium carbonate, sodium hydroxide or potassium hydroxide, or by treatment with ammonia for example using a methanolic ammonia solution
10 such as 7N ammonia in methanol.

The protecting groups used in the processes above may in general be chosen from any of the groups described in the literature or known to the skilled chemist as appropriate for the protection of the group in question and may be introduced by conventional methods. Protecting groups may be removed by any convenient method as described in the literature or
15 known to the skilled chemist as appropriate for the removal of the protecting group in question, such methods being chosen so as to effect removal of the protecting group with minimum disturbance of groups elsewhere in the molecule.

Specific examples of protecting groups are given below for the sake of convenience, in which "lower", as in, for example, lower alkyl, signifies that the group to which it is applied
20 preferably has 1-4 carbon atoms. It will be understood that these examples are not exhaustive. Where specific examples of methods for the removal of protecting groups are given below these are similarly not exhaustive. The use of protecting groups and methods of deprotection not specifically mentioned are, of course, within the scope of the invention.

A carboxy protecting group may be the residue of an ester-forming aliphatic or
25 arylaliphatic alcohol or of an ester-forming silanol (the said alcohol or silanol preferably containing 1-20 carbon atoms). Examples of carboxy protecting groups include straight or branched chain (1-12C)alkyl groups (for example isopropyl, and tert-butyl); lower alkoxy- lower alkyl groups (for example methoxymethyl, ethoxymethyl and isobutoxymethyl); lower acyloxy-lower alkyl groups, (for example acetoxymethyl, propionyloxymethyl, butyryloxymethyl and pivaloyloxymethyl); lower
30 alkoxycarbonyloxy-lower alkyl groups (for example 1-methoxycarbonyloxyethyl and 1-ethoxycarbonyloxyethyl); aryl-lower alkyl groups (for example benzyl, 4-methoxybenzyl, 2-nitrobenzyl, 4-nitrobenzyl, benzhydryl and phthalidyl); tri(lower alkyl)silyl groups (for

example trimethylsilyl and tert-butyldimethylsilyl); tri(lower alkyl)silyl-lower alkyl groups (for example trimethylsilylethyl); and (2-6C)alkenyl groups (for example allyl). Methods particularly appropriate for the removal of carboxyl protecting groups include for example acid-, base-, metal- or enzymically-catalysed cleavage.

- 5 Examples of hydroxy protecting groups include lower alkyl groups (for example tert-butyl), lower alkenyl groups (for example allyl); lower alkanoyl groups (for example acetyl); lower alkoxycarbonyl groups (for example tert-butoxycarbonyl); lower alkenyloxycarbonyl groups (for example allyloxycarbonyl); aryl-lower alkoxycarbonyl groups (for example benzyloxycarbonyl, 4-methoxybenzyloxycarbonyl, 10 2-nitrobenzyloxycarbonyl and 4-nitrobenzyloxycarbonyl); tri(lower alkyl)silyl (for example trimethylsilyl and tert-butyldimethylsilyl) and aryl-lower alkyl (for example benzyl) groups.

- Examples of amino protecting groups include formyl, aryl-lower alkyl groups (for example benzyl and substituted benzyl, 4-methoxybenzyl, 2-nitrobenzyl and 2,4-dimethoxybenzyl, and triphenylmethyl); di-4-anisylmethyl and furylmethyl groups; lower 15 alkoxycarbonyl (for example tert-butoxycarbonyl); lower alkenyloxycarbonyl (for example allyloxycarbonyl); aryl-lower alkoxycarbonyl groups (for example benzyloxycarbonyl, 4-methoxybenzyloxycarbonyl, 2-nitrobenzyloxycarbonyl and 4-nitrobenzyloxycarbonyl); lower alkanoyloxyalkyl groups (for example pivaloyloxymethyl); trialkylsilyl (for example trimethylsilyl and tert-butyldimethylsilyl); alkylidene (for example methyldiene) and 20 benzyldiene and substituted benzyldiene groups.

- Methods appropriate for removal of hydroxy and amino protecting groups include, for example, acid-, base-, metal- or enzymically-catalysed hydrolysis for groups such as 2-nitrobenzyloxycarbonyl, hydrogenation for groups such as benzyl and photolytically for groups such as 2-nitrobenzyloxycarbonyl. For example a tert butoxycarbonyl protecting group 25 may be removed from an amino group by an acid catalysed hydrolysis using trifluoroacetic acid.

- The reader is referred to Advanced Organic Chemistry, 4th Edition, by J. March, published by John Wiley & Sons 1992, for general guidance on reaction conditions and reagents and to Protective Groups in Organic Synthesis, 2nd Edition, by T. Green *et al.*, also 30 published by John Wiley & Son, for general guidance on protecting groups.

 It will be appreciated that certain of the various ring substituents in the compounds of the present invention may be introduced by standard aromatic substitution reactions or generated by conventional functional group modifications either prior to or immediately

following the processes mentioned above, and as such are included in the process aspect of the invention. Such reactions and modifications include, for example, introduction of a substituent by means of an aromatic substitution reaction, reduction of substituents, alkylation of substituents and oxidation of substituents. The reagents and reaction conditions for such procedures are well known in the chemical art. Particular examples of aromatic substitution reactions include the introduction of a nitro group using concentrated nitric acid, the introduction of an acyl group using, for example, an acyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; the introduction of an alkyl group using an alkyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; and the introduction of a halogeno group. Particular examples of modifications include the oxidation of alkylthio to alkylsulfinyl or alkylsulfonyl; the substitution of an NH or OH group in R¹ by the reaction with an optionally substituted alkyl halide or an optionally substituted alkyl-tosylate. An analogous procedure may be used to introduce optionally substituted (2-6C)alkenyloxy or (2-6C)alkynyloxy groups into R¹.

When a pharmaceutically-acceptable salt of a quinazoline derivative of the formula I is required, for example an acid-addition salt, it may be obtained by, for example, reaction of said quinazoline derivative with a suitable acid using a conventional procedure.

As mentioned hereinbefore some of the compounds according to the present invention may contain one or more chiral centers and may therefore exist as stereoisomers (for example when Q¹ is pyrrolidin-2-yl). Stereoisomers may be separated using conventional techniques, e.g. chromatography or fractional crystallisation. The enantiomers may be isolated by separation of a racemate for example by fractional crystallisation, resolution or HPLC. The diastereoisomers may be isolated by separation by virtue of the different physical properties of the diastereoisomers, for example, by fractional crystallisation, HPLC or flash chromatography. Alternatively particular stereoisomers may be made by chiral synthesis from chiral starting materials under conditions which will not cause racemisation or epimerisation, or by derivatisation, with a chiral reagent. When a specific stereoisomer is isolated it is suitably isolated substantially free for other stereoisomers, for example containing less than 20%, particularly less than 10% and more particularly less than 5% by weight of other stereoisomers.

In the section above relating to the preparation of the compound of formula I, the expression "inert solvent" refers to a solvent which does not react with the starting materials,

reagents, intermediates or products in a manner which adversely affects the yield of the desired product.

Persons skilled in the art will appreciate that, in order to obtain compounds of the invention in an alternative and in some occasions, more convenient manner, the individual process steps mentioned hereinbefore may be performed in different order, and/or the individual reactions may be performed at different stage in the overall route (i.e. chemical transformations may be performed upon different intermediates to those associated hereinbefore with a particular reaction).

Biological Assays

10 The following assays may be used to measure the effects of the compounds of the present invention as inhibitors of the erbB-tyrosine kinases, as inhibitors *in-vitro* of the proliferation of KB cells (human naso-pharyngeal carcinoma cells) and H16 N-2 cells, and as inhibitors *in vivo* on the growth in nude mice of xenografts of LoVo tumour cells (colorectal adenocarcinoma).

15 a) Protein Tyrosine Kinase phosphorylation Assays

This test measures the ability of a test compound to inhibit the phosphorylation of a tyrosine containing polypeptide substrate by EGFR tyrosine kinase enzyme.

Recombinant intracellular fragments of EGFR, erbB2 and erbB4 (accession numbers X00588, X03363 and L07868 respectively) were cloned and expressed in the
20 baculovirus/Sf21 system. Lysates were prepared from these cells by treatment with ice-cold lysis buffer (20mM N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid (HEPES) pH7.5, 150mM NaCl, 10% glycerol, 1% Triton X-100, 1.5mM MgCl₂, 1mM ethylene glycol-bis(β-aminoethyl ether) N',N',N',N'-tetraacetic acid (EGTA), plus protease inhibitors and then cleared by centrifugation.

25 Constitutive kinase activity of the recombinant protein was determined by its ability to phosphorylate a synthetic peptide (made up of a random co-polymer of Glutamic Acid, Alanine and Tyrosine in the ratio of 6:3:1). Specifically, MaxisorbTM 96-well immunoplates were coated with synthetic peptide (0.2μg of peptide in a 100μl phosphate buffered saline (PBS) solution and incubated at 4°C overnight). Plates were washed in PBS-T (phosphate
30 buffered saline with 0.5% Tween 20) then in 50mM HEPES pH 7.4 at room temperature to remove any excess unbound synthetic peptide. EGFR, ErbB2 or ErbB4 tyrosine kinase activity was assessed by incubation in peptide coated plates for 20 minutes at 22°C in 100mM

HEPES pH 7.4, adenosine triphosphate (ATP) at K_m concentration for the respective enzyme, 10mM $MnCl_2$, 0.1mM Na_3VO_4 , 0.2mM DL-dithiothreitol (DTT), 0.1% Triton X-100 with test compound in DMSO (final concentration of 2.5%). Reactions were terminated by the removal of the liquid components of the assay followed by washing of the plates with

5 PBS-T.

The immobilised phospho-peptide product of the reaction was detected by immunological methods. Firstly, plates were incubated for 90 minutes at room temperature with anti-phosphotyrosine primary antibodies that were raised in the mouse (4G10 from Upstate Biotechnology). Following extensive washing, plates were treated with Horseradish

10 Peroxidase (HRP) conjugated sheep anti-mouse secondary antibody (NXA931 from Amersham) for 60 minutes at room temperature. After further washing, HRP activity in each well of the plate was measured colorimetrically using 22'-Azino-di-[3-ethylbenzthiazoline sulfonate (6)] diammonium salt crystals (ABTSTM from Roche) as a substrate.

Quantification of colour development and thus enzyme activity was achieved by the

15 measurement of absorbance at 405nm on a Molecular Devices ThermoMax microplate reader. Kinase inhibition for a given compound was expressed as an IC_{50} value. This was determined by calculation of the concentration of compound that was required to give 50% inhibition of phosphorylation in this assay. The range of phosphorylation was calculated from the positive (vehicle plus ATP) and negative (vehicle minus ATP) control values.

20 **b) KB cell proliferation assay**

This assay measures the ability of a test compound to inhibit the EGF driven proliferation of KB cells (human naso-pharyngeal carcinoma obtained from the American Type Culture Collection (ATCC)).

KB cells were cultured in Dulbecco's modified Eagle's medium (DMEM) containing

25 10% foetal calf serum, 2 mM glutamine and non-essential amino acids at 37°C in a 7.5% CO_2 air incubator. Cells were harvested from the stock flasks using Trypsin/ethylenediaminetetraacetic acid (EDTA). Cell density was measured using a haemocytometer and viability was calculated using trypan blue solution before being seeded at a density of 1.25×10^3 cells per well of a 96 well plate in DMEM containing 2.5% charcoal

30 stripped serum, 2mM glutamine and non-essential amino acids at 37°C in 7.5% CO_2 and allowed to settle for 4 hours.

Following adhesion to the plate, the cells are treated with or without EGF (final concentration of 1ng/ml) and with or without compound at a range of concentrations in dimethylsulfoxide (DMSO) (1% final) in a final volume of 200µl before incubation for 4 days at 37°C in 7.5% CO₂. Following the incubation period, cell numbers were determined by
5 addition of 50µl of 3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) (stock 5mg/ml) and incubated at 37°C in a 7.5% CO₂ air incubator for 2 hours. MTT solution was then removed from the cells by aspiration, which were then allowed to air dry and were dissolved upon the addition of 100µl of DMSO.

Absorbance of this solubilised cells was read at 540nm to quantify cell biomass.

- 10 Inhibition of proliferation was expressed as an IC₅₀ value. This was determined by calculation of the concentration of compound that was required to give 50% inhibition of proliferation. The range of proliferation was calculated from the positive (vehicle plus EGF) and negative (vehicle minus EGF) control values.

c) H16N-2 cell proliferation assay

- 15 This assay measures the ability of a test compound to inhibit heregulin β or EGF driven proliferation of H16N-2 cells. These non-neoplastic epithelial cells respond in a proliferative manner to stimulation with either EGF or heregulin β (Ram, G.R. and Ethier, S.P. (1996) *Cell Growth and Differentiation*, 7, 551-561) were isolated human mammary tissue (Band, V. and Sager, R. Tumour progression in breast cancer. In: J. S. Rhim and A.
20 Dritschilo (eds.), *Neoplastic Transformation in human Cell Culture*, pp 169-178. Clifton, NJ: Humana Press, 1991) and were obtained from the Dana-Farber Cancer Institute, 44 Binney Street, Boston, Massachusetts 02115.

- H16N-2 cells were routinely cultured in culture medium (a 1:1 mix of Gibco F12 and Ham's αMEM media containing 1 % foetal calf serum, 10mM HEPES, 1µg/ml Insulin,
25 12.5ng/ml EGF, 2.8µM Hydrocortisone, 2nM Estradiol 5µM Ascorbic Acid, 10µg/ml Transferrin, 0.1mM Phosphoethanolamine, 15nM Sodium Selenite, 2mM Glutamine, 10nM Tri-iodo-thyrone, 35µg/ml Bovine pituitary Extract and 0.1mM Ethanolamine) at 37°C in a 7.5% CO₂ air incubator. Cells were harvested from the stock flasks using Trypsin/ethylaminetetraacetic acid (EDTA). Cell density was measured using a
30 haemocytometer and viability was calculated using trypan blue solution before being seeded at a density of 1.0x10³ cells per well of a 96 well plate in the above media at 37°C in 7.5% CO₂ and allowed to settle for 72 hours.

Following this, the cells were starved of serum for 24 hours upon the addition of starvation medium (a 1:1 mix of Gibco F12 and Ham's α MEM media containing, 10mM HEPES, 2nM Estradiol, 5 μ M Ascorbic Acid, 10 μ g/ml Transferrin, 0.1mM Phosphoethanolamine, 15nM Sodium Selenite, 2mM Glutamine, and 0.1mM Ethanolamine) and incubated at 37°C in 7.5% CO₂. The cells were then treated with or without compound at a range of concentrations in dimethylsulfoxide (DMSO) (1% final) for two hours before the addition of exogenous ligand (at a final concentration of 100ng/ml of heregulin β or 5ng/ml of EGF) and incubation with both ligand and compound in a total volume of 200 μ l for 4 days at 37°C in 7.5% CO₂. Following the incubation period, cell numbers were determined by addition of 50 μ l of 3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) (stock 5mg/ml) and incubated at 37°C in a 7.5% CO₂ air incubator for 2 hours. MTT solution was then removed from the cells by aspiration, which were then allowed to air dry and were dissolved upon the addition of 100 μ l of DMSO.

Absorbance of this solubilised cells was read at 540nm to quantify cell biomass. Inhibition of proliferation was expressed as an IC₅₀ value. This was determined by calculation of the concentration of compound that was required to give 50% inhibition of proliferation. The range of proliferation was calculated from the positive (vehicle plus ligand) and negative (vehicle minus ligand) control values.

(d) Clone 24 phospho-erbB2 cell assay

This immunofluorescence end point assay measures the ability of a test compound to inhibit the phosphorylation of erbB2 in a MCF7 (breast carcinoma) derived cell line which was generated by transfecting MCF7 cells with the full length erbB2 gene using standard methods to give a cell line that overexpresses full length wild type erbB2 protein (hereinafter 'Clone 24' cells).

Clone 24 cells were cultured in Growth Medium (phenol red free Dulbecco's modified Eagle's medium (DMEM) containing 10% foetal bovine serum, 2 mM glutamine and 1.2mg/ml G418) in a 7.5% CO₂ air incubator at 37°C. Cells were harvested from T75 stock flasks by washing once in PBS (phosphate buffered saline, pH7.4, Gibco No. 10010-015) and harvested using 2mls of Trypsin (1.25mg/ml) / ethylenediaminetetraacetic acid (EDTA) (0.8mg/ml) solution. The cells were resuspended in Growth Medium. Cell density was measured using a haemocytometer and viability was calculated using Trypan Blue

solution before being further diluted in Growth Medium and seeded at a density of 1×10^4 cells per well (in 100ul) into clear bottomed 96 well plates (Packard, No. 6005182).

3 days later, Growth Medium was removed from the wells and replaced with 100ul Assay Medium (phenol red free DMEM, 2mM glutamine, 1.2mg/ml G418) either with or
5 without erbB inhibitor compound. Plates were returned to the incubator for 4hrs and then 20ul of 20% formaldehyde solution in PBS was added to each well and the plate was left at room temperature for 30 minutes. This fixative solution was removed with a multichannel pipette, 100ul of PBS was added to each well and then removed with a multichannel pipette and then 50ul PBS was added to each well. Plates were then sealed and stored for up to 2 weeks at 4°C.

10 Immunostaining was performed at room temperature. Wells were washed once with 200ul PBS / Tween 20 (made by adding 1 sachet of PBS / Tween dry powder (Sigma, No. P3563) to 1L of double distilled H₂O) using a plate washer then 200ul Blocking Solution (5% Marvel dried skimmed milk (Nestle) in PBS / Tween 20) was added and incubated for 10 minutes. Blocking Solution was removed using a plate washer and 200ul of 0.5% Triton X-
15 100 / PBS was added to permeabilise the cells. After 10 minutes, the plate was washed with 200ul PBS / Tween 20 and then 200ul Blocking Solution was added once again and incubated for 15 minutes. Following removal of the Blocking Solution with a plate washer, 30ul of rabbit polyclonal anti-phospho ErbB2 IgG antibody (epitope phospho-Tyr 1248, SantaCruz, No. SC-12352-R), diluted 1:250 in Blocking Solution, was added to each well and incubated
20 for 2 hours. Then this primary antibody solution was removed from the wells using a plate washer followed by two 200ul PBS / Tween 20 washes using a plate washer. Then 30ul of Alexa-Fluor 488 goat anti-rabbit IgG secondary antibody (Molecular Probes, No. A-11008), diluted 1:750 in Blocking Solution, was added to each well. From now onwards, wherever possible, plates were protected from light exposure, at this stage by sealing with black backing
25 tape. The plates were incubated for 45 minutes and then the secondary antibody solution was removed from the wells followed by two 200ul PBS / Tween 20 washes using a plate washer. Then 100ul PBS was added to each plate, incubated for 10 minutes and then removed using a plate washer. Then a further 100ul PBS was added to each plate and then, without prolonged incubation, removed using a plate washer. Then 50ul of PBS was added to each well and
30 plates were resealed with black backing tape and stored for up to 2 days at 4°C before analysis.

The Fluorescence signal in each well was measured using an Acumen Explorer Instrument (Acumen Bioscience Ltd.), a plate reader that can be used to rapidly quantitate features of images generated by laser-scanning. The instrument was set to measure the number of fluorescent objects above a pre-set threshold value and this provided a measure of the phosphorylation status of erbB2 protein. Fluorescence dose response data obtained with each compound was exported into a suitable software package (such as Origin) to perform curve fitting analysis. Inhibition of erbB2 phosphorylation was expressed as an IC₅₀ value. This was determined by calculation of the concentration of compound that was required to give 50% inhibition of erbB2 phosphorylation signal.

10 e) *In vivo* Xenograft assay

This assay measures the ability of a test compound to inhibit the growth of a LoVo tumour (colorectal adenocarcinoma obtained from the ATCC) in Female Swiss athymic mice (Alderley Park, *nu/nu* genotype).

Female Swiss athymic (*nu/nu* genotype) mice were bred and maintained in Alderley Park in negative pressure Isolators (PFI Systems Ltd.). Mice were housed in a barrier facility with 12hr light/dark cycles and provided with sterilised food and water *ad libitum*. All procedures were performed on mice of at least 8 weeks of age. LoVo tumour cell (colorectal adenocarcinoma obtained from the ATCC) xenografts were established in the hind flank of donor mice by sub cutaneous injections of 1×10^7 freshly cultured cells in 100µl of serum free media per animal. On day 5 post-implant, mice were randomised into groups of 7 prior to the treatment with compound or vehicle control that was administered once daily at 0.1ml/10g body weight. Tumour volume was assessed twice weekly by bilateral Vernier calliper measurement, using the formula $(\text{length} \times \text{width}) \times \sqrt{(\text{length} \times \text{width}) \times (\pi/6)}$, where length was the longest diameter across the tumour, and width was the corresponding perpendicular. Growth inhibition from start of study was calculated by comparison of the mean changes in tumour volume for the control and treated groups, and statistical significance between the two groups was evaluated using a Students *t* test.

Although the pharmacological properties of the compounds of the formula I vary with structural change as expected, in general activity possessed by compounds of the formula I, may be demonstrated at the following concentrations or doses in one or more of the above tests (a), (b), (c) and (d):-

Test (a):- IC₅₀ in the range, for example, 0.001 - 10 µM;

Test (b):- IC_{50} in the range, for example, 0.001 - 10 μM ;

Test (c):- IC_{50} in the range, for example, 0.001 - 10 μM ;

Test (d):- IC_{50} in the range, for example, 0.001 - 10 μM ;

Test (e):- activity in the range, for example, 1-200 mg/kg/day;

- 5 No physiologically unacceptable toxicity was observed in Test (e) at the effective dose for compounds tested of the present invention. Accordingly no untoward toxicological effects are expected when a compound of formula I, or a pharmaceutically-acceptable salt thereof, as defined hereinbefore is administered at the dosage ranges defined hereinafter.

- By way of example, Table A illustrates the activity of representative compounds
 10 according to the invention. Column 2 of Table A shows IC_{50} data from Test (a) for the inhibition of EGFR tyrosine kinase protein phosphorylation; column 3 shows IC_{50} data from Test (a) for the inhibition of erbB2 tyrosine kinase protein phosphorylation; and column 4 shows IC_{50} data for inhibition of proliferation of KB cells in test (b) described above:

Table A –

Example Number	IC_{50} (nM) Test (a): Inhibition of EGFR tyrosine kinase protein phosphorylation	IC_{50} (nM) Test (a): Inhibition of erbB2 tyrosine kinase protein phosphorylation	IC_{50} (nM) Test (b): EGFR driven KB cell proliferation assay
2	70	1852	141
3	9	275	91
4	39	252	138
6	57	1631	130
13	1	358	28
16	1	52	128
18	8	85	99
22	1	61	75
23	56	1727	288
25	32	1910	156
31	42	532	112
32	18	215	118

According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a quinazoline derivative of the formula I, or a pharmaceutically-acceptable thereof, as defined hereinbefore in association with a
5 pharmaceutically-acceptable diluent or carrier.

The compositions of the invention may be in a form suitable for oral use (for example as tablets, lozenges, hard or soft capsules, aqueous or oily suspensions, emulsions, dispersible powders or granules, syrups or elixirs), for topical use (for example as creams, ointments, gels, or aqueous or oily solutions or suspensions), for administration by inhalation (for
10 example as a finely divided powder or a liquid aerosol), for administration by insufflation (for example as a finely divided powder) or for parenteral administration (for example as a sterile aqueous or oily solution for intravenous, subcutaneous, intramuscular or intramuscular dosing or as a suppository for rectal dosing).

The compositions of the invention may be obtained by conventional procedures using
15 conventional pharmaceutical excipients, well known in the art. Thus, compositions intended for oral use may contain, for example, one or more colouring, sweetening, flavouring and/or preservative agents.

The amount of active ingredient that is combined with one or more excipients to produce a single dosage form will necessarily vary depending upon the host treated and the
20 particular route of administration. For example, a formulation intended for oral administration to humans will generally contain, for example, from 0.5 mg to 0.5 g of active agent (more suitably from 0.5 to 100 mg, for example from 1 to 30 mg) compounded with an appropriate and convenient amount of excipients which may vary from about 5 to about 98 percent by weight of the total composition.

25 The size of the dose for therapeutic or prophylactic purposes of a quinazoline derivative of the formula I will naturally vary according to the nature and severity of the conditions, the age and sex of the animal or patient and the route of administration, according to well known principles of medicine.

In using a quinazoline derivative of the formula I for therapeutic or prophylactic
30 purposes it will generally be administered so that a daily dose in the range, for example, 0.1 mg/kg to 75 mg/kg body weight is received, given if required in divided doses. In general lower doses will be administered when a parenteral route is employed. Thus, for example, for intravenous administration, a dose in the range, for example, 0.1 mg/kg to 30 mg/kg body

weight will generally be used. Similarly, for administration by inhalation, a dose in the range, for example, 0.05 mg/kg to 25 mg/kg body weight will be used. Oral administration is however preferred, particularly in tablet form. Typically, unit dosage forms will contain about 0.5 mg to 0.5 g of a compound of this invention.

5 We have found that the compounds of the present invention possess anti-proliferative properties such as anti-cancer properties that are believed to arise from their erbB family receptor tyrosine kinase inhibitory activity, particularly inhibition of the EGF receptor (erbB1) tyrosine kinase. Furthermore, certain of the compounds according to the present invention possess substantially better potency against the EGF receptor tyrosine kinase, than against
10 other tyrosine kinase enzymes, for example erbB2. Such compounds possess sufficient potency against the EGF receptor tyrosine kinase that they may be used in an amount sufficient to inhibit EGF receptor tyrosine kinase whilst demonstrating little, or significantly lower, activity against other tyrosine kinase enzymes such as erbB2. Such compounds are likely to be useful for the selective inhibition of EGF receptor tyrosine kinase and are likely to
15 be useful for the effective treatment of, for example EGF driven tumours.

Accordingly, the compounds of the present invention are expected to be useful in the treatment of diseases or medical conditions mediated alone or in part by erbB receptor tyrosine kinases (especially EGF receptor tyrosine kinase), i.e. the compounds may be used to produce an erbB receptor tyrosine kinase inhibitory effect in a warm-blooded animal in need
20 of such treatment. Thus the compounds of the present invention provide a method for the treatment of malignant cells characterised by inhibition of one or more of the erbB family of receptor tyrosine kinases. Particularly the compounds of the invention may be used to produce an anti-proliferative and/or pro-apoptotic and/or anti-invasive effect mediated alone or in part by the inhibition of erbB receptor tyrosine kinases. Particularly, the compounds of
25 the present invention are expected to be useful in the prevention or treatment of those tumours that are sensitive to inhibition of one or more of the erbB receptor tyrosine kinases, such as EGF and/or erbB2 and/or erbB4 receptor tyrosine kinases (especially EGF receptor tyrosine kinase) that are involved in the signal transduction steps which drive proliferation and survival of these tumour cells. Accordingly the compounds of the present invention are expected to be
30 useful in the treatment of psoriasis, benign prostatic hyperplasia (BPH), atherosclerosis and restenosis and/or cancer by providing an anti-proliferative effect, particularly in the treatment of erbB receptor tyrosine kinase sensitive cancers. Such benign or malignant tumours may affect any tissue and include non-solid tumours such as leukaemia, multiple myeloma or

lymphoma, and also solid tumours, for example bile duct, bone, bladder, brain/CNS, breast, colorectal, cervical, endometrial, gastric, head and neck, hepatic, lung, muscle, neuronal, oesophageal, ovarian, pancreatic, pleural/peritoneal membranes, prostate, renal, skin, testicular, thyroid, uterine and vulval tumours. According to this aspect of the invention there is provided a quinazoline derivative of the formula I, or a pharmaceutically acceptable salt thereof, for use as a medicament.

According to a further aspect of the invention there is provided a compound of the formula I, or a pharmaceutically acceptable salt thereof, for use in the production of an anti-proliferative effect in a warm-blooded animal such as man.

10 Thus according to this aspect of the invention there is provided the use of a quinazoline derivative of the formula I, or a pharmaceutically-acceptable salt thereof, as defined hereinbefore in the manufacture of a medicament for use in the production of an anti-proliferative effect in a warm-blooded animal such as man.

According to a further feature of this aspect of the invention there is provided a
15 method for producing an anti-proliferative effect in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a quinazoline derivative of the formula I, or a pharmaceutically acceptable salt thereof, as hereinbefore defined.

According to a further aspect of the present invention there is provided the use of a
20 quinazoline derivative of the formula I, or a pharmaceutically-acceptable salt thereof, as defined hereinbefore in the manufacture of a medicament for use in the treatment of a disease state or medical condition (for example a cancer as mentioned herein) mediated alone or in part by one or more erbB receptor tyrosine kinases (such as EGFR and/or erbB2 and/or erbB4, especially EGFR).

25 According to a further feature of this aspect of the invention there is provided a method for treating a disease or medical condition (for example a cancer as mentioned herein) mediated alone or in part by one or more erbB receptor tyrosine kinases (such as EGFR and/or erbB2 and/or erbB4, especially EGFR) in a warm-blooded animal, such as man, in need of such treatment, which comprises administering to said animal an effective amount of a
30 quinazoline derivative of the formula I, or a pharmaceutically-acceptable salt thereof, as defined hereinbefore.

According to a further aspect of the invention there is provided a quinazoline derivative of the formula I, or a pharmaceutically acceptable salt thereof, for use in the

treatment of a disease or medical condition (for example a cancer as mentioned herein) mediated alone or in part by one or more erbB receptor tyrosine kinases (such as EGFR and/or erbB2 and/or erbB4, especially EGFR).

According to a further aspect of the invention there is provided the use of a
5 quinazoline derivative of the formula I, or a pharmaceutically-acceptable salt thereof, as defined hereinbefore in the manufacture of a medicament for use in the prevention or treatment of those tumours which are sensitive to inhibition of one or more of the erbB receptor tyrosine kinases, such as EGFR and/or erbB2 and/or erbB4 (especially EGFR), that are involved in the signal transduction steps which lead to the proliferation of tumour cells.

10 According to a further feature of this aspect of the invention there is provided a method for the prevention or treatment of those tumours which are sensitive to inhibition of one or more of the erbB family of receptor tyrosine kinases, such as EGFR and/or erbB2 and/or erbB4 (especially EGFR), that are involved in the signal transduction steps which lead to the proliferation and/or survival of tumour cells in a warm-blooded animal, such as man, in
15 need of such treatment, which comprises administering to said animal an effective amount of a quinazoline derivative of the formula I, or a pharmaceutically-acceptable salt thereof, as defined hereinbefore.

According to a further feature of this aspect of the invention there is provided a compound of the formula I, or a pharmaceutically acceptable salt thereof, for use in the
20 prevention or treatment of those tumours which are sensitive to inhibition of erbB receptor tyrosine kinases, such as EGFR and/or erbB2 and/or erbB4 (especially EGFR), that are involved in the signal transduction steps which lead to the proliferation of tumour cells.

According to a further aspect of the invention there is provided the use of a quinazoline derivative of the formula I, or a pharmaceutically-acceptable salt thereof, as
25 defined hereinbefore in the manufacture of a medicament for use in providing a EGFR and/or erbB2 and/or erbB4 (especially a EGFR) tyrosine kinase inhibitory effect.

According to a further feature of this aspect of the invention there is provided a method of treatment for providing a EGFR and/or an erbB2 and or an erbB4 (especially a EGFR) tyrosine kinase inhibitory effect in a warm-blooded animal, such as man, in need of
30 such treatment, which comprises administering to said animal an effective amount of a quinazoline derivative of the formula I, or a pharmaceutically-acceptable salt thereof, as defined hereinbefore.

According to a further feature of this aspect of the invention there is provided a compound of the formula I, or a pharmaceutically acceptable salt thereof, for use in providing a EGFR and/or erbB2 and/or erbB4 (especially a EGFR) tyrosine kinase inhibitory effect.

According to a further feature of the present invention there is provided the use of a
5 quinazoline derivative of the formula I, or a pharmaceutically-acceptable salt thereof, as defined hereinbefore in the manufacture of a medicament for use in providing a selective EGFR tyrosine kinase inhibitory effect.

According to a further feature of this aspect of the invention there is provided a method of treatment for providing a selective EGFR tyrosine kinase inhibitory effect in a
10 warm-blooded animal, such as man, in need of such treatment, which comprises administering to said animal an effective amount of a quinazoline derivative of the formula I, or a pharmaceutically-acceptable salt thereof, as defined hereinbefore.

According to a further feature of this aspect of the invention there is provided a compound of the formula I, or a pharmaceutically acceptable salt thereof, for use in providing
15 a selective EGFR tyrosine kinase inhibitory effect.

By "a selective EGFR kinase inhibitory effect" is meant that the quinazoline derivative of formula I is more potent against EGF receptor tyrosine kinase than it is against other kinases. In particular some of the compounds according to the invention are more potent against EGF receptor kinase than it is against other tyrosine kinases such as other erbB
20 receptor tyrosine kinases, particularly erbB2. Particularly a selective EGFR kinase inhibitor according to the invention is at least 5 times, preferably at least 10 times more potent against EGFR tyrosine kinase driven proliferation than it is against erbB2 receptor tyrosine kinase driven proliferation. The relative potency against EGFR and erbB2 may be determined by comparison of the relative IC₅₀ values obtained in suitable assays, such as an in-vitro cellular
25 assay, for example by comparison of the IC₅₀ values obtained from the H16N-2 assay described above under heregulin β and EGF driven proliferative conditions. Alternatively by comparison of the IC₅₀ values from the KB cell proliferation assay (test (b) above) and the Clone 24 phospho-erbB2 cell assay (test (c) above).

According to a further aspect of the present invention there is provided the use of a
30 quinazoline derivative of the formula I, or a pharmaceutically-acceptable salt thereof, as defined hereinbefore in the manufacture of a medicament for use in the treatment of a cancer, for example a cancer selected from leukaemia, multiple myeloma, lymphoma, bile duct, bone, bladder, brain/CNS, breast, colorectal, cervical, endometrial, gastric, head and neck, hepatic,

lung, muscle, neuronal, oesophageal, ovarian, pancreatic, pleural/peritoneal membranes, prostate, renal, skin, testicular, thyroid, uterine and vulval cancer.

According to a further feature of this aspect of the invention there is provided a method for treating a cancer, for example a cancer selected from leukaemia, multiple
5 myeloma, lymphoma, bile duct, bone, bladder, brain/CNS, breast, colorectal, cervical, endometrial, gastric, head and neck, hepatic, lung, muscle, neuronal, oesophageal, ovarian, pancreatic, pleural/peritoneal membranes, prostate, renal, skin, testicular, thyroid, uterine and vulval cancer in a warm-blooded animal, such as man, in need of such treatment, which comprises administering to said animal an effective amount of a quinazoline derivative of the
10 formula I, or a pharmaceutically-acceptable salt thereof, as defined hereinbefore.

According to a further aspect of the invention there is provided a quinazoline derivative of the formula I, or a pharmaceutically acceptable salt thereof, for use in the treatment of a cancer, for example a cancer selected from leukaemia, multiple myeloma, lymphoma, bile duct, bone, bladder, brain/CNS, breast, colorectal, cervical, endometrial,
15 gastric, head and neck, hepatic, lung, muscle, neuronal, oesophageal, ovarian, pancreatic, pleural/peritoneal membranes, prostate, renal, skin, testicular, thyroid, uterine and vulval cancer.

As mentioned above the size of the dose required for the therapeutic or prophylactic treatment of a particular disease will necessarily be varied depending upon, amongst other
20 things, the host treated, the route of administration and the severity of the illness being treated.

The anti-proliferative treatment defined hereinbefore may be applied as a sole therapy or may involve, in addition to the quinazoline derivative of the invention, conventional surgery or radiotherapy or chemotherapy. Such chemotherapy may include one or more of the following categories of anti-tumour agents :-

- 25 (i) antiproliferative/antineoplastic drugs and combinations thereof, as used in medical oncology, such as alkylating agents (for example cis-platin, carboplatin, cyclophosphamide, nitrogen mustard, melphalan, chlorambucil, busulfan and nitrosoureas); antimetabolites (for example antifolates such as fluoropyrimidines like 5-fluorouracil and tegafur, raltitrexed, methotrexate, cytosine arabinoside and hydroxyurea; antitumour antibiotics (for example
30 anthracyclines like adriamycin, bleomycin, doxorubicin, daunomycin, epirubicin, idarubicin, mitomycin-C, dactinomycin and mithramycin); antimitotic agents (for example vinca alkaloids like vincristine, vinblastine, vindesine and vinorelbine and taxoids like taxol and

- taxotere); and topoisomerase inhibitors (for example epipodophyllotoxins like etoposide and teniposide, amsacrine, topotecan and camptothecin);
- (ii) cytostatic agents such as antioestrogens (for example tamoxifen, toremifene, raloxifene, droloxifene and idoxifene), oestrogen receptor down regulators (for example
5 fulvestrant), antiandrogens (for example bicalutamide, flutamide, nilutamide and cyproterone acetate), LHRH antagonists or LHRH agonists (for example goserelin, leuprorelin and buserelin), progestogens (for example megestrol acetate), aromatase inhibitors (for example as anastrozole, letrozole, vorazole and exemestane) and inhibitors of 5 α -reductase such as finasteride;
- 10 (iii) agents which inhibit cancer cell invasion (for example metalloproteinase inhibitors like marimastat and inhibitors of urokinase plasminogen activator receptor function);
- (iv) inhibitors of growth factor function, for example such inhibitors include growth factor antibodies, growth factor receptor antibodies (for example the anti-erbB2 antibody trastuzumab [HerceptinTM] and the anti-erbB1 antibody cetuximab [C225]), farnesyl
15 transferase inhibitors, tyrosine kinase inhibitors and serine/threonine kinase inhibitors, for example other inhibitors of the epidermal growth factor family (for example EGFR family tyrosine kinase inhibitors such as N-(3-chloro-4-fluorophenyl)-7-methoxy-6-(3-morpholinopropoxy)quinazolin-4-amine (gefitinib, AZD1839), N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)quinazolin-4-amine (erlotinib, OSI-774) and 6-acrylamido-N-(3-chloro-
20 4-fluorophenyl)-7-(3-morpholinopropoxy)quinazolin-4-amine (CI 1033)), for example inhibitors of the platelet-derived growth factor family and for example inhibitors of the hepatocyte growth factor family;
- (v) antiangiogenic agents such as those which inhibit the effects of vascular endothelial growth factor, (for example the anti-vascular endothelial cell growth factor antibody
25 bevacizumab [AvastinTM], compounds such as those disclosed in International Patent Applications WO 97/22596, WO 97/30035, WO 97/32856 and WO 98/13354) and compounds that work by other mechanisms (for example linomide, inhibitors of integrin $\alpha v \beta 3$ function and angiostatin);
- (vi) vascular damaging agents such as Combretastatin A4 and compounds disclosed in
30 International Patent Applications WO 99/02166, WO00/40529, WO 00/41669, WO01/92224, WO02/04434 and WO02/08213;

- (vii) antisense therapies, for example those which are directed to the targets listed above, such as ISIS 2503, an anti-ras antisense;
- (viii) gene therapy approaches, including for example approaches to replace aberrant genes such as aberrant p53 or aberrant BRCA1 or BRCA2, GDEPT (gene-directed enzyme pro-drug
5 therapy) approaches such as those using cytosine deaminase, thymidine kinase or a bacterial nitroreductase enzyme and approaches to increase patient tolerance to chemotherapy or radiotherapy such as multi-drug resistance gene therapy; and
- (ix) immunotherapy approaches, including for example ex-vivo and in-vivo approaches to increase the immunogenicity of patient tumour cells, such as transfection with cytokines such
10 as interleukin 2, interleukin 4 or granulocyte-macrophage colony stimulating factor, approaches to decrease T-cell anergy, approaches using transfected immune cells such as cytokine-transfected dendritic cells, approaches using cytokine-transfected tumour cell lines and approaches using anti-idiotypic antibodies.

Such conjoint treatment may be achieved by way of the simultaneous, sequential or
15 separate dosing of the individual components of the treatment. Such combination products employ the compounds of this invention within the dosage range described hereinbefore and the other pharmaceutically-active agent within its approved dosage range.

According to this aspect of the invention there is provided a pharmaceutical product comprising a quinazoline derivative of the formula I as defined hereinbefore and an additional
20 anti-tumour agent as defined hereinbefore for the conjoint treatment of cancer.

Although the compounds of the formula I are primarily of value as therapeutic agents for use in warm-blooded animals (including man), they are also useful whenever it is required to inhibit the effects of the erbB receptor tyrosine protein kinases. Thus, they are useful as pharmacological standards for use in the development of new biological tests and in the search
25 for new pharmacological agents.

In the above pharmaceutical composition, process, method, use and medicament manufacture features of the present invention, the alternative and preferred embodiments of the compounds of the invention described herein also apply.

The invention will now be illustrated by the following non limiting examples in which,
30 unless stated otherwise:


- (i) temperatures are given in degrees Celsius (°C); operations were carried out at room or ambient temperature, that is, at a temperature in the range of 18-25°C;

- (ii) organic solutions were dried over anhydrous magnesium sulfate; evaporation of solvent was carried out using a rotary evaporator under reduced pressure (600-4000 Pascals; 4.5-30mmHg) with a bath temperature of up to 60°C;
- (iii) chromatography means flash chromatography on silica gel; thin layer chromatography
- 5 (TLC) was carried out on silica gel plates;
- (iv) in general, the course of reactions was followed by TLC and / or analytical LC-MS, and reaction times are given for illustration only;
- (v) final products had satisfactory proton nuclear magnetic resonance (NMR) spectra and/or mass spectral data;
- 10 (vi) yields are given for illustration only and are not necessarily those which can be obtained by diligent process development; preparations were repeated if more material was required;
- (vii) when given, NMR data is in the form of delta values for major diagnostic protons, given in parts per million (ppm) relative to tetramethylsilane (TMS) as an internal standard, determined at 300 MHz using perdeuterio dimethyl sulfoxide (DMSO-d₆) as solvent unless
- 15 otherwise indicated; the following abbreviations have been used: s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet; b, broad;
- (viii) chemical symbols have their usual meanings; SI units and symbols are used;
- (ix) solvent ratios are given in volume:volume (v/v) terms; and
- (x) mass spectra were run with an electron energy of 70 electron volts in the chemical
- 20 ionization (CI) mode using a direct exposure probe; where indicated ionization was effected by electron impact (EI), fast atom bombardment (FAB) or electrospray (ESP); values for m/z are given; generally, only ions which indicate the parent mass are reported; and unless otherwise stated, the mass ion quoted is (MH)⁺ which refers to the protonated mass ion; reference to M⁺ is to the mass ion generated by loss of an electron; and reference to M-H⁺ is to
- 25 the mass ion generated by loss of a proton;
- (xi) unless stated otherwise compounds containing an asymmetrically substituted carbon and/or sulfur atom have not been resolved;
- (xii) where a synthesis is described as being analogous to that described in a previous example the amounts used are the millimolar ratio equivalents to those used in the previous example;
- 30 (xiii) the following abbreviations have been used:
- | | |
|-----|--------------------------------|
| THF | tetrahydrofuran; |
| DMF | <i>N,N</i> -dimethylformamide; |
| DMA | <i>N,N</i> -dimethylacetamide; |

	DCM	dichloromethane;
	DMSO	dimethylsulfoxide;
	IPA	Isopropyl alcohol;
	ether	diethyl ether; and
5	HPLC	high pressure liquid chromatography.

Example 1**2-[(2S)-2-[(4-[3-Chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]pyrrolidin-1-yl]-2-oxoethanol**

(Process (a))



Glycolic acid (20 mg) and *O*-(7-azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (HATU, 109 mg) were added sequentially, each in one portion, to a stirred solution of *N*-(3-chloro-4-fluorophenyl)-7-methoxy-5-[(2*S*)-pyrrolidin-2-ylmethoxy]-quinazolin-4-amine (96 mg) and *N,N*-diisopropylethylamine (83 μ l) in DCM (3 ml) at room temperature under a nitrogen atmosphere. The reaction mixture was stirred for 16 hours 30 minutes and then diluted with DCM (10 ml) and washed with saturated aqueous sodium hydrogen carbonate and water. The organic layer was dried (MgSO_4) and concentrated *in vacuo* to leave a yellow oil, which was purified by column chromatography using 0-10% methanol in dichloromethane as eluent to give the title compound as a white solid (55 mg, 15 50%); NMR Spectrum (DMSO- d_6) 9.90 (s, 1H), 8.42 (s, 1H), 8.15 (dd, 1H), 7.70-7.60 (m, 1H), 7.40 (t, 1H), 6.88 (d, 1H), 6.78 (d, 1H), 4.65-4.53 (m, 2H), 4.40 (dd, 1H), 4.17 (dd, 1H), 4.03 (d, 1H), 3.90 (s, 3H), 3.45-3.35 (m, 3H), 2.10-1.82 (m, 4H); Mass spectrum MH^+ 461.

The *N*-(3-chloro-4-fluorophenyl)-7-methoxy-5-[(2*S*)-pyrrolidin-2-ylmethoxy]-quinazolin-4-amine used as starting material was prepared as follows:

20 DMF (12 ml) was added dropwise over 5 minutes to a slurry of 5,7-difluoroquinazolin-4(3*H*)-one (15 g) (obtained as described in WO01/94341 Example 4, foot note 5) in thionyl chloride (120 ml) at room temperature under a nitrogen atmosphere and the grey mixture was heated at 110°C for 16 hours. The reaction mixture was cooled to room temperature and concentrated *in vacuo*. There was thus obtained 4-chloro-5,7-difluoro-25 quinazoline as a brown solid (16.5 g) which was used without further purification.

3-Chloro-4-fluoroaniline (12 g) and 1N HCl in dioxane (40 ml) were added sequentially, each in one portion, to a mixture of 4-chloro-5,7-difluoroquinazoline (16.5 g) in IPA (250 ml) and the mixture was stirred at 80°C under a nitrogen atmosphere for 17 hours. The reaction mixture was cooled to 0°C and the precipitated solid was filtered and dried to

leave *N*-(3-chloro-4-fluorophenyl)-5,7-difluoroquinazolin-4-amine as a green solid (18.2 g, 71%); Mass spectrum MH^+ 310.

Methanol (6.5 ml) and potassium *tert*-butoxide (18 g) were added sequentially, each in one portion, to a mixture of *N*-(3-chloro-4-fluorophenyl)-5,7-difluoroquinazolin-4-amine (13.8 g) in THF (270 ml) at room temperature under a nitrogen atmosphere. The mixture was heated at 75 °C for 18 hours 30 minutes and then cooled to room temperature and concentrated *in vacuo* to leave a brown solid. The solid was cooled to 0°C and cold water was added and then the precipitate was filtered and dried to leave *N*-(3-chloro-4-fluorophenyl)-5,7-dimethoxyquinazolin-4-amine as a light brown solid (10.4g, 69%); Mass spectrum MH^+ 334.

Pyridine hydrochloride (2.33g) was added in one portion to a stirred mixture of *N*-(3-chloro-4-fluorophenyl)-5,7-dimethoxyquinazolin-4-amine (6.4 g) in pyridine (75 ml) at room temperature. The reaction mixture was stirred at 115 °C under a nitrogen atmosphere for 23 hours and then the brown solution was cooled to room temperature and the yellow precipitate was filtered and washed with ice cold water to leave a light yellow solid. The solid was taken up in 7N ammonia in methanol and stirred at room temperature for 10 minutes. The solution was filtered and the filtrate was concentrated *in vacuo* to leave 4-[(3-chloro-4-fluorophenyl)-amino]-7-methoxyquinazolin-5-ol as a light brown solid (4.2g, 68%); Mass spectrum MH^+ 320.

tert-Butyl (2*S*)-(-)-2-(hydroxymethyl)pyrrolidine-1-carboxylate (840 mg) and a solution of di-*tert*-butyl azodicarboxylate (1.32 g) in DCM (10 ml) were added sequentially, each in one portion, to a stirred mixture of 4-[(3-chloro-4-fluorophenyl)-amino]-7-methoxyquinazolin-5-ol (1.22 g) and triphenylphosphine (1.50 g) in DCM (50 ml) at 0 °C under a nitrogen atmosphere. The mixture was warmed to room temperature and stirred for 17 hours. The mixture was filtered, concentrated *in vacuo*, and purified by column chromatography using 0-10% methanol in DCM as eluent to give *tert*-butyl (2*S*)-2-[(4-[3-chloro-4-fluoro-anilino]-7-methoxyquinazolin-5-yl)oxy)methyl]pyrrolidine-1-carboxylate as a thick yellow oil (1.6 g, containing 6% triphenylphosphine oxide); Mass spectrum MH^+ 503.

Trifluoroacetic acid (25 ml) was added in one portion to *tert*-butyl (2*S*)-2-[(4-[3-chloro-4-fluoro-anilino]-7-methoxyquinazolin-5-yl)oxy)methyl]pyrrolidine-1-carboxylate (1.6 g) and the solution was stirred at room temperature for 1 hour 30 minutes. The solution was then concentrated *in vacuo* and azeotroped three times with ether to leave an orange oil. 7N Ammonia in methanol (5 ml) was added and the mixture was stirred for 10 minutes. The


mixture was then concentrated *in vacuo* to leave an orange oil and purified by column chromatography using 0-10% methanol in DCM as eluent to give *N*-(3-chloro-4-fluorophenyl)-7-methoxy-5-[(2*S*)-pyrrolidin-2-ylmethoxy]-quinazolin-4-amine as a light yellow solid (1.15g, 75% over two steps); Mass spectrum MH^+ 403.

5

Example 2

N-(3-Chloro-4-fluorophenyl)-5-[(2*S*)-1-[(dimethylamino)acetyl]pyrrolidin-2-yl]methoxy)-7-methoxyquinazolin-4-amine

(Process (a))




The procedure described in Example 1 was repeated using *N*-(3-chloro-4-fluorophenyl)-7-methoxy-5-[(2*S*)-pyrrolidin-2-ylmethoxy]-quinazolin-4-amine (96 mg) and *N,N*-dimethylglycine (27 mg) to give the title compound as a white solid in 73% yield; NMR Spectrum (DMSO- d_6) 9.90 (s, 1H), 8.40 (s, 1H), 8.15 (dd, 1H), 7.72-7.62 (m, 1H), 7.40 (t, 1H), 6.88 (d, 1H), 6.78 (d, 1H), 4.66-4.58 (m, 1H), 4.40 (dd, 1H), 4.19 (dd, 1H), 3.90 (s, 3H), 3.66-3.40 (m, 2H), 3.20-3.06 (m, 2H), 2.40 (s, 6H), 2.10-1.82 (m, 4H); Mass spectrum MH^+ 488.

Example 3

2-[(2*R*)-2-[(4-[3-Chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]pyrrolidin-1-yl]-2-oxoethanol

(Process (a))



The procedure described in Example 1 was repeated using *N*-(3-chloro-4-fluorophenyl)-7-methoxy-5-[(2*R*)-pyrrolidin-2-ylmethoxy]-quinazolin-4-amine (90 mg) and glycolic acid (19 mg) to give the title compound in 47% yield; NMR Spectrum (DMSO- d_6) 9.90 (s, 1H), 8.42 (s, 1H), 8.15 (dd, 1H), 7.70-7.60 (m, 1H), 7.40 (t, 1H), 6.90 (d, 1H), 6.78 (d,

1H), 4.65-4.53 (m, 2H), 4.42-4.37 (m, 1H), 4.20-4.11 (m, 1H), 4.03 (d, 1H), 3.90 (s, 3H), 3.45-3.35 (m, 3H), 2.10-1.82 (m, 4H); Mass spectrum MH^+ 461.


The *N*-(3-chloro-4-fluorophenyl)-7-methoxy-5-[(2*R*)-pyrrolidin-2-ylmethoxy]-quinazolin-4-amine used as starting material was prepared in 79% yield by reacting 4-[(3-chloro-4-fluorophenyl)-amino]-7-methoxy-quinazolin-5-ol (obtained as described in Example 1, preparation of starting materials) and *tert*-butyl (2*R*)-(+)-2-(hydroxymethyl)pyrrolidine-1-carboxylate using an analogous procedure to that described in the preparation of *N*-(3-chloro-4-fluorophenyl)-7-methoxy-5-[(2*S*)-pyrrolidin-2-ylmethoxy]-quinazolin-4-amine in Example 1; Mass spectrum MH^+ 403.

10

Example 4

N-(3-Chloro-4-fluorophenyl)-5-[(2*R*)-1-[(dimethylamino)acetyl]pyrrolidin-2-yl]methoxy-7-methoxyquinazolin-4-amine

(Process (a))



15


The procedure described in Example 1 was repeated using *N*-(3-chloro-4-fluorophenyl)-7-methoxy-5-[(2*R*)-pyrrolidin-2-ylmethoxy]-quinazolin-4-amine (90 mg) (obtained as described in Example 3, preparation of starting materials) and *N,N*-dimethylglycine (25 mg) to give the title compound as a white solid in 53% yield; NMR Spectrum (DMSO- d_6) 9.90 (s, 1H), 8.40 (s, 1H), 8.15 (dd, 1H), 7.72-7.62 (m, 1H), 7.40 (t, 1H), 6.86 (d, 1H), 6.78 (d, 1H), 4.64-4.56 (m, 1H), 4.38 (dd, 1H), 4.18 (dd, 1H), 3.90 (s, 3H), 3.58-3.50 (m, 2H), 3.15 (q, 2H), 2.20 (s, 6H), 2.10-1.82 (m, 4H); Mass spectrum MH^+ 488.

20

Example 5

2-[3-[(4-[3-Chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]pyrrolidin-1-yl]-2-oxoethanol

(Process (a))




The procedure described in Example 1 was repeated using *N*-(3-chloro-4-fluorophenyl)-7-methoxy-5-(pyrrolidin-3-ylmethoxy)quinazolin-4-amine (100 mg) and glycolic acid (21 mg) to give the title product in 45% yield; NMR spectrum (DMSO- d_6 at 373K) 9.69 (s, 1H), 8.52 (s, 1H), 8.21-8.20 (m, 1H), 7.66-7.63 (m, 1H), 7.40 (t, 1H), 6.90 (d, 1H), 6.83 (d, 1H), 4.41-4.39 (m, 2H), 4.20-4.10 (m, 1H), 4.05-4.00 (m, 2H), 3.97 (s, 3H), 3.73 (dd, 1H), 3.61-3.58 (m, 1H), 3.51-3.47 (m, 1H), 3.42 (dd, 1H), 2.26-2.14 (br m, 1H), 2.00-1.87 (br m, 1H); Mass spectrum MH^+ 461.

The *N*-(3-chloro-4-fluorophenyl)-7-methoxy-5-(pyrrolidin-3-ylmethoxy)quinazolin-4-amine used as starting material was prepared in 80% yield by reacting 4-[(3-chloro-4-fluorophenyl)-amino]-7-methoxyquinazolin-5-ol (obtained as described in Example 1, preparation of starting materials) and *tert*-butyl 3-(hydroxymethyl)pyrrolidine-1-carboxylate using an analogous procedure to that described in the preparation of *N*-(3-chloro-4-fluorophenyl)-7-methoxy-5-[(2*S*)-pyrrolidin-2-ylmethoxy]quinazolin-4-amine in Example 1; NMR spectrum ($CDCl_3$) 9.83 (s, 1H), 8.56 (s, 1H), 7.89 (dd, 1H), 7.46-7.41 (m, 1H), 7.15 (t, 1H), 6.86 (d, 1H), 6.50 (d, 1H), 4.15-4.12 (m, 2H), 3.93 (s, 3H), 3.50 (s, 1H), 3.20 (dd, 1H), 3.10-3.02 (m, 1H), 3.00-2.91 (m, 2H), 2.82-2.76 (m, 1H), 2.17-2.05 (m, 1H), 1.77-1.65 (m, 1H); Mass spectrum MH^+ 403.

20 Example 6

N-(3-Chloro-4-fluorophenyl)-7-methoxy-5-[(2*S*)-1-(pyrrolidin-1-ylacetyl)pyrrolidin-2-yl]methoxyquinazolin-4-amine

(Process (b))



25 Chloroacetyl chloride (21 μ l) was added in one portion to a stirred solution of *N*-(3-chloro-4-fluorophenyl)-7-methoxy-5-[(2*S*)-pyrrolidin-2-ylmethoxy]quinazolin-4-amine (96


mg) (obtained as described in Example 1, preparation of starting materials) and *N,N*-diisopropylethylamine (83 μ l) in DCM (3 ml) at room temperature. The reaction mixture was stirred for 40 minutes and then pyrrolidine (80 μ l) was added in one portion and the reaction mixture was stirred for 15 hours 30 minutes under a nitrogen atmosphere. The reaction

5 mixture was diluted with DCM (10 ml) and washed with saturated aqueous sodium hydrogen carbonate and water. The organic layer was dried (MgSO_4) and concentrated *in vacuo* to leave a yellow oil, which was purified by column chromatography using 0-10% methanol in DCM as eluent to give the title compound as a white solid (75 mg, 61 %); NMR Spectrum (DMSO- d_6) 9.90 (s, 1H), 8.40 (s, 1H), 8.10 (dd, 1H), 7.72-7.62 (m, 1H), 7.39 (t, 1H), 6.82 (d, 1H), 6.78 (d, 1H), 4.62-4.54 (m, 1H), 4.34 (dd, 1H), 4.16 (dd, 1H), 3.90 (s, 3H), 3.58-3.48 (m, 2H), 3.28-3.16 (m, 2H), 2.48 (br s, 4H), 2.10-1.82 (m, 4H), 1.6 (br s, 4H); Mass spectrum MH^+ 514.

Example 7

15 *N*-(3-Chloro-4-fluorophenyl)-7-methoxy-5-[(2*S*)-1-(morpholin-4-ylacetyl)pyrrolidin-2-yl]methoxy]quinazolin-4-amine

(Process (b))



The procedure described in Example 6 was repeated using *N*-(3-chloro-4-fluoro-


20 phenyl)-7-methoxy-5-[(2*S*)-pyrrolidin-2-ylmethoxy]-quinazolin-4-amine and morpholine to give the title compound as a white solid in 38% yield; NMR Spectrum (DMSO- d_6) 9.90 (s, 1H), 8.40 (s, 1H), 8.10 (dd, 1H), 7.72-7.62 (m, 1H), 7.39 (t, 1H), 6.82 (d, 1H), 6.78 (d, 1H), 4.62-4.54 (m, 1H), 4.34 (dd, 1H), 4.16 (dd, 1H), 3.90 (s, 3H), 3.58-3.50 (m, 2H), 3.50-3.42 (m, 4H), 3.10 (q, 2H), 2.40 (m, 4H), 2.08-1.82 (m, 4H); Mass spectrum MH^+ 530.

25

Example 8

N-(3-Chloro-4-fluorophenyl)-7-methoxy-5-[(2*S*)-1-[(4-methylpiperazin-1-yl)acetyl]pyrrolidin-2-yl]methoxy]quinazolin-4-amine

(Process (b))




- The procedure described in Example 6 was repeated using *N*-(3-chloro-4-fluorophenyl)-7-methoxy-5-[(2*S*)-pyrrolidin-2-ylmethoxy]-quinazolin-4-amine and *N*-methylpiperazine to give the title compound as a white solid in 36% yield; NMR Spectrum (DMSO-*d*₆) 9.90 (s, 1H), 8.40 (s, 1H), 8.15 (dd, 1H), 7.72-7.62 (m, 1H), 7.40 (t, 1H), 6.88 (d, 1H), 6.78 (d, 1H), 4.60-4.52 (m, 1H), 4.34 (dd, 1H), 4.16 (dd, 1H), 3.88 (s, 3H), 3.58-3.50 (m, 2H), 3.08 (q, 2H), 2.46-2.20 (m, 8H), 2.15 (s, 3H), 2.05-1.82 (m, 4H); Mass spectrum MH⁺ 543.

10 Example 9

(3*S*)-1-(2-[(2*S*)-2-[(4-[3-Chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]pyrrolidin-1-yl)-2-oxoethyl)pyrrolidin-3-ol

(Process (b))




- The procedure described in Example 6 was repeated using *N*-(3-chloro-4-fluorophenyl)-7-methoxy-5-[(2*S*)-pyrrolidin-2-ylmethoxy]-quinazolin-4-amine and (*S*)-3-pyrrolidinol to give the title compound as a white solid in 38% yield; Mass spectrum MH⁺ 530.

20 Example 10

N-(3-Chloro-4-fluorophenyl)-7-methoxy-5-[(2*S*)-1-[(2-methoxyethyl)(methyl)amino]-acetyl]pyrrolidin-2-ylmethoxy]quinazolin-4-amine

(Process (b))




The procedure described in Example 6 was repeated using *N*-(3-chloro-4-fluorophenyl)-7-methoxy-5-[(2*S*)-pyrrolidin-2-ylmethoxy]-quinazolin-4-amine and *N*-(2-methoxyethyl)methylamine to give the title compound as a white solid in 34% yield; NMR

- 5 Spectrum (DMSO-*d*₆) 9.90 (s, 1H), 8.40 (s, 1H), 8.10 (dd, 1H), 7.72-7.62 (m, 1H), 7.40 (t, 1H), 6.84 (d, 1H), 6.78 (d, 1H), 4.63-4.57 (m, 1H), 4.35 (dd, 1H), 4.15 (dd, 1H), 3.87 (s, 3H), 3.54 (t, 2H), 3.32 (t, 2H), 3.30-3.15 (m, 2H), 3.15 (s, 3H), 2.56 (t, 2H), 2.20 (s, 3H), 2.04-1.82 (m, 4H); Mass spectrum MH⁺ 532.

10 Example 11

N-(3-Chloro-4-fluorophenyl)-7-methoxy-5-[[[(2*S*)-1-[(2-hydroxyethyl)(methyl)amino]-acetyl]pyrrolidin-2-yl]methoxy]quinazolin-4-amine

(Process (b))




- 15 The procedure described in Example 6 was repeated using *N*-(3-chloro-4-fluorophenyl)-7-methoxy-5-[(2*S*)-pyrrolidin-2-ylmethoxy]-quinazolin-4-amine and 2-(methylamino)ethanol to give the title compound as a white solid in 39% yield; NMR
Spectrum (DMSO-*d*₆) 9.90 (s, 1H), 8.40 (s, 1H), 8.13 (dd, 1H), 7.70-7.62 (m, 1H), 7.40 (t, 1H), 6.85-6.77 (m, 2H), 4.62-4.54 (m, 1H), 4.40-4.28 (m, 2H), 4.14 (dd, 1H), 3.90 (s, 3H),
 20 3.55 (t, 2H), 3.44-3.37 (m, 2H), 3.24-3.10 (m, 2H), 2.54-2.40 (m, 2H), 2.20 (s, 3H), 2.10-1.82 (m, 4H); Mass spectrum MH⁺ 518.

Example 12

N-(3-Chloro-4-fluorophenyl)-7-methoxy-5-[[[(2*R*)-1-(methoxyacetyl)pyrrolidin-2-

- 25 yl]methoxy]quinazolin-4-amine

(Process (a))




Methoxyacetic acid (27 mg) and 4-(4,6-dimethoxy[1.3.5]triazin-2-yl)-4-methylmorpholinium chloride hydrate (103 mg) were added sequentially, each in one portion, to a stirred mixture of *N*-(3-chloro-4-fluorophenyl)-7-methoxy-5-[(2*R*)-pyrrolidin-2-ylmethoxy]-quinazolin-4-amine (100 mg) in DCM (2 ml) at room temperature. The reaction mixture was stirred for 1 hour and then diluted with DCM (10 ml) and washed with saturated aqueous sodium hydrogen carbonate. The organic layer was dried (MgSO₄) and concentrated *in vacuo* to leave a yellow solid, which was triturated with 1:1 ether/hexanes and filtered to give the title compound as a white solid (97 mg, 82%); NMR Spectrum (DMSO-d₆) 9.90 (br s, 1H), 8.40 (s, 1H), 8.15 (m, 1H), 7.65 (m, 1H), 7.40 (m, 1H), 6.80 (m, 2H), 4.60 (m, 1H), 4.50 (m, 1H), 4.25 (m, 1H), 4.0 (m, 3H), 3.95 (s, 3H), 3.50 (m, 2H), 3.20 (m, 2H), 2.0-1.85 (m, 4H); Mass spectrum MH⁺ 475.

The *N*-(3-chloro-4-fluorophenyl)-7-methoxy-5-[(2*R*)-pyrrolidin-2-ylmethoxy]-quinazolin-4-amine used as starting material was prepared as described in Example 3.

Example 13

5-[[[(2*R*)-1-Acetylpyrrolidin-2-yl]methoxy]-*N*-(3-chloro-4-fluorophenyl)-7-methoxyquinazolin-4-amine

(Process (a))



N,N-Diisopropylethylamine (52 μ l) and acetyl chloride (21 μ l) were added sequentially, each in one portion, to a stirred mixture of *N*-(3-chloro-4-fluorophenyl)-7-methoxy-5-[(2*R*)-pyrrolidin-2-ylmethoxy]-quinazolin-4-amine (100 mg) in DCM (2 ml) at room temperature. The reaction mixture was stirred for 1 hour and then diluted with DCM (10 ml) and washed with saturated aqueous sodium hydrogen carbonate. The organic layer was dried (MgSO₄) and concentrated *in vacuo* to leave a yellow solid, which was purified by

column chromatography using 2-4% methanol in DCM as eluent, followed by trituration with ether to give the title compound as a white solid (92 mg, 83%); NMR Spectrum (DMSO-d₆) 9.90 (br s, 1H), 8.40 (s, 1H), 8.15 (m, 1H), 7.65 (m, 1H), 7.40 (m, 1H), 6.80 (m, 2H), 4.60 (m, 1H), 4.50 (m, 1H), 4.25 (m, 1H), 3.95 (s, 3H), 3.50 (m, 1H), 3.30 (s, 3H), 2.0-1.85 (m, 5H);


5 Mass spectrum MH⁺ 445.

The *N*-(3-chloro-4-fluorophenyl)-7-methoxy-5-[(2*R*)-pyrrolidin-2-ylmethoxy]-quinazolin-4-amine used as starting material was prepared as described in Example 3.

Example 14

10 *N*-(3-Chloro-4-fluorophenyl)-7-methoxy-5-[(2*R*)-1-(morpholin-4-ylacetyl)pyrrolidin-2-yl]methoxy}quinazolin-4-amine

(Process (b))




Chloroacetyl chloride (77 μ l) was added in one portion to a stirred solution of *N*-(3-chloro-4-fluorophenyl)-7-methoxy-5-[(2*R*)-pyrrolidin-2-ylmethoxy]-quinazolin-4-amine (300 mg) and *N,N*-diisopropylethylamine (168 μ l) in DCM (4 ml) at room temperature. The reaction mixture was stirred for 1 hour and then was split into two equal portions. Morpholine (2 ml) was added to one portion and the reaction mixture was stirred at room temperature for 2 hours. The reaction mixture was diluted with DCM (10 ml) and washed with saturated aqueous sodium hydrogen carbonate. The organic layer was dried (MgSO₄) and concentrated *in vacuo* to leave a yellow oil, which was purified by column chromatography using 2-10% methanol in DCM with 1% 7 N ammonia in methanol as eluent to give the title compound as a white solid (142 mg, 72 %); NMR Spectrum (DMSO-d₆ at 373 K) 9.80 (br s, 1H), 8.40 (s, 1H), 8.15 (m, 1H), 7.65 (m, 1H), 7.33 (m, 1H), 6.80 (m, 2H), 4.60 (m, 1H), 4.50 (m, 1H), 4.25 (m, 1H), 3.95 (s, 3H), 3.60-3.45 (m, 6H), 3.20-3.0 (m, 2H), 2.40 (m, 4H), 2.10-1.85 (m, 4H); Mass spectrum MH⁺ 530.

The *N*-(3-chloro-4-fluorophenyl)-7-methoxy-5-[(2*R*)-pyrrolidin-2-ylmethoxy]-quinazolin-4-amine used as starting material was prepared as described in Example 3.

Example 15**N-(3-Chloro-4-fluorophenyl)-7-methoxy-5-[(2R)-1-[(4-methylpiperazin-1-yl)acetyl]pyrrolidin-2-yl]methoxy)quinazolin-4-amine**


(Process (b))



The procedure described in Example 14 was repeated using *N*-(3-chloro-4-fluorophenyl)-7-methoxy-5-[(2*R*)-pyrrolidin-2-ylmethoxy]-quinazolin-4-amine and 4-methylpiperazine to give the title compound as a white solid in 88% yield; NMR Spectrum (DMSO-*d*₆ at 373K); 9.80 (br s, 1H), 8.40 (s, 1H), 8.15 (m, 1H), 7.65 (m, 1H), 7.33 (m, 1H), 6.80 (m, 2H), 4.60 (m, 1H), 4.50 (m, 1H), 4.25 (m, 1H), 3.95 (s, 3H), 3.70 (m, 1H), 3.50 (m, 1H), 3.20-3.0 (m, 2H), 2.45 (m, 4H), 2.25 (m, 4H), 2.10 (m, 3H), 2.0-1.85 (m, 4H); Mass spectrum MH⁺ 543.

Example 16**2-[(2R)-2-[[4-[3-Chloro-4-fluoroanilino]-7-(2-hydroxyethoxy)quinazolin-5-yl]oxy)methyl]pyrrolidin-1-yl]-2-oxoethanol**

(Process (d))



2-Bromoethanol (51 μ l) and potassium carbonate (86 mg) were added to a mixture of 4-[3-chloro-4-fluoroanilino]-5-[[[(2*R*)-1-glycoloylpyrrolidin-2-yl]methoxy]-quinazolin-7-ol (70 mg) in DMF and the reaction mixture was heated at 70°C for 16 hours. The reaction mixture was cooled to room temperature and then water (5 ml) was added to leave a white precipitate. The precipitate was filtered, washed with water and then purified by column chromatography using 2-5% methanol in DCM as eluent. The product was purified further by acidic reverse phase HPLC, and then the combined fractions were reduced to a third of the

initial concentration, basified with sodium carbonate and extracted with ethyl acetate. The organic layer was dried (Na₂SO₄) and concentrated *in vacuo* to give the title compound as a white solid (35 mg, 45%); NMR Spectrum (DMSO-d₆ at 373K) 9.80 (br s, 1H), 8.40 (s, 1H), 8.15 (m, 1H), 7.65 (m, 1H), 7.33 (m, 1H), 6.80 (m, 2H), 4.60 (m, 1H), 4.50 (m, 1H), 4.30-4.10 (m, 4H), 4.0 (m, 2H), 3.75 (m, 2H), 3.60 (m, 1H), 3.50-3.37 (m, 2H), 2.10-1.85 (m, 4H); Mass spectrum MH⁺ 491.

The 4-[3-chloro-4-fluoroanilino]-5-[[*(2R)*-1-glycoloylpyrrolidin-2-yl]-methoxy]-quinazolin-7-ol used as starting material was prepared as follows:

Di-*tert*-butyl azodicarboxylate (11.0 g) was added portionwise, over 5 minutes, to a stirred solution of [7-(benzyloxy)-5-hydroxy-4-oxoquinazolin-3(*4H*)-yl]methyl pivalate (12.2 g) (obtained as described in WO01/94341 Example 15, foot note [8]), *tert*-butyl (*2R*)-(-)-2-(hydroxymethyl)pyrrolidine-1-carboxylate (7.4 g) and triphenylphosphine (11.7 g) in DCM (100 ml) at 0 °C under a nitrogen atmosphere. The mixture mixture was warmed to room temperature and stirred for 16 hours. The mixture was concentrated *in vacuo* to leave the crude *tert*-butyl (*2R*)-2-[[7-(benzyloxy)-3-[[*(2,2*-dimethylpropanoyl)oxy]methyl]-4-oxo-3,4-dihydroquinazolin-5-yl]oxy]methyl]pyrrolidine-1-carboxylate, which was used without further purification; Mass spectrum MH⁺ 566.

The crude *tert*-butyl (*2R*)-2-[[7-(benzyloxy)-3-[[*(2,2*-dimethylpropanoyl)oxy]methyl]-4-oxo-3,4-dihydroquinazolin-5-yl]oxy]methyl]pyrrolidine-1-carboxylate (assumed 18 g) was dissolved in 7N ammonia in methanol and stirred at room temperature for 16 hours. The reaction mixture was concentrated *in vacuo* and purified by column chromatography using 0-10% methanol in DCM as eluent to give a mixture of two products. The mixture was dissolved in DCM, washed with 20% aqueous sodium hydroxide solution (2x) and then dried (MgSO₄) and concentrated *in vacuo* to leave *tert*-butyl (*2R*)-2-[[7-(benzyloxy)-4-oxo-3,4-dihydroquinazolin-5-yl]oxy]methyl]pyrrolidine-1-carboxylate as a beige foam (6.8 g, 48% for the two steps); Mass spectrum MH⁺ 452.

Phosphorous oxychloride (520 µl) was added dropwise to a mixture of *tert*-butyl (*2R*)-2-[[7-(benzyloxy)-4-oxo-3,4-dihydroquinazolin-5-yl]oxy]methyl]pyrrolidine-1-carboxylate (500 mg) and *N,N*-diisopropylethylamine (52 µl) in DCM (10 ml) and the reaction mixture was heated at 50°C for 62 hours. The reaction mixture was concentrated *in vacuo* and then water and saturated aqueous sodium carbonate (cooled to 4°C) was added. The aqueous mixture was then extracted with DCM and the organic layer was dried (MgSO₄) and

concentrated *in vacuo* to leave *tert*-butyl (2*R*)-2-([7-(benzyloxy)-4-chloroquinazolin-5-yl]oxy)methylpyrrolidine-1-carboxylate as a brown oil, which was used without further purification.

The crude *tert*-butyl (2*R*)-2-([7-(benzyloxy)-4-chloroquinazolin-5-yl]oxy)methylpyrrolidine-1-carboxylate was taken up in THF (5 ml) and 3-chloro-4-fluoroaniline (160 mg) and *N,N*-diisopropylethylamine (580 μ l) were each added in one portion and the solution was heated at 70°C for 18 hours. The reaction mixture was cooled to room temperature and concentrated *in vacuo* to leave a thick brown oil. Purification by flash chromatography, using 1.5% methanol in DCM as eluent, gave the *tert*-butyl (2*R*)-2-([7-(benzyloxy)-4-[3-chloro-4-fluoroanilino]quinazolin-5-yl]oxy)methylpyrrolidine-1-carboxylate as a beige foam (350 mg, 54% over 2 steps); Mass spectrum MH^+ 481.

tert-Butyl (2*R*)-2-([7-(benzyloxy)-4-[3-chloro-4-fluoroanilino]quinazolin-5-yl]oxy)methylpyrrolidine-1-carboxylate (1.5g) was taken up in trifluoroacetic acid (15 ml) and the solution was heated at 70°C for 18 hours. The solution was cooled to room temperature and then concentrated *in vacuo* to leave a brown residue. Saturated aqueous sodium hydrogen carbonate was added to the residue and the aqueous mixture was extracted with ethyl acetate. The layers were separated and the organic layer was dried ($MgSO_4$) and concentrated *in vacuo* to leave a brown oil. Purification by flash chromatography, using 5:1:84 methanol:7N ammonia in methanol:DCM \rightarrow 20:1:79 methanol:7N ammonia in methanol:DCM as eluent, followed by trituration with ether gave 4-[3-chloro-4-fluoroanilino]-5-[(2*R*)-pyrrolidin-2-ylmethoxy]quinazolin-7-ol as an off-white solid (200 mg, 20%); NMR Spectrum (DMSO- d_6) 10.40 (brs, 1H), 8.40 (s, 1H), 8.15 (m, 1H), 7.95 (m, 1H), 7.40 (m, 1H), 6.75 (m, 2H), 4.30 (m, 1H), 4.05 (m, 1H), 3.60-3.50 (m, 1H), 3.0-2.80 (m, 2H), 1.95-1.80 (m, 1H), 1.75-1.65 (m, 2H), 1.50-1.40 (m, 1H); Mass spectrum MH^+ 391.


Glycolic acid (50 mg) and *O*-(7-azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (HATU, 250 mg) were added sequentially, each in one portion, to a stirred solution of 4-[3-chloro-4-fluoroanilino]-5-[(2*R*)-pyrrolidin-2-ylmethoxy]quinazolin-7-ol (197 mg) and *N,N*-diisopropylethylamine (177 μ l) in DCM (4 ml) at room temperature. The reaction mixture was stirred for 60 hours and then an additional portion of glycolic acid (20 mg) and *O*-(7-azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (HATU, 100 mg) were added and the mixture was stirred for a further 5 hours. The reaction mixture was then diluted with DCM (10 ml) and washed with saturated aqueous sodium

hydrogen carbonate. The organic layer was dried (MgSO₄) and concentrated *in vacuo* to leave a yellow oil, which was purified by column chromatography using 5% methanol in DCM as eluent to give the 4-[3-chloro-4-fluoro-anilino]-5-[[*(2R)*-1-glycoloylpyrrolidin-2-yl]-methoxy]-quinazolin-7-ol as a grey solid (96 mg, 42%); NMR Spectrum (DMSO-d₆ at 373K) 10.10 (brs, 1H), 9.60 (brs, 1H), 8.40 (s, 1H), 8.15 (m, 1H), 7.65 (m, 1H), 7.40 (m, 1H), 6.70 (s, 2H), 4.60 (m, 1H), 4.40 (m, 1H), 4.20 (m, 1H), 4.05 (m, 2H), 3.55-3.40 (m, 2H), 2.15-1.85 (m, 4H); Mass spectrum MH⁺ 449.

Example 17

10 2-[(2*R*)-2-({[4-[3-Chloro-4-fluoroanilino]-7-(2-methoxyethoxy)quinazolin-5-yl]oxy)methyl]pyrrolidin-1-yl]-2-oxoethanol

(Process (d))




2-Bromoethyl methyl ether (22 μ l) and potassium carbonate (43 mg) were added to a mixture of 4-[3-chloro-4-fluoroanilino]-5-[[*(2R)*-1-glycoloylpyrrolidin-2-yl]-methoxy]-quinazolin-7-ol (70 mg) in DMF and the reaction mixture was heated at 40°C for 16 hours. The reaction mixture was cooled to room temperature and then water (5 ml) was added to leave a white precipitate. The precipitate was filtered, washed with water and then purified by column chromatography using 2-5% methanol in DCM as eluent to give the title compound as a white solid (51 mg, 64%); NMR Spectrum (DMSO-d₆ at 373K) 9.80 (br s, 1H), 8.40 (s, 1H), 8.15 (m, 1H), 7.65 (m, 1H), 7.33 (m, 1H), 6.80 (m, 2H), 4.60 (m, 1H), 4.50 (m, 1H), 4.30-4.10 (m, 4H), 4.0 (m, 2H), 3.75 (t, 2H), 3.50-3.37 (m, 2H), 3.35 (s, 3H), 2.10-1.85 (m, 4H); Mass spectrum MH⁺ 505.

The 4-[3-chloro-4-fluoroanilino]-5-[[*(2R)*-1-glycoloylpyrrolidin-2-yl]-methoxy]-quinazolin-7-ol used as starting material was prepared as described in Example 16.

Example 18

2-[(2*R*)-2-({[4-[3-Chloro-4-fluoroanilino]-7-(2-ethoxyethoxy)quinazolin-5-yl]oxy)methyl]pyrrolidin-1-yl]-2-oxoethanol

30 (Process (d))




The procedure described in Example 17 was repeated using 4-[3-chloro-4-fluoro-anilino]-5-[(2*R*)-1-glycoloylpyrrolidin-2-yl]-methoxy}-quinazolin-7-ol (70 mg) and 2-bromoethyl ethyl ether (26 μ l) to give the title compound as a white solid in 48% yield; NMR Spectrum (DMSO- d_6 at 373K) 9.80 (br s, 1H), 8.40 (s, 1H), 8.15 (m, 1H), 7.65 (m, 1H), 7.33 (m, 1H), 6.80 (m, 2H), 4.60 (m, 1H), 4.50 (m, 1H), 4.30-4.10 (m, 4H), 4.0 (m, 2H), 3.75 (t, 2H), 3.55 (q, 2H), 3.48-3.37 (m, 2H), 2.10-1.85 (m, 4H), 1.15 (t, 3H); Mass spectrum MH^+ 519.

10 Example 19

2-[(2*R*)-2-[(4-[3-Chloro-4-fluoroanilino]-7-ethoxyquinazolin-5-yl)oxy]methyl]-pyrrolidin-1-yl]-2-oxoethanol

(Process (a))



15 Glycolic acid (22 mg) and *O*-(7-azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (HATU, 110 mg) were added sequentially, each in one portion, to a stirred solution of *N*-(3-chloro-4-fluorophenyl)-7-ethoxy-5-[(2*R*)-pyrrolidin-2-ylmethoxy]-quinazolin-4-amine (81 mg) and *N,N*-diisopropylethylamine (68 μ l) in DCM (2.5 ml) at room temperature. The reaction mixture was stirred for 16 hours and then concentrated *in vacuo* to
 20 leave a brown residue. The residue was taken up in a 7:2:1 mixture of DMSO:acetonitrile: water, filtered and then purified by reverse phase HPLC. The fractions containing the required product were combined, basified by the addition of potassium carbonate and then extracted with DCM. The organic layer was dried ($MgSO_4$) and then concentrated *in vacuo* to give the title compound as a white solid (36 mg, 39%); NMR Spectrum (DMSO- d_6 at 373K)
 25 9.80 (br s, 1H), 8.40 (s, 1H), 8.15 (m, 1H), 7.65 (m, 1H), 7.33 (m, 1H), 6.80 (m, 2H), 4.60 (m, 1H), 4.50 (m, 1H), 4.30-4.10 (m, 4H), 4.0 (m, 2H), 3.50-3.40 (m, 2H), 2.10-1.85 (m, 4H), 1.40 (t, 3H); Mass spectrum MH^+ 475.

The *N*-(3-chloro-4-fluorophenyl)-7-ethoxy-5-[(2*R*)-pyrrolidin-2-ylmethoxy]-quinazolin-4-amine used as starting material was prepared as follows:


A mixture of *tert*-butyl (2*R*)-2-[(4-[3-chloro-4-fluoroanilino]-quinazolin-5-yl)oxy)methyl]pyrrolidine-1-carboxylate (230 mg) (obtained as described in Example 16) and 5% palladium on carbon in ethanol (5 ml) was stirred under hydrogen (5 bar pressure) at 25°C for 16 hours. The mixture was then filtered and concentrated *in vacuo* to leave the *tert*-butyl (2*R*)-2-[(4-[3-chloro-4-fluoroanilino]-7-hydroxyquinazolin-5-yl)oxy)methyl]pyrrolidine-1-carboxylate as a light brown oil, which was used without further purification.

10 The crude *tert*-butyl (2*R*)-2-[(4-[3-chloro-4-fluoroanilino]-7-hydroxyquinazolin-5-yl)oxy)methyl]pyrrolidine-1-carboxylate (120 mg) was dissolved in DMA (1 ml), and bromoethane (27 µl) and potassium carbonate (68 mg) were each added in one portion. The reaction mixture was heated at 40°C for 16 hours and then cooled to room temperature. Trifluoroacetic acid (0.5 ml) was then added and the reaction mixture was stirred for 4 hours
15 at room temperature. Further trifluoroacetic acid (1 ml) was added and the reaction mixture was stirred at room temperature for 16 hours. The reaction mixture was then poured onto saturated aqueous sodium hydrogen carbonate and then extracted with ethyl acetate. The organic layer was dried (MgSO₄) and then concentrated *in vacuo* to leave a green oil. Purification by flash chromatography, using 2-5% methanol in DCM as eluent, gave the *N*-(3-
20 chloro-4-fluorophenyl)-7-ethoxy-5-[(2*R*)-pyrrolidin-2-ylmethoxy]quinazolin-4-amine as a pale green solid (83 mg, 52% over three steps); NMR Spectrum (DMSO-d₆) 10.2 (s, 1H), 9.50 (s, 1H), 8.30-8.20 (m, 1H), 7.90-7.80 (m, 1H), 7.50-7.40 (m, 1H), 6.80-6.70 (m, 1H), 4.50-4.40 (m, 1H), 4.30-4.20 (m, 1H), 4.30-4.10 (m, 3H), 3.90-3.80 (m, 1H), 3.15-3.05 (m, 2H), 2.10-2.00 (m, 1H), 1.90-1.80 (m, 2H), 1.70-1.60 (m, 1H), 1.40 (t, 3H); Mass spectrum
25 MH⁺ 417.

Example 20

(3*S*,5*R*)-5-[(4-[3-Chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]-1-glycoloylpyrrolidin-3-ol

30 (Process (a))



- O*-(7-Azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (HATU, 171 mg) was added in one portion to a stirred solution of (3*S*,5*R*)-5-[(4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]pyrrolidin-3-ol (157 mg), *N,N*-diisopropylethylamine (131 μ l) and glycolic acid (31 μ l) in DCM (5 ml), under an atmosphere of nitrogen. The reaction mixture was stirred at room temperature for 4 hours and then diluted with DCM and saturated aqueous sodium hydrogen carbonate was added. The organic layer was separated, dried (Na_2SO_4) and then concentrated *in vacuo* to leave the crude product as a yellow waxy solid. Purification by column chromatography, using 0 to 10% methanol in DCM as eluent, gave the title compound as a white solid (144 mg, 80%); NMR spectrum (DMSO- d_6) 10.17 (s, 1H), 8.53 (s, 1H), 8.14 (dd, 1H), 7.74-7.69 (m, 1H), 7.47 (t, 1H), 6.92 (d, 1H), 6.83 (d, 1H), 5.14-5.10 (m, 1H), 4.71-4.67 (m, 1H), 4.50-4.44 (m, 1H), 4.40-4.38 (m, 1H), 4.28-4.23 (m, 1H), 4.06-4.03 (m, 2H), 3.95 (s, 3H), 3.53-3.48 (m, 2H), 2.05-2.02 (m, 2H); Mass spectrum MH^+ 477.
- 15 The (3*S*,5*R*)-5-[(4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]pyrrolidin-3-ol used as starting material was prepared as follows:
- Di-*tert*-butylazodicarboxylate (DTAD, 623 mg) was added in one portion to a mixture of 4-[(3-chloro-4-fluorophenyl)-amino]-7-methoxyquinazolin-5-ol (578 mg, prepared as described in Example 1, preparation of starting materials), *tert*-butyl (2*R*,4*S*)-4-[[*tert*-butyl(dimethyl)silyl]oxy]-2-(hydroxymethyl)pyrrolidine-1-carboxylate (750 mg), and triphenylphosphine (710 mg) in DCM (100 ml) at room temperature, under an atmosphere of nitrogen. The reaction mixture was stirred at room temperature for 24 hours and then concentrated *in vacuo* to leave a brown oil. Purification by column chromatography, using 0-50% ethyl acetate in hexane as eluent, gave *tert*-butyl (2*R*,4*S*)-4-[[*tert*-butyl(dimethyl)silyl]oxy]-2-[(4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]-pyrrolidine-1-carboxylate as a white solid (671 mg, 59%); NMR spectrum (DMSO- d_6) 8.46 (s, 1H), 8.03 (br s, 1H), 7.67 (br s, 1H), 7.43 (t, 1H), 6.85-6.81 (m, 2H), 4.57-4.20 (m, 4H), 3.94 (s, 3H), 3.47-3.40 (m, 1H), 2.10-2.00 (m, 2H), 1.33 (m, 10H), 0.86 (s, 9H), 0.07 (s, 6H); Mass Spectrum MH^+ 633.
- 25

tert-Butyl (2*R*,4*S*)-4-{[*tert*-butyl(dimethyl)silyl]oxy}-2-[(4-[3-chloro-4-fluoro-anilino]-7-methoxyquinazolin-5-yl)oxy)methyl]pyrrolidine-1-carboxylate (636 mg) was dissolved in trifluoroacetic acid (30 ml) and stirred at room temperature for 72 hours. The reaction mixture was then concentrated *in vacuo*, and excess water was added, followed by the
5 careful addition of saturated aqueous sodium hydrogen carbonate. The mixture was extracted with 3% methanol in DCM and the organic layer was separated, dried (Na₂SO₄) and concentrated *in vacuo* to leave a light brown foam. Titration with cold acetonitrile gave (3*S*,5*R*)-5-[(4-[3-chloro-4-fluoroanilino]-7-methoxy-quinazolin-5-yl)oxy)methyl]-pyrrolidin-3-ol as an off-white solid (370 mg, 88%); NMR spectrum (DMSO-d₆) 10.50-10.00 (br, 1H),
10 8.49 (s, 1H), 8.25 (dd, 1H), 7.90-7.84 (m, 1H), 7.44 (t, 1H), 6.82 (d, 1H), 6.76 (d, 1H), 4.92 (br s, 1H), 4.38 (dd, 1H), 4.32 (br s, 1H), 3.92-3.86 (m, 4H), 3.77-3.23 (br s, 1H), 3.09 (dd, 1H), 2.83 (d, 1H), 1.88 (dd, 1H), 1.74-1.65 (m, 1H); Mass spectrum MH⁺ 419.

The *tert*-butyl (2*R*,4*S*)-4-{[*tert*-butyl(dimethyl)silyl]oxy}-2-(hydroxymethyl)-pyrrolidine-1-carboxylate used as starting material was prepared as follows:

15 (Trimethylsilyl)diazomethane (2*N* solution in hexane, 2.9 ml) was added dropwise over 5 minutes to a stirred solution of (4*S*)-1-(*tert*-butoxycarbonyl)-4-hydroxy-D-proline (1.0 g) in DCM (20 ml) and methanol (5 ml) at room temperature until a permanent faint yellow colour persisted. The reaction mixture was concentrated *in vacuo* and purified by column chromatography, using 0-50% ethyl acetate in hexane as eluent, to give 1-*tert*-butyl 2-methyl
20 (2*R*,4*S*)-4-hydroxypyrrolidine-1,2-dicarboxylate as a clear oil (1.03 g, 97%); NMR spectrum (CDCl₃) 4.90-4.37 (m, 2H), 3.73 (s, 3H), 3.64 (dd, 1H), 3.53-3.43 (m, 1H), 2.34-2.27 (m, 1H), 2.12-1.97 (m, 2H), 1.46 (s, 3H), 1.41 (s, 6H).


tert-Butyldimethylsilyl triflate (1.44 ml) was added to a stirred solution of 1-*tert*-butyl 2-methyl (2*R*,4*S*)-4-hydroxypyrrolidine-1,2-dicarboxylate (1.03 g) and triethylamine (1.76 ml)
25 in DCM (50 ml) at room temperature, under an atmosphere of nitrogen. The reaction mixture was stirred for 2 hours and then the mixture was washed with saturated aqueous potassium hydrogensulfate, dried (Na₂SO₄), and concentrated *in vacuo* to give a clear oil. Purification by column chromatography, using 0-30% ethyl acetate in hexane as eluent, gave 1-*tert*-butyl 2-methyl (2*R*,4*S*)-4-{[*tert*-butyl(dimethyl)silyl]oxy}pyrrolidine-1,2-dicarboxylate as a clear oil
30 (891 mg, 59%); NMR spectrum (CDCl₃) 4.43-4.32 (m, 2H), 3.75-3.72 (m, 2H), 3.65-3.56 (m, 1H), 3.43-3.31 (m, 1H), 2.21-2.15 (m, 1H), 2.07-1.98 (m, 1H), 1.47 (s, 3H), 1.42 (s, 6H), 0.88 (s, 9H), 0.07 (s, 6H).

Lithium borohydride (2M solution in THF, 6.19 ml) was added dropwise to a stirred solution of 1-*tert*-butyl 2-methyl (2*R*,4*S*)-4-[[*tert*-butyl(dimethyl)silyl]oxy]pyrrolidine-1,2-dicarboxylate (891 mg) in ether (100 ml) at 0°C under an atmosphere of nitrogen. The mixture was allowed to warm to room temperature and stirred for 3 hours. A saturated solution of potassium hydrogencarbonate (5 ml) was added slowly, followed by water (20 ml). The ether layer was separated, and the aqueous layer washed with DCM (3x). The combined organic layers were dried (Na₂SO₄) and concentrated *in vacuo* to give *tert*-butyl (2*R*,4*S*)-4-[[*tert*-butyl(dimethyl)silyl]oxy]-2-(hydroxymethyl)pyrrolidine-1-carboxylate as a clear oil (750 mg, 88%), which was used without further purification; NMR spectrum (CDCl₃) 4.82 (d, 1H), 4.26-4.28 (m, 1H), 4.13-4.15 (m, 1H), 3.67-3.73 (m, 1H), 3.55 (ddd, 1H), 3.42-3.46 (m, 1H), 3.35 (dd, 1H), 1.92-2.01 (m, 1H), 1.48 (s, 9H), 0.91-0.97 (m, 1H), 0.88 (s, 9H), 0.07 (s, 6H).

Example 21

15 (3*S*,5*R*)-5-[(4-[3-Chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]-1-(*N,N*-dimethylglycyl)pyrrolidin-3-ol

(Process (a))




The procedure described in Example 20 was repeated using (3*S*,5*R*)-5-[(4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]pyrrolidin-3-ol (185 mg) with *N,N*-dimethylglycine (50 mg) to give the title compound as a white solid in 73% yield; NMR spectrum (DMSO-*d*₆) 9.93 (s, 1H), 8.45 (s, 1H), 8.15 (dd, 1H), 7.76-7.70 (m, 1H), 7.45 (t, 1H), 6.83 (s, 2H), 5.09-5.08 (m, 1H), 4.70-4.66 (m, 1H), 4.45-4.40 (m, 2H), 4.28-4.23 (m, 1H), 3.93 (s, 3H), 3.62-3.49 (m, 2H), 3.20 (d, 1H), 3.00 (d, 1H), 2.21 (s, 6H), 2.06-2.02 (m, 2H); Mass spectrum MH⁺ 504.

Example 22

(3*S*,5*R*)-1-Acetyl-5-[(4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]pyrrolidin-3-ol

30 (Process (a))




The procedure described in Example 20 was repeated using (3*S*,5*R*)-5-[(4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]pyrrolidin-3-ol (120 mg) with acetic acid (18 μ l) to give the title compound as a white solid in 59% yield; NMR spectrum (DMSO- d_6) 9.95 (s, 1H), 8.46 (s, 1H), 8.16-8.14 (m, 1H), 7.73-7.70 (m, 1H), 7.46 (t, 1H), 6.85 (s, 1H), 6.84 (s, 1H), 5.10 (m, 1H), 4.67-4.63 (m, 1H), 4.47-4.38 (m, 2H), 4.26-4.21 (m, 1H), 3.94 (s, 3H), 3.59 (dd, 1H), 3.41 (dd, 1H), 2.11-2.05 (m, 2H), 2.00 (s, 3H); Mass spectrum MH^+ 461.

Example 23

10 (3*S*,5*R*)-5-[(4-[3-Chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]-1-(methoxyacetyl)pyrrolidin-3-ol

(Process (a))




The procedure described in Example 20 was repeated using (3*S*,5*R*)-5-[(4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]pyrrolidin-3-ol (120 mg) with methoxyacetic acid (24 μ l) to give the title compound as a white solid in 60% yield; NMR spectrum (DMSO- d_6) 9.96 (s, 1H), 8.45 (s, 1H), 8.17-8.14 (m, 1H), 7.76-7.72 (m, 1H), 7.45 (t, 1H), 6.83 (s, 2H), 5.10 (m, 1H), 4.73-4.69 (m, 1H), 4.47-4.38 (m, 2H), 4.25 (dd, 1H), 4.10-3.98 (m, 2H), 3.94 (s, 3H), 3.51 (dd, 1H), 3.40 (s, 1H), 3.29 (s, 3H), 2.05-2.02 (m, 2H); Mass spectrum MH^+ 491.

Example 24

25 (3*S*,5*R*)-5-[(4-[3-Chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]-1-(4-methylpiperazin-1-yl)acetylpyrrolidin-3-ol

(Process (a))




The procedure described in Example 20 was repeated using (3*S*,5*R*)-5-[(4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]pyrrolidin-3-ol (120 mg) with *N*-methylpiperazin-1-ylacetic acid (50 mg). Purification by column chromatography, using 0 to 10% 7*N* ammonia in methanol in DCM as eluent, gave the title compound as a white solid (82 mg, 51%); NMR spectrum (CDCl₃) 9.62 (s, 1H), 8.51 (s, 1H), 7.86-7.82 (m, 1H), 7.55-7.50 (m, 1H), 7.16-7.10 (m, 1H), 6.84 (br s, 1H), 6.54 (br s, 1H), 4.90-4.80 (m, 1H), 4.51 (m, 1H), 4.37-4.33 (m, 1H), 4.20-4.10 (m, 1H), 3.90 (s, 3H), 3.86-3.80 (m, 1H), 3.54-3.50 (m, 1H), 3.21 (d, 1H), 3.02 (d, 1H), 2.69-2.17 (br m, 13H), 2.05-1.90 (m, 1H); Mass spectrum MH⁺ 559.

Example 25

(3*R*,5*S*)-5-[(4-[3-Chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]-1-glycoloylpyrrolidin-3-ol

15 (Process (a))



The procedure described in Example 20 was repeated using (3*R*,5*S*)-5-[(4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]pyrrolidin-3-ol (209 mg) with glycolic acid (42 mg) to give the title compound as a white solid in 53% yield; NMR spectrum (DMSO-*d*₆) 10.17 (s, 1H), 8.53 (s, 1H), 8.14 (dd, 1H), 7.74-7.69 (m, 1H), 7.47 (t, 1H), 6.92 (d, 1H), 6.83 (d, 1H), 5.14-5.10 (m, 1H), 4.71-4.67 (m, 1H), 4.50-4.44 (m, 1H), 4.40-4.38 (m, 1H), 4.28-4.23 (m, 1H), 4.06-4.03 (m, 2H), 3.95 (s, 3H), 3.53-3.48 (m, 3H), 2.05-2.02 (m, 2H); Mass spectrum MH⁺ 477.

The (3*R*,5*S*)-5-[(4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]pyrrolidin-3-ol used as starting material was prepared as follows:

Di-*tert*-butylazodicarboxylate (DTAD, 692 mg) was added in one portion to a mixture of 4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-ol (640 mg), *tert*-butyl (2*S*,4*R*)-4-[[*tert*-butyl(dimethyl)silyl]oxy]-2-(hydroxymethyl)-pyrrolidine-1-carboxylate (830 mg), and triphenylphosphine (788 mg) in DCM (50 ml) at room temperature, under an atmosphere of nitrogen. The reaction mixture was stirred at room temperature for 24 hours and then concentrated *in vacuo* to leave a brown oil. Purification by column chromatography, using 0-50% ethyl acetate in hexane as eluent, gave *tert*-butyl (2*S*,4*R*)-4-[[*tert*-butyl(dimethyl)silyl]oxy]-2-[[4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl]oxy)methyl]-pyrrolidine-1-carboxylate as a white solid (884 mg, 70%); Mass spectrum MH^+ 633.


10 *tert*-Butyl (2*S*,4*R*)-4-[[*tert*-butyl(dimethyl)silyl]oxy]-2-[[4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl]oxy)methyl]pyrrolidine-1-carboxylate (860 mg) was dissolved in trifluoroacetic acid (40 ml) and stirred at room temperature for 24 hours. The reaction mixture was then concentrated *in vacuo*, and excess water was added, followed by the careful addition of saturated aqueous sodium hydrogen carbonate. The mixture was extracted with 3% methanol in DCM and the organic layer was separated, dried (Na₂SO₄) and concentrated *in vacuo* to leave a light brown foam. Trituration with cold acetonitrile gave (3*R*,5*S*)-5-[[4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl]oxy)methyl]-pyrrolidin-3-ol as an off-white solid (394 mg, 69%); NMR spectrum (DMSO-*d*₆) 10.50-10.00 (br, 1H), 8.49 (s, 1H), 8.25 (dd, 1H), 7.90-7.84 (m, 1H), 7.44 (t, 1H), 6.82 (d, 1H), 6.76 (d, 1H), 4.92 (br s, 1H), 4.38 (dd, 1H), 4.32 (br s, 1H), 3.92-3.86 (m, 4H), 3.77-3.23 (br s, 1H), 3.09 (dd, 20 1H), 2.83 (d, 1H), 1.88 (dd, 1H), 1.74-1.65 (m, 1H); Mass spectrum MH^+ 419.

The *tert*-butyl (2*S*,4*R*)-4-[[*tert*-butyl(dimethyl)silyl]oxy]-2-(hydroxymethyl)-pyrrolidine-1-carboxylate used as a starting material is commercially available or can be prepared using an analogous process to that described in Example 20 for the preparation of 25 *tert*-butyl (2*R*,4*S*)-4-[[*tert*-butyl(dimethyl)silyl]oxy]-2-(hydroxymethyl)-pyrrolidine-1-carboxylate but using (4*R*)-1-(*tert*-butoxycarbonyl)-4-hydroxy-L-proline.

Example 26

(3*R*,5*S*)-5-[[4-[3-Chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl]oxy)methyl]-1-(*N,N*-dimethylglycyl)pyrrolidin-3-ol

30 (Process (a))




The procedure described in Example 25 was repeated using (3*R*,5*S*)-5-[(4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]pyrrolidin-3-ol (185 mg) with *N,N*-dimethylglycine (50 mg) to give the title compound as a white solid in 65% yield; NMR spectrum (DMSO-*d*₆) 9.93 (s, 1H), 8.45 (s, 1H), 8.15 (dd, 1H), 7.76-7.70 (m, 1H), 7.45 (t, 1H), 6.83 (s, 2H), 5.09-5.08 (m, 1H), 4.70-4.66 (m, 1H), 4.45-4.40 (m, 2H), 4.28-4.23 (m, 1H), 3.93 (s, 3H), 3.62-3.49 (m, 2H), 3.20 (d, 1H), 3.00 (d, 1H), 2.21 (s, 6H), 2.06-2.02 (m, 2H); Mass spectrum MH⁺ 504.

10 Example 27

(3*R*,5*S*)-1-Acetyl-5-[(4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]pyrrolidin-3-ol

(Process (a))




The procedure described in Example 25 was repeated using (3*R*,5*S*)-5-[(4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]pyrrolidin-3-ol (209 mg) with acetic acid (32 μ l) to give the title compound as a white solid in 92% yield; NMR spectrum (DMSO-*d*₆) 9.98 (s, 1H), 8.47 (s, 1H), 8.16-8.14 (m, 1H), 7.73-7.70 (m, 1H), 7.46 (t, 1H), 6.85 (s, 1H), 6.84 (s, 1H), 5.10 (m, 1H), 4.67-4.63 (m, 1H), 4.47-4.38 (m, 2H), 4.26-4.21 (m, 1H), 3.94 (s, 3H), 3.59 (dd, 1H), 3.41 (dd, 1H), 2.11-2.05 (m, 2H), 2.00 (s, 3H); Mass spectrum MH⁺ 461.

Example 28

(3*R*,5*S*)-5-[(4-[3-Chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]-1-(methoxyacetyl)pyrrolidin-3-ol

25 (Process (a))




The procedure described in Example 25 was repeated using (3*R*,5*S*)-5-[(4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]pyrrolidin-3-ol (209 mg) with methoxyacetic acid (42 μ l) to give the title compound as a white solid in 91% yield; NMR spectrum (DMSO- d_6) 9.96 (s, 1H), 8.45 (s, 1H), 8.17-8.14 (m, 1H), 7.76-7.72 (m, 1H), 7.45 (t, 1H), 6.83 (s, 2H), 5.10 (m, 1H), 4.73-4.69 (m, 1H), 4.47-4.38 (m, 2H), 4.25 (dd, 1H), 4.10-3.98 (m, 2H), 3.94 (s, 3H), 3.51 (dd, 1H), 3.40 (s, 1H), 3.29 (s, 3H), 2.05-2.02 (m, 2H); Mass spectrum MH^+ 491.

10 Example 29

(3*R*,5*S*)-5-[(4-[3-Chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)-methyl]-1-[(4-methylpiperazin-1-yl)acetyl]pyrrolidin-3-ol


(Process (a))



15 The procedure described in Example 25 was repeated using (3*R*,5*S*)-5-[(4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]pyrrolidin-3-ol (210 mg) with *N*-methylpiperazin-1-ylacetic acid (84 mg). Purification by column chromatography, using 0 to 10% 7*N* ammonia in methanol in DCM, gave the title compound as a white solid in 72% yield; NMR spectrum (DMSO- d_6) 9.91 (s, 1H), 8.45 (s, 1H), 8.14 (dd, 1H), 7.75-7.70 (m, 1H), 7.45 (t, 1H), 6.82 (s, 2H), 5.05 (m, 1H), 4.68-4.64 (m, 1H), 4.43-4.26 (m, 3H), 3.93 (s, 3H), 3.56 (m, 2H), 3.18 (d, 1H), 3.00 (d, 1H), 2.38 (br, 4H), 2.22 (br, 4H), 2.13 (s, 3H), 2.07-2.05 (m, 2H); Mass spectrum MH^+ 559.

Example 30**(3*S*,5*S*)-5-[(4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]-1-glycoloylpyrrolidin-3-ol**

(Process (a))



The procedure described in Example 20 was repeated using (3*S*,5*S*)-5-[(4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]pyrrolidin-3-ol (157 mg) with glycolic acid (31 mg) to give the title compound as a white solid in 81% yield; NMR spectrum (DMSO-*d*₆) 10.01 (s, 1H), 8.48 (s, 1H), 8.22 (dd, 1H), 7.74-7.69 (m, 1H), 7.43 (t, 1H), 6.84 (d, 1H), 6.78 (d, 1H), 5.29-5.28 (m, 1H), 4.73-4.68 (m, 1H), 4.64-4.59 (m, 2H), 4.44-4.42 (m, 1H), 4.34-4.31 (m, 1H), 4.06-4.05 (m, 2H), 3.93 (s, 3H), 3.64 (dd, 1H), 3.45 (s, 1H), 2.25-2.12 (m, 1H), 1.99-1.95 (m, 1H); Mass spectrum MH⁺ 477.

The (3*S*,5*S*)-5-[(4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]pyrrolidin-3-ol used as starting material was prepared as follows:

15 Di-*tert*-butylazodicarboxylate (DTAD, 1.38 g) was added in one portion to a mixture of 4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-ol (1.28 g), *tert*-butyl (2*S*,4*S*)-4-[[*tert*-butyl(dimethyl)silyl]oxy]-2-(hydroxymethyl)-pyrrolidine-1-carboxylate (1.66 g), and triphenylphosphine (1.58 g) in DCM (100 ml) at room temperature, under an atmosphere of nitrogen. The reaction mixture was stirred at room temperature for 28 hours and then

20 concentrated *in vacuo* to leave a brown oil. Purification by column chromatography, using 0-50% ethyl acetate in hexane as eluent, gave *tert*-butyl (2*S*,4*S*)-4-[[*tert*-butyl(dimethyl)silyl]oxy]-2-[(4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]-pyrrolidine-1-carboxylate as a white solid (1.13 g, 45%); Mass spectrum MH⁺ 633.

tert-Butyl (2*S*,4*S*)-4-[[*tert*-butyl(dimethyl)silyl]oxy]-2-[(4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]pyrrolidine-1-carboxylate (1.13 g) was dissolved in trifluoroacetic acid (50 ml) and stirred at room temperature for 72 hours. The reaction mixture was then concentrated *in vacuo*, and excess water was added, followed by the careful addition of saturated aqueous sodium hydrogen carbonate. The mixture was extracted with 3% methanol in DCM and the organic layer was separated, dried (Na₂SO₄) and

30 concentrated *in vacuo* to leave a pink solid. Trituration with cold acetonitrile gave (3*S*,5*S*)-5-


[[{4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl}oxy)methyl]pyrrolidin-3-ol as a brown solid (380 mg, 51%); NMR spectrum (DMSO-d₆) 10.38-10.54 (br s, 1H), 8.52 (s, 1H), 8.36 (dd, 1H), 7.94-7.99 (m, 1H), 7.43 (t, 1H), 6.84 (d, 1H), 6.79 (d, 1H), 4.75 (d, 1H), 4.41 (dd, 1H), 4.26-4.34 (m, 1H), 4.17 (t, 1H), 3.94 (s, 3H), 3.54-3.73 (m, 1H), 2.96 (dd, 1H), 2.79 (dd, 1H), 2.11-2.20 (m, 1H), 1.41-1.49 (m, 1H); Mass spectrum MH⁺ 419.

The *tert*-butyl (2*S*,4*S*)-4-[[*tert*-butyl(dimethyl)silyl]oxy]-2-(hydroxymethyl)-pyrrolidine-1-carboxylate used as starting material was prepared in an analogous method to *tert*-butyl (2*R*,4*S*)-4-[[*tert*-butyl(dimethyl)silyl]oxy]-2-(hydroxymethyl)-pyrrolidine-1-carboxylate described in Example 20, preparation of starting materials, starting from 1-*tert*-butyl 2-methyl (2*S*,4*S*)-4-hydroxypyrrolidine-1,2-dicarboxylate (commercially available or can be prepared using an analogous process to that described in Example 20 for the preparation of 1-*tert*-butyl 2-methyl (2*R*,4*S*)-4-hydroxypyrrolidine-1,2-dicarboxylate but using (4*S*)-1-(*tert*-butoxycarbonyl)-4-hydroxy-L-proline.

15 Example 31

(3*S*,5*S*)-5-[(4-[3-Chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl}oxy)methyl]-1-(*N,N*-dimethylglycyl)pyrrolidin-3-ol

(Process (a))




20 The procedure described in Example 30 was repeated using (3*S*,5*S*)-5-[(4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl}oxy)methyl]pyrrolidin-3-ol (157 mg) with *N,N*-dimethylglycine (43 mg) to give the title compound as a white solid in 83% yield; NMR spectrum (DMSO-d₆) 10.03 (s, 1H), 8.46 (s, 1H), 8.19 (dd, 1H), 7.76-7.71 (m, 1H), 7.43 (t, 1H), 6.83 (d, 1H), 6.75 (d, 1H), 5.27-5.26 (m, 1H), 4.72-4.67 (m, 1H), 4.61-4.56 (m, 1H), 4.41-4.40 (m, 1H), 4.34-4.29 (m, 1H), 3.93 (s, 3H), 3.78 (dd, 1H), 3.52 (br d, 1H), 3.14 (d, 1H), 2.99 (d, 1H), 2.26-2.17 (m, 7H), 1.95-1.90 (m, 1H); Mass spectrum MH⁺ 504.

Example 32

(3*R*,5*R*)-5-[(4-[3-Chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl}oxy)methyl]-1-

30 glycoloylpyrrolidin-3-ol

(Process (a))



The procedure described in Example 20 was repeated using (3*R*,5*R*)-5-[(4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]pyrrolidin-3-ol (209 mg) with
 5 glycolic acid (42 mg) to give the title compound as a white solid in 46% yield; NMR spectrum (DMSO-*d*₆) 10.01 (s, 1H), 8.48 (s, 1H), 8.22 (dd, 1H), 7.74-7.69 (m, 1H), 7.43 (t, 1H), 6.84 (d, 1H), 6.78 (d, 1H), 5.29-5.28 (m, 1H), 4.73-4.68 (m, 1H), 4.64-4.59 (m, 2H), 4.44-4.42 (m, 1H), 4.34-4.31 (m, 1H), 4.06-4.05 (m, 2H), 3.93 (s, 3H), 3.64 (dd, 1H), 3.45 (s, 1H), 2.25-2.12 (m, 1H), 1.99-1.95 (m, 1H); Mass spectrum MH⁺ 477.

10 The (3*R*,5*R*)-5-[(4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)-oxy)methyl]pyrrolidin-3-ol used as starting material was prepared as follows:

Di-*tert*-butylazodicarboxylate (DTAD, 914 mg) was added in one portion to a mixture of 4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-ol (848 mg), *tert*-butyl (2*R*,4*R*)-4-
 15 {*tert*-butyl(dimethyl)silyl}oxy}-2-(hydroxymethyl)-pyrrolidine-1-carboxylate (1.10 g), and triphenylphosphine (1.04 g) in DCM (100 ml) at room temperature, under an atmosphere of nitrogen. The reaction mixture was stirred at room temperature for 24 hours and then concentrated *in vacuo* to leave a brown oil. Purification by column chromatography, using 0-
 50% ethyl acetate in hexane as eluent, gave *tert*-butyl (2*R*,4*R*)-4-{*tert*-butyl(dimethyl)silyl}-oxy}-2-[(4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]-pyrrolidine-
 20 1-carboxylate as a white solid (839 mg, 50%); Mass spectrum MH⁺ 633.

tert-Butyl (2*R*,4*R*)-4-{*tert*-butyl(dimethyl)silyl}oxy}-2-[(4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]pyrrolidine-1-carboxylate (839 mg) was dissolved in trifluoroacetic acid (50 ml) and stirred at room temperature for 72 hours. The reaction mixture was then concentrated *in vacuo*, and excess water was added, followed by the
 25 careful addition of saturated aqueous sodium hydrogen carbonate. The mixture was extracted with 3% methanol in DCM and the organic layer was separated, dried (Na₂SO₄) and concentrated *in vacuo* to leave a green solid. Trituration with cold acetonitrile gave (3*R*,5*R*)-5-[(4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]pyrrolidin-3-ol as a
 30 grey solid (540 mg, 98%); NMR spectrum (DMSO-*d*₆) 10.54-10.02 (br s, 1H), 8.48 (s, 1H), 8.32 (dd, 1H), 7.93-7.88 (m, 1H), 7.38 (t, 1H), 6.79 (d, 1H), 6.69 (d, 1H), 5.05-4.74 (br s, 1H),

4.40-4.32 (m, 2H), 4.16 (t, 1H), 3.91 (s, 3H), 3.80-3.40 (br m, 2H), 2.98 (dd, 1H), 2.86 (dd, 1H), 2.22-2.14 (m, 1H), 1.52-1.44 (m, 1H); Mass spectrum MH^+ 419.


The *tert*-butyl (2*R*,4*R*)-4-[[*tert*-butyl(dimethyl)silyl]oxy]-2-(hydroxymethyl)-pyrrolidine-1-carboxylate used as starting material was prepared in an analogous method to
 5 *tert*-butyl (2*R*,4*S*)-4-[[*tert*-butyl(dimethyl)silyl]oxy]-2-(hydroxymethyl)-pyrrolidine-1-carboxylate described in Example 20, preparation of starting materials, starting from (4*R*)-1-(*tert*-butoxycarbonyl)-4-hydroxy-D-proline). This gave *tert*-butyl (2*R*,4*R*)-4-[[*tert*-butyl(dimethyl)silyl]oxy]-2-(hydroxy-methyl)-pyrrolidine-1-carboxylate as a clear oil in 39%
 over three steps; NMR spectrum ($CDCl_3$) 4.57 (d, 1H), 4.32-4.25 (m, 1H), 4.10-4.00 (m, 1H),
 10 3.83-3.51 (m, 3H), 3.39-3.23 (m, 1H), 2.26-2.18 (m, 1H), 1.48 (s, 9H), 0.97-0.90 (m, 10H), 0.09 (s, 6H).

Example 33

(3*R*,5*R*)-5-[(4-[3-Chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)-methyl]-1-

15 (*N,N*-dimethylglycyl)pyrrolidin-3-ol


(Process (a))



The procedure described in Example 32 was repeated using (3*R*,5*R*)-5-[(4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]pyrrolidin-3-ol (209 mg) with *N,N*-
 20 dimethylglycine (57 mg) to give the title compound as a white solid in 72% yield; NMR spectrum ($DMSO-d_6$) 10.03 (s, 1H), 8.46 (s, 1H), 8.19 (dd, 1H), 7.76-7.71 (m, 1H), 7.43 (t, 1H), 6.83 (d, 1H), 6.75 (d, 1H), 5.27-5.26 (m, 1H), 4.72-4.67 (m, 1H), 4.61-4.56 (m, 1H), 4.41-4.40 (m, 1H), 4.34-4.29 (m, 1H), 3.93 (s, 3H), 3.78 (dd, 1H), 3.52 (br d, 1H), 3.14 (d, 1H), 2.99 (d, 1H), 2.26-2.17 (m, 7H), 1.95-1.90 (m, 1H); Mass spectrum MH^+ 504.

CLAIMS

1. A quinazoline derivative of the formula I:




I

5

wherein:

- X^1 is $C(R^3)_2$, wherein each R^3 , which may be the same or different, is selected from hydrogen and (1-4C)alkyl;
- Q^1 is pyrrolidinyl;
- 10 and wherein Q^1 is linked to the group X^1-O by a ring carbon atom,
and wherein Q^1 is substituted at the 1-position by the group of the formula $ZC(O)$;
- Z is selected from (1-4C)alkyl, halogeno-(1-4C)alkyl, hydroxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, amino-(1-4C)alkyl, (1-4C)alkylamino-(1-4C)alkyl and di-[(1-4C)alkyl]amino-(1-4C)alkyl,
- 15 or Z is Q^2 , wherein Q^2 is heterocyclyl or heterocyclyl-(1-4C)alkyl,
and wherein any CH_2 or CH_3 group within a Z group, other than a CH_2 group within a heterocyclyl ring, optionally bears on each said CH_2 or CH_3 group one or more halogeno or (1-6C)alkyl substituents or a substituent selected from hydroxy, cyano, amino, carboxy, carbamoyl, formyl, mercapto, sulfamoyl, (2-6C)alkenyl, (2-6C)alkynyl, (1-6C)alkoxy,
- 20 (1-6C)alkylthio, (1-6C)alkylsulfinyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, \underline{N} -(1-6C)alkylcarbamoyl, $\underline{N,N}$ -di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, \underline{N} -(1-6C)alkyl-(2-6C)alkanoylamino, \underline{N} -(1-6C)alkylsulfamoyl, $\underline{N,N}$ -di-[(1-6C)alkyl]sulfamoyl, (1-6C)alkanesulfonylamino and
- 25 \underline{N} -(1-6C)alkyl-(1-6C)alkanesulfonylamino,
and wherein any heterocyclyl group within Q^1 or Z optionally bears one or more substituents, which may be the same or different selected from halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, sulfamoyl, formyl, mercapto, (1-6C)alkyl,

(2-6C)alkenyl, (2-6C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulfinyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, 5 N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulfamoyl, N,N-di-[(1-6C)alkyl]sulfamoyl, (1-6C)alkanesulfonylamino, and N-(1-6C)alkyl-(1-6C)alkanesulfonylamino, or from a group of the formula:




wherein X^2 is a direct bond or is selected from O, CO and $N(R^5)$, wherein R^5 is 10 hydrogen or (1-6C)alkyl, and R^4 is halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl, carboxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl, cyano-(1-6C)alkyl, amino-(1-6C)alkyl, N-(1-6C)alkylamino-(1-6C)alkyl, N,N-di-[(1-6C)alkyl]amino-(1-6C)alkyl, (2-6C)alkanoylamino-(1-6C)alkyl, (1-6C)alkoxycarbonylamino-(1-6C)alkyl, carbamoyl-(1-6C)alkyl, N-(1-6C)alkylcarbamoyl-(1-6C)alkyl, 15 N,N-di-[(1-6C)alkyl]carbamoyl-(1-6C)alkyl, (2-6C)alkanoyl-(1-6C)alkyl, (2-6C)alkanoyloxy-(1-6C)alkyl and (1-6C)alkoxycarbonyl-(1-6C)alkyl,

and wherein any heterocyclyl group within Q^1 or Z optionally bears 1 or 2 oxo or thioxo substituents;

m is 0, 1 or 2 and R^1 when present is located at the 6- and/or 7- position;

20 **each R^1 group**, which may be the same or different, is selected from hydroxy, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, or from a group of the formula :




wherein X^3 is a direct bond or is O, and Q^3 is (3-7C)cycloalkyl, (3-7C)cycloalkyl-(1-6C)alkyl, (3-7C)cycloalkenyl, (3-7C)cycloalkenyl-(1-6C)alkyl, heterocyclyl or heterocyclyl-(1-6C)alkyl, 25 or $(R^1)_m$ is (1-3C)alkylenedioxy,

and wherein adjacent carbon atoms in any (2-6C)alkylene chain within a R^1 substituent are optionally separated by the insertion into the chain of a group selected from O, S, SO, SO_2 , $N(R^6)$, CO, $CH(OR^6)$, $CON(R^6)$, $N(R^6)CO$, $SO_2N(R^6)$, $N(R^6)SO_2$, $CH=CH$ and $C\equiv C$ wherein R^6 is hydrogen or (1-6C)alkyl,

30 and wherein any $CH_2=CH-$ or $HC\equiv C-$ group within a R^1 substituent optionally bears at the terminal $CH_2=$ or $HC\equiv$ position a substituent selected from halogeno, carboxy, carbamoyl, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl,

amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl and di-[(1-6C)alkyl]amino-(1-6C)alkyl or from a group of the formula :




wherein X^4 is a direct bond or is selected from CO and $N(R^7)CO$, wherein R^7 is hydrogen or
 5 (1-6C)alkyl, and Q^4 is heterocyclyl or heterocyclyl-(1-6C)alkyl,

and wherein any CH_2 or CH_3 group within a R^1 substituent, other than a CH_2 group within a heterocyclyl ring, optionally bears on each said CH_2 or CH_3 group one or more halogeno or (1-6C)alkyl substituents or a substituent selected from hydroxy, cyano, amino, carboxy, carbamoyl, sulfamoyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulfinyl,
 10 (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulfamoyl, N,N-di-[(1-6C)alkyl]sulfamoyl, (1-6C)alkanesulfonylamino and N-(1-6C)alkyl-(1-6C)alkanesulfonylamino, or from a group of the formula:



wherein X^5 is a direct bond or is selected from O, S, SO, SO_2 , $N(R^8)$, CO, $CH(OR^8)$, $CON(R^8)$, $N(R^8)CO$, $SO_2N(R^8)$, $N(R^8)SO_2$, $C(R^8)_2O$, $C(R^8)_2S$ and $C(R^8)_2N(R^8)$, wherein R^8 is hydrogen or (1-6C)alkyl, and Q^5 is (3-7C)cycloalkyl, (3-7C)cycloalkyl-
 (1-6C)alkyl, (3-7C)cycloalkenyl, (3-7C)cycloalkenyl-(1-6C)alkyl, heterocyclyl or
 20 heterocyclyl-(1-6C)alkyl,

and wherein any heterocyclyl group within a substituent on R^1 optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, formyl, mercapto, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy,
 25 (1-6C)alkylthio, (1-6C)alkylsulfinyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulfamoyl, N,N-di-[(1-6C)alkyl]sulfamoyl, (1-6C)alkanesulfonylamino, and
 30 N-(1-6C)alkyl-(1-6C)alkanesulfonylamino, or from a group of the formula:



wherein X^6 is a direct bond or is selected from O, $N(R^{10})$ and C(O), wherein R^{10} is hydrogen or (1-6C)alkyl, and R^9 is halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl, carboxy-(1-6C)alkyl,

- (1-6C)alkoxy-(1-6C)alkyl, cyano-(1-6C)alkyl, amino-(1-6C)alkyl,
 (1-6C)alkylamino-(1-6C)alkyl, di-[(1-6C)alkyl]amino-(1-6C)alkyl,
 (2-6C)alkanoylamino-(1-6C)alkyl, (1-6C)alkoxycarbonylamino-(1-6C)alkyl,
 carbamoyl-(1-6C)alkyl, N-(1-6C)alkylcarbamoyl-(1-6C)alkyl,
 5 N,N-di-[(1-6C)alkyl]carbamoyl-(1-6C)alkyl, (2-6C)alkanoyl-(1-6C)alkyl or
 (1-6C)alkoxycarbonyl-(1-6C)alkyl,

and wherein any heterocyclyl group within a substituent on R¹ optionally bears 1 or 2
 oxo or thioxo substituents;

Y is selected from halogeno, cyano, trifluoromethyl, (1-6C)alkyl, (2-6C)alkenyl,

- 10 (2-6C)alkynyl and (1-6C)alkoxy;

- each R², which may be the same or different, is selected from halogeno, cyano, nitro,
 hydroxy, amino, carboxy, carbamoyl, sulfamoyl, trifluoromethyl, (1-6C)alkyl, (2-8C)alkenyl,
 (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylthio,
 (1-6C)alkylsulfinyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino,
 15 (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl,
N-(1-6C)alkylsulfamoyl, N,N-di-[(1-6C)alkyl]sulfamoyl; and

a is 0, 1, 2, 3 or 4;

or a pharmaceutically acceptable salt thereof.


- 20 2. A quinazoline derivative of the formula I according to claim 1, or a pharmaceutically
 acceptable salt thereof, wherein each of X¹, Q¹, R¹, R², a, m and Y are as defined in claim 1;
 and

- Z is selected from halogeno-(1-4C)alkyl, hydroxy-(1-4C)alkyl, (1-
 4C)alkoxy-(1-4C)alkyl, amino-(1-4C)alkyl, (1-4C)alkylamino-(1-4C)alkyl and
 25 di-[(1-4C)alkyl]amino-(1-4C)alkyl,

or Z is Q², wherein Q² is heterocyclyl or heterocyclyl-(1-4C)alkyl, and wherein the
 heterocyclyl in Q² contains at least 1 nitrogen heteroatom, and optionally 1 or 2 further
 heteroatoms selected from oxygen, nitrogen and sulfur,

- and wherein any CH₂ or CH₃ group within a Z group, other than a CH₂ group within a
 30 heterocyclyl ring, optionally bears on each said CH₂ or CH₃ group one or more halogeno or
 (1-6C)alkyl substituents or a substituent selected from hydroxy, cyano, amino, carboxy,
 carbamoyl, formyl, mercapto, sulfamoyl, (2-6C)alkenyl, (2-6C)alkynyl, (1-6C)alkoxy,


- (1-6C)alkylthio, (1-6C)alkylsulfinyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulfamoyl,
- 5 N,N-di-[(1-6C)alkyl]sulfamoyl, (1-6C)alkanesulfonylamino and N-(1-6C)alkyl-(1-6C)alkanesulfonylamino,
- and wherein any heterocyclyl group within Q¹ or Z optionally bears one or more (for example 1, 2 or 3) substituents, which may be the same or different selected from halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, sulfamoyl, formyl,
- 10 mercapto, (1-6C)alkyl, (2-6C)alkenyl, (2-6C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulfinyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulfamoyl,
- 15 N,N-di-[(1-6C)alkyl]sulfamoyl, (1-6C)alkanesulfonylamino, and N-(1-6C)alkyl-(1-6C)alkanesulfonylamino, or from a group of the formula:



- wherein X² is a direct bond or is selected from O, CO and N(R⁵), wherein R⁵ is hydrogen or (1-6C)alkyl, and R⁴ is halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl, carboxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl, cyano-(1-6C)alkyl, amino-(1-6C)alkyl, N-(1-6C)alkylamino-(1-6C)alkyl, N,N-di-[(1-6C)alkyl]amino-(1-6C)alkyl, (2-6C)alkanoylamino-(1-6C)alkyl, (1-6C)alkoxycarbonylamino-(1-6C)alkyl, carbamoyl-(1-6C)alkyl, N-(1-6C)alkylcarbamoyl-(1-6C)alkyl, N,N-di-[(1-6C)alkyl]carbamoyl-(1-6C)alkyl, (2-6C)alkanoyl-(1-6C)alkyl,
- 20 (2-6C)alkanoyloxy-(1-6C)alkyl and (1-6C)alkoxycarbonyl-(1-6C)alkyl,
- and wherein any heterocyclyl group within Q¹ or Z optionally bears 1 or 2 oxo or thioxo substituents.

3. A quinazoline derivative of the formula I according to either claim 1 or claim 2, or a
- 30 pharmaceutically acceptable salt thereof, wherein Q¹ is pyrrolidin-2-yl substituted at the 1-position by the group of the formula ZC(O).

4. A quinazoline derivative of the formula I according to any one of the preceding claims, or a pharmaceutically acceptable salt thereof, wherein Z is selected from hydroxy-(1-2C)alkyl, (1-4C)alkoxy-(1-2C)alkyl, amino-(1-2C)alkyl, (1-4C)alkylamino-(1-2C)alkyl and di-[(1-4C)alkyl]amino-(1-2C)alkyl,
- 5 or Z is Q², wherein Q² is heterocyclyl or heterocyclyl-(1-2C)alkyl, and wherein the heterocyclyl in Q² is a fully saturated 4, 5, 6 or 7 membered monocyclic heterocyclyl group which contains at least 1 nitrogen heteroatom, and optionally 1 or 2 further heteroatoms selected from oxygen, nitrogen and sulfur,
- and wherein Q² is linked to the (1-2C)alkyl or C(O) group by a ring nitrogen,
- 10 and wherein any CH₂ or CH₃ group within a Z group, other than a CH₂ group within a heterocyclyl ring, optionally bears on each said CH₂ or CH₃ group one or more (1-4C)alkyl substituents or a substituent selected from cyano, carbamoyl, formyl, (2-6C)alkenyl, (2-6C)alkynyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl and (2-6C)alkanoyl,
- and wherein any CH₂ which is attached to 2 carbon atoms or a CH₃ group which is
- 15 attached to a carbon atom within a Z group, other than a CH₂ group within a heterocyclyl ring, optionally bears on each said CH₂ or CH₃ group one or more halogeno substituents or a substituent selected from hydroxy, amino, mercapto, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulfinyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (2-6C)alkanoyloxy, (2-6C)alkanoylamino and N-(1-6C)alkyl-(2-6C)alkanoylamino,
- 20 and wherein any heterocyclyl group within Z optionally bears one or more (for example 1, 2 or 3) substituents, which may be the same or different selected from halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carbamoyl, formyl, (1-6C)alkyl, (2-6C)alkenyl, (2-6C)alkynyl, (1-6C)alkoxy, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl,
- 25 (2-6C)alkanoyl, N-(1-6C)alkylsulfamoyl and N,N-di-[(1-6C)alkyl]sulfamoyl, or from a group of the formula:



- wherein X² is a direct bond or is selected from O, CO and N(R⁵), wherein R⁵ is hydrogen or (1-4C)alkyl, and R⁴ is halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl,
- 30 (1-6C)alkoxy-(1-6C)alkyl, cyano-(1-6C)alkyl, amino-(1-6C)alkyl, N-(1-6C)alkylamino-(1-6C)alkyl, N,N-di-[(1-6C)alkyl]amino-(1-6C)alkyl, carbamoyl-(1-6C)alkyl, N-(1-6C)alkylcarbamoyl-(1-6C)alkyl and N,N-di-[(1-6C)alkyl]carbamoyl-(1-6C)alkyl,


and wherein any heterocyclyl group within Z optionally bears 1 or 2 oxo substituents.

5. A quinazoline derivative of the formula I according to any one of claims 1 to 3, or a pharmaceutically acceptable salt thereof, wherein Z is selected from (1-2C)alkyl, 5 hydroxy-(1-2C)alkyl, amino-(1-2C)alkyl, (1-4C)alkylamino-(1-2C)alkyl and di-[(1-4C)alkyl]amino-(1-2C)alkyl, or Z is Q², wherein Q² is heterocyclyl-(1-2C)alkyl, and wherein the heterocyclyl in Q² is selected from pyrrolidin-1-yl, piperidino and piperazin-1-yl, and wherein Q² is linked to the (1-2C)alkyl group by a ring nitrogen, 10 and wherein any heterocyclyl group within Z optionally bears one or more (for example 1, 2 or 3) substituents, which may be the same or different selected from fluoro, chloro, hydroxy, (1-3C)alkyl, (1-3C)alkoxy, or from a group of the formula:
- $$-X^2-R^4$$
- wherein X² is a direct bond or is O, and R⁴ is halogeno-(1-3C)alkyl, 15 hydroxy-(1-3C)alkyl or (1-3C)alkoxy-(1-3C)alkyl, and wherein any heterocyclyl group within Z optionally bears an oxo substituent.

6. A quinazoline derivative of the formula I according to claim 1, or a pharmaceutically acceptable salt thereof, wherein Q¹-X¹ is selected from (2R)-pyrrolidin-2-ylmethyl and (2S)- 20 pyrrolidin-2-ylmethyl, and wherein the group of the formula ZC(O) is attached at the 1-position of the pyrrolidin-2-yl group, Z is selected from methyl, hydroxymethyl, aminomethyl, (1-2C)alkylaminomethyl, di-[(1-2C)alkyl]aminomethyl and pyrrolidin-1-ylmethyl, 25 and wherein any heterocyclyl group within Z or Q¹ optionally bears one or more (for example 1, 2 or 3) substituents, which may be the same or different selected from hydroxy, (1-3C)alkyl and (1-3C)alkoxy, and wherein any heterocyclyl group within Z optionally bears an oxo substituent.

7. A quinazoline derivative of the formula I according to any one of the preceding claims, 30 or a pharmaceutically acceptable salt thereof, wherein m is 1 and R¹ is located at the 7-position.

8. A quinazoline derivative of the formula I according to claim 7, or a pharmaceutically acceptable salt thereof, wherein R^1 is selected from (1-4C)alkoxy, hydroxy-(1-4C)alkoxy and (1-4C)alkoxy(1-4C)alkoxy.
- 5 9. A quinazoline derivative of the formula I according to any one of the preceding claims, or a pharmaceutically acceptable salt thereof, wherein the aniline at the 4-position in the quinazoline ring in formula I is selected from 3-chloro-4-fluoroanilino, 3-chloro-2-fluoroanilino, 2-fluoro-5-chloroanilino, 3-bromoanilino, 3-methylanilino and 3-ethynylanilino.
- 10 10. A quinazoline derivative of the formula I according to claim 1, or a pharmaceutically acceptable salt thereof, wherein:
- m is 1 and R^1 is located at the 7-position and is selected from (1-4C)alkoxy, hydroxy-(1-4C)alkoxy, methoxy-(1-4C)alkoxy and ethoxy-(1-4C)alkoxy;
- Q^1-X^1 is pyrrolidin-2-ylmethyl,
- 15 and wherein Q^1 optionally bears one or more substituents, which may be the same or different selected from hydroxy and (1-3C)alkoxy,
- and wherein Q^1 optionally bears an oxo substituent,
- and wherein the group of the formula $ZC(O)$ is attached at the 1-position in Q^1 ;
- Z is selected from hydroxymethyl, aminomethyl, methylaminomethyl,
- 20 di-methylaminomethyl and pyrrolidin-1-ylmethyl,
- and wherein any heterocyclyl group within Z optionally bears 1 or 2 substituents, which may be the same or different selected from hydroxy, amino, methyl, ethyl, methoxy, methylamino and di-methylamino;
- Y is halogeno;
- 25 each R^2 , which may be the same or different, is selected from halogeno; and
- a is 0, 1 or 2;
- or a pharmaceutically acceptable salt thereof.
11. A quinazoline derivative of the formula I according to claim 1 of the formula Ia:



Ia

wherein:

m is 0 or **m** is 1 and **R**^{1a} is selected from (1-4C)alkoxy, hydroxy-(1-4C)alkoxy and

5 (1-4C)alkoxy(1-4C)alkoxy;

R¹¹ is selected from hydrogen, hydroxy, (1-4C)alkyl, (1-4C)alkoxy, hydroxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, hydroxy-(1-4C)alkoxy and (1-4C)alkoxy-(1-4C)alkoxy;

Z¹ is selected from (1-2C)alkyl, hydroxy-(1-2C)alkyl, (1-2C)alkoxy-(1-2C)alkyl,
10 amino-(1-2C)alkyl, (1-4C)alkylamino-(1-2C)alkyl and di-[(1-4C)alkyl]amino-(1-2C)alkyl,
or **Z**¹ is **Q**², wherein **Q**² is pyrrolidinyl-(1-2C)alkyl,

and wherein any pyrrolidinyl group within **Z**¹ optionally bears one or more
substituents, which may be the same or different selected from fluoro, chloro, hydroxy,
(1-3C)alkyl, (1-3C)alkoxy, or from a group of the formula:

15 $-X^2-R^4$

wherein **X**² is a direct bond or is O, and **R**⁴ is halogeno-(1-3C)alkyl,
hydroxy-(1-3C)alkyl or (1-3C)alkoxy-(1-3C)alkyl,

and wherein any pyrrolidinyl group within **Z**¹ optionally bears an oxo substituent;

Y¹ is selected from fluoro, chloro, bromo, cyano, methyl, ethyl, isopropyl, methoxy,
20 ethoxy, vinyl, allyl, ethynyl, 1-propynyl and 2-propynyl;

each **R**^{2a}, which may be the same or different, is selected from fluoro, chloro, bromo,
cyano, hydroxy, trifluoromethyl, (1-4C)alkyl, (2-4C)alkynyl and (1-4C)alkoxy; and

a is 0, 1 or 2;

or a pharmaceutically acceptable salt thereof.

25

12. A quinazoline derivative of the formula Ia according to claim 11, or a
pharmaceutically acceptable salt thereof, wherein:

m is 1 and **R^{1a}** is selected from methoxy, ethoxy, 2-hydroxyethoxy, 2-methoxyethoxy and 2-ethoxyethoxy;

R¹¹ is selected from hydrogen, hydroxy or methoxy;

Z¹ is selected from methyl, hydroxymethyl, aminomethyl, methylaminomethyl,

5 dimethylaminomethyl and pyrrolidin-1-ylmethyl;

Y¹ is selected from fluoro, chloro, bromo and ethynyl;

R^{2a} is selected from fluoro, chloro and bromo; and


a is 0 or 1.

- 10 13. A quinazoline derivative of the formula Ia according to claim 1 selected from:
- 2-[(2*R*)-2-[(4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]pyrrolidin-1-yl]-2-oxoethanol;
- N*-(3-chloro-4-fluorophenyl)-5-[(2*R*)-1-[(dimethylamino)acetyl]pyrrolidin-2-yl]methoxy-7-methoxyquinazolin-4-amine;
- 15 *N*-(3-chloro-4-fluorophenyl)-7-methoxy-5-[(2*S*)-1-(pyrrolidin-1-ylacetyl)pyrrolidin-2-yl]methoxyquinazolin-4-amine;
- (3*S*,5*S*)-5-[(4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]-1-(*N,N*-dimethylglycyl)pyrrolidin-3-ol;
- (3*S*,5*R*)-5-[(4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]-1-
- 20 glycoloylpyrrolidin-3-ol;
- (3*R*,5*R*)-5-[(4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]-1-glycoloylpyrrolidin-3-ol;
- (3*R*,5*R*)-5-[(4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]-1-(*N,N*-dimethylglycyl)pyrrolidin-3-ol;
- 25 (3*S*,5*R*)-1-acetyl-5-[(4-[3-chloro-4-fluoroanilino]-7-methoxyquinazolin-5-yl)oxy)methyl]pyrrolidin-3-ol;
- 2-[(2*R*)-2-[(4-[3-chloro-4-fluoroanilino]-7-(2-hydroxyethoxy)quinazolin-5-yl)oxy)methyl]pyrrolidin-1-yl]-2-oxoethanol;
- 2-[(2*R*)-2-[(4-[3-chloro-4-fluoroanilino]-7-(2-ethoxyethoxy)quinazolin-5-
- 30 yl)oxy)methyl]pyrrolidin-1-yl]-2-oxoethanol;
- 2-[(2*R*)-2-[(4-[3-chloro-4-fluoroanilino]-7-ethoxyquinazolin-5-yl)oxy)methyl]pyrrolidin-1-yl]-2-oxoethanol; and

N-(3-chloro-4-fluorophenyl)-7-methoxy-5-[(2*R*)-1-(methoxyacetyl)pyrrolidin-2-yl]methoxy}quinazolin-4-amine;
or a pharmaceutically acceptable salt thereof.

14. A process for the preparation of a quinazoline derivative of the formula 1, or a pharmaceutically acceptable salt thereof, according to claim 1 which comprises:

Process (a) the coupling, conveniently in the presence of a suitable base, of a quinazoline of the formula **II**:




II

wherein R^1 , R^2 , X^1 , Y , Q^1 , a and m have any of the meanings defined in claim 1 except that any functional group is protected if necessary, with a carboxylic acid of the formula $Z\text{-COOH}$, or a reactive derivative thereof, wherein Z has any of the meanings defined in claim 1
except that any functional group is protected if necessary;

or

Process (b): for the preparation of those compounds of the formula **I** wherein Z is selected from amino-(1-4C)alkyl, (1-4C)alkylamino-(1-4C)alkyl, di-[(1-4C)alkyl]amino-(1-4C)alkyl, or Z is Q^2 , wherein Q^2 is heterocyclyl-(1-4C)alkyl, and wherein the heterocyclyl in Q^2 contains at least 1 nitrogen heteroatom, and optionally 1 or 2 further heteroatoms selected from oxygen, nitrogen and sulfur, and wherein Q^2 is attached by a ring nitrogen atom,

the reaction of a quinazoline of the formula **III**:



III


wherein R^1 , R^2 , X^1 , Y , Q^1 , a and m have any of the meanings defined in claim 1 except that any functional group is protected if necessary, Z^1 is a direct bond or (1-4C)alkyl and L^1 is a displaceable group, with an amine or a compound of the formula Q^2H ;

or


Process (c): for the preparation of those compounds of the formula I wherein R^1 is a hydroxy group, the cleavage of a quinazoline derivative of the formula I wherein R^1 is a (1-6C)alkoxy;

or

10 Process (d): for the preparation of those compounds of the formula I wherein m is 1 and R^1 is optionally substituted (1-6C)alkoxy, optionally substituted (2-6C)alkenyloxy, optionally substituted (2-6C)alkynyloxy or a group of the formula :



wherein X^3 is O, and Q^3 is as defined in claim 1 except that any functional group is protected if necessary, the reaction, conveniently in the presence of a suitable base, of a quinazoline of the formula IV:



IV

wherein R^2 , X^1 , Y , Q^1 , Z and a have any of the meanings defined in claim 1 except that any functional group is protected if necessary, with a compound of the formula $R^{1a}-L^1$ or Q^3-L^1 wherein Q^3 is as defined in claim 1 except any functional group is protected if necessary, R^{1a} is selected from optionally substituted (1-6C)alkyl, optionally substituted (2-6C)alkenyl and optionally substituted (2-6C)alkynyl and L^1 is a displaceable group;

or

Process (e): for the preparation of those compounds of the formula I wherein m is 1 and R¹ is optionally substituted (1-6C)alkoxy, optionally substituted (2-6C)alkenyloxy, optionally substituted (2-6C)alkynyloxy or a group of the formula :



wherein X³ is O, and Q³ is as defined in claim 1, the coupling of the a quinazoline of the formula IV as hereinbefore defined with an alcohol of the formula R^{1a}OH or Q³OH wherein Q³ is as defined in claim 1 except any functional group is protected if necessary and R^{1a} is as defined in process (d);

10 and thereafter, if necessary:

- (i) converting a quinazoline derivative of the formula I into another quinazoline derivative of the formula I;
- (ii) removing any protecting group that is present by conventional means;
- (iii) forming a pharmaceutically acceptable salt.

15

15. A pharmaceutical composition which comprises a quinazoline derivative of the formula I, or a pharmaceutically-acceptable thereof, as defined in claim 1 or claim 2 in association with a pharmaceutically-acceptable diluent or carrier.

20 16. A quinazoline derivative of the formula I, or a pharmaceutically acceptable salt thereof, as defined in claim 1 or claim 2 for use as a medicament.

17. The use of a quinazoline derivative of the formula I, or a pharmaceutically-acceptable salt thereof, as defined in claim 1 or claim 2 in the manufacture of a medicament for use in the
25 treatment of a disease state or medical condition mediated alone or in part by one or more erbB receptor tyrosine kinases.

18. The use of a quinazoline derivative of the formula I, or a pharmaceutically-acceptable salt thereof, as defined in claim 1 or claim 2 in the manufacture of a medicament for use in the
30 prevention or treatment of those tumours which are sensitive to inhibition of one or more of the erbB receptor tyrosine kinases, that are involved in the signal transduction steps which lead to the proliferation of tumour cells.

19. The use of a quinazoline derivative of the formula I, or a pharmaceutically-acceptable salt thereof, as defined hereinbefore in claim 1 or claim 2 in the manufacture of a medicament for use in providing a selective EGFR tyrosine kinase inhibitory effect.

5 20. A method for treating a disease or medical condition mediated alone or in part by one or more erbB receptor tyrosine kinases in a warm-blooded animal, such as man, in need of such treatment, which comprises administering to said animal an effective amount of a quinazoline derivative of the formula I, or a pharmaceutically-acceptable salt thereof, as defined in claim 1 or claim 2.

10

21. A method for the prevention or treatment of those tumours which are sensitive to inhibition of one or more of the erbB family of receptor tyrosine kinases, that are involved in the signal transduction steps which lead to the proliferation and/or survival of tumour cells in a warm-blooded animal, such as man, in need of such treatment, which comprises

15 administering to said animal an effective amount of a quinazoline derivative of the formula I, or a pharmaceutically-acceptable salt thereof, as defined in claim 1 or claim 2.

22. A method of treatment for providing a selective EGFR tyrosine kinase inhibitory effect in a warm-blooded animal, such as man, in need of such treatment, which comprises

20 administering to said animal an effective amount of a quinazoline derivative of the formula I, or a pharmaceutically-acceptable salt thereof, as defined hereinbefore.

INTERNATIONAL SEARCH REPORT

national Application No

PCT/GB2004/001799

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 A61K31/517 C07D403/12 C07D403/14 A61P35/00		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 7 C07D		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, WPI Data, PAJ, BEILSTEIN Data, CHEM ABS Data		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, Y	WO 03/040108 A (PASS MARTIN ; ASTRAZENECA UK LTD (GB); BRADBURY ROBERT HUGH (GB); HENN) 15 May 2003 (2003-05-15) see claims 1 and 24 and example 25.10 on page 155	1-22
Y	WO 96/09294 A (WELLCOME FOUND ; HUDSON ALAN THOMAS (GB); VILE SADIE (GB); BARRACLOUGH) 28 March 1996 (1996-03-28) cited in the application claims 1,26	1-22
Y	WO 01/94341 A (PLE PATRICK ; HENNEQUIN LAURENT FRANCOIS AND (FR); ASTRAZENECA UK LTD) 13 December 2001 (2001-12-13) cited in the application claims 1,16	1-22
----- -/-		
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : *A* document defining the general state of the art which is not considered to be of particular relevance *E* earlier document but published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. *Z* document member of the same patent family		
Date of the actual completion of the international search		Date of mailing of the international search report
4 August 2004		17/08/2004
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer Bérillon, L

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB2004/001799

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>WO 96/15118 A (THOMAS ANDREW PETER ; ZENECA LTD (GB); BROWN DEARG SUTHERLAND (GB); MO) 23 May 1996 (1996-05-23) cited in the application claims 1,12.</p>	1-22

INTERNATIONAL SEARCH REPORT

International application No.
PCT/GB2004/001799

Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 20-22 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.

☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB2004/001799

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 03040108	A	15-05-2003	CA 2465068 A1	15-05-2003
			CA 2465100 A1	15-05-2003
			WO 03040108 A1	15-05-2003
			WO 03040109 A2	15-05-2003
WO 9609294	A	28-03-1996	AU 3482495 A	09-04-1996
			EP 0782570 A1	09-07-1997
			WO 9609294 A1	28-03-1996
			JP 10505600 T	02-06-1998
			TR 960233 A2	21-06-1996
			ZA 9507853 A	18-03-1997
WO 0194341	A	13-12-2001	AU 6048201 A	17-12-2001
			BG 107332 A	31-07-2003
			BR 0111335 A	10-06-2003
			CA 2407371 A1	13-12-2001
			CN 1434821 T	06-08-2003
			CZ 20023961 A3	12-03-2003
			EE 200200673 A	15-06-2004
			EP 1292594 A1	19-03-2003
			WO 0194341 A1	13-12-2001
			HU 0301046 A2	28-08-2003
			JP 2003535859 T	02-12-2003
			NO 20025792 A	02-12-2002
			SK 17022002 A3	02-05-2003
WO 9615118	A	23-05-1996	AT 175962 T	15-02-1999
			AU 703328 B2	25-03-1999
			AU 3813095 A	06-06-1996
			CA 2200871 A1	23-05-1996
			DE 69507505 D1	04-03-1999
			DE 69507505 T2	02-06-1999
			DK 790986 T3	13-09-1999
			EP 0790986 A1	27-08-1997
			ES 2128092 T3	01-05-1999
			FI 971970 A	07-05-1997
			WO 9615118 A1	23-05-1996
			JP 10508616 T	25-08-1998
			NO 972152 A	12-05-1997
			NZ 294917 A	25-02-1999
			US 5821246 A	13-10-1998
			ZA 9509572 A	13-05-1996